INDEX OF SURGICAL PROGRESS.

BONES, JOINTS, ORTHOPÆDIC.

I. The Treatment of Advanced Conditions of Equinovarus. By R. SWAN (Dublin). When, says the author, the person afflicted with equino-varus has reached the age of ten years, either with early treatment and subsequent relapse, or without treatment, features are presented which preclude the hope of success being attended by tenotomy and extension. The tarsal bones are thickened, and the typical distortion is increased. The large adventitious bursa is dense, and includes in its base, the cuboid and the tarsal end of the fifth metatarsal bones, and may extend to a greater or less degree on the dorsum of the foot. In such case ablation of a portion of the tarsal bones is called for, provided an atrophied condition of the anterior part of the foot does not exist. The portion removed will consist of the anterior three-fourths of the cuboid bone, all the external cuneiform except its posterior part, the proximal extremities of the third, fourth and fifth metatarsal bones, and the apex of the wedge will consist of a portion of the middle cuneiform. Attention is drawn to the fact that after this operation the mobility of the foot will remain almost unimpaired. As the metatarsal bones have their epiphyses at the distal end, it is probable that the development of the foot would not be interfered with. The author had operated at the time of the reading of his paper, upon thirty-four selected cases since 1876.—*The Dublin Journal of Medical Science*, May, 1888.

H. PERCY DUNN (London).

II. Rare Fracture of Olecranon. By JORDAN LLOYD (Birmingham.) This was a specimen obtained by excising the elbow for an unreduced dislocation of 18 months standing. The olecranon was