Conclusion.

The present case has, then, much in common, clinically and pathologically, with more typical cases of acute red degeneration; it differs in that the infection proceeded to a well-established septicaemia, but as in puerperal sapraemia infection may remain localised to the uterus, so we suggest it may remain localised in necrobiosis with hæmolysis. We think that the theory of anaerobic infection as the primary cause of this condition is more satisfactory than the current theories of primary thrombosis or aseptic necrosis, and the immunity from red degeneration which ovarian fibroids show, as has been pointed out by Sir John Bland-Sutton,⁵ is well explained by this theory in virtue of their non-contact with the uterine cavity. A final point is the fact that although all varieties of uterine fibroids may become necrobiotic, interstitial and submucous tumours are most liable to become affected.

For permission to describe this case we wish to record our thanks to Mr. T. H. Kellock, surgeon to the Middlesex Hospital, and to Prof. James McIntosh, of the Bland-Sutton Institute of Pathology, Middlesex Hospital. We are indebted for the illustration to Mr. S. A. Sewell.

References.—1. Joannovics, G.: *Ztschr. f. Heilk.*, Wien u. Leipz., 1904 (25), p. 25. 2. Stadelmann, E.: *Ikterus*, Stuttgart, 1891. 3. Eppinger, H.: *Ergeb. d. inn. Med. u. Kinderh.*, 1908, i., p. 107. 4. Murray, H. Leith: *Jl. Obst. and Gynaecol.*, 1910, vol. xvii., p. 534. 5. Bland-Sutton, Sir J.: *B.M.J.*, 1909, i., 1471. 6. Bland-Sutton, Sir J.: *Tumours*, sixth edition, p. 224. 7. Smith, J. Lorrain, and Shaw, W. Fletcher: *Jl. Obst. and Gynaecol.*, 1909, vol. xv., p. 225. 8. Hartmann, H., and Mignot, R.: *Annales de Gynécologie*, 1896, vol. xiv., p. 425. 9. Lea, A. W. W., and Sidebotham, E. J.: *Jl. Obst. and Gynaecol.*, 1909, vol. xv., p. 26 (with bibliography).

CHRONIC DEEP INFECTION OF THE JAWS.

By STANLEY COLYER, M.D., M.R.C.P. LOND.,

RADIOLOGIST TO THE MILD MAY MISSION HOSPITAL AND
ASSISTANT RADIOLOGIST TO THE ROYAL DENTAL
HOSPITAL, LONDON.

It is well known to many clinical observers that the results obtained in general diseases by the radical treatment of oral sepsis, their assumed cause, are by no means uniform. Anyone who has followed the fortunes of this ætiological factor knows of brilliant results, complete failures, partial failures, apparent cures with subsequent relapses, and of advanced cases of pyorrhœa alveolaris with little or no demonstrable general symptoms. Like many others who believe in the importance of the septic mouth, I was at a loss to account for these obvious failures, and not until I took up the study of radiology did I obtain a workable explanation.

Type of Infection only Diagnosable from Radiograms.

The examination of several hundred dental radiograms made it quite clear to me that in many cases infection is by no means limited to the teeth and their immediate neighbourhood, but may pass into the very body of the jaws and there form breeding-grounds for germs which in time produce changes in the bone sufficiently coarse to alter the radiographic shadows peculiar to normal cancellous tissue. It was plain, also, that, even in edentulous parts of jaws from which septic teeth had been removed, infection of the bone might continue, so that it did not follow, when the offending teeth had been extracted, that nature was always able to deal with the residual infection. Further, in certain patients radiographed for their own satisfaction—patients declared clinically fit by their dentists—I frequently found unexpected infection, which showed that such conditions of the jaw may be undiagnosable save from a radiogram. There was still another class, those with advanced marginal osteitis, who showed no obvious general conditions; these usually presented markedly sclerotic edges to their receding alveolar edge, and either but little bone infection or infection that was practically cured.

It will thus be seen that the results of these findings was to give the workable explanation I spoke of above, and I may add that it seemed to support the view of those who believe that the damage done by oral sepsis is chiefly via the lymphatic rather than via the alimentary system. It gave me also an explanation of that somewhat rare condition of the jaws characterised by persistent shrinkage.

I do not wish to indicate that infection of the jaws has not been recognised, because it is clear that if the periodontal membrane is infected the bone immediately around must be infected; it is a pathological necessity. But it has been thought by most workers that this infection was limited to the bone immediately around the roots, and it does not seem to have been recognised, or if recognised its importance has been overlooked, that infection may pass deeply and there settle. Moreover, I do not wish in any way to minimise the importance of the more superficial infections, such as marginal osteitis, infected and thickened periodontal membranes, and peri-apical inflammatory changes—these have their importance and possibly greater importance than the deep infection, but I want in this paper, as far as possible, to confine my attention to the latter only, and after describing shortly its radiographic appearances to discuss its possible relationship to the failures attributed to oral sepsis.

Necessity of Minute Examination.

To discover infection of the jaw one must study closely the finer changes in the cancellous tissue. To do so it is often necessary to use a magnifying glass, and it is *always* necessary to view radiograms in a strong transmitted light. To hold up negatives at arm's length against the daylight and expect to see fine changes is to hope for the impossible. Careful examination shows that these changes vary considerably in appearance, but all are essentially dependent upon the two inflammatory changes of bone known as rarefaction and sclerosis—the varying appearances being due, it would seem, to the structure of the cancellous tissue, to the nature of the infection, and to the resistance and age of the patient.

Let us suppose that the radiograms have been suitably mounted close together and are now in position on a view-box. On glancing at them the radiologist gains a general impression—in other words, he knows at once without looking far for details that the patient is suffering either from localised areas of infection or that he has a more general infection of one or both jaws—has, in fact, a chronic pan-alveolitis of the jaws. It is just this general impression of infection which cannot be exactly translated into words; it depends upon accumulated experiences and upon the examination of numerous radiograms produced by a known technique. It is the familiarity of a radiologist with his own technique which makes him careful in expressing an opinion upon the work of others. With a full consciousness of the difficulties in mind we find that the general impression resolves itself into two smaller impressions: (1) a suggestion of woolliness or fleeciness about the negatives, and (2) the presence of blotches or patches of either increased or decreased translucency. If we now look more closely into the radiograms we shall note that the fleeciness appears to be due to a thickening of the trabeculae of the cancellous tissue and to a decreased translucency of the cancellous spaces, as though lime salts were diffused throughout the medullary substance. The blotches are due to (1) largish or smallish areas, or even streaks, of increased translucency which may be translated as areas of rarefaction, regular or irregular in outline; the bony tissue immediately around them may or may not be sclerosed, and the streaks referred to are seen in more severe cases of infection, generally in young people, where reparative changes are insignificant, and their direction is that of the lymphatic flow; (2) largish or smallish areas of sclerosis, brought about by irritation of toxins from some local source. The small areas may be nothing

but dots which may represent the thickened junction of cancellous trabeculae or a healed "small darkish spot." In severe cases the entire picture of the structure of cancellous tissue is altered; in less severe cases the structure is altered near obviously infected spots, such as a septic root.

If the negatives be now looked at still more closely, with the aid of a magnifying glass, one of the most important features of chronic infection may be discovered—namely, the presence of small, roundish, blackish spots surrounded by a layer of sclerosed bone. I can give these spots no name. From analogy with a similar condition often found at the apex of infected inter-radicular bony septa I assume them to be small masses of inflammatory tissue dependent upon a low grade infection. To be of diagnostic value they must be present in fair numbers, as a very similar appearance is shown by the minute foramina of nutrient vessels from the periosteum. These spots may occur singly or in clusters; the latter appearance is very characteristic. The proofs of their infective nature is indirect, and are these: (1) They may often be seen in the bone immediately around the apices of infected teeth. (2) They may sometimes be seen in the substances of interdental bony septa in a state of chronic inflammation from a marginal osteitis. (3) In an otherwise normal-looking septum they may sometimes be seen right up against an exostosis affecting what appears to be a healthy tooth—a most significant fact.

Clinical Importance of Bone Infections.

Let us now turn from a description of these bone infections and endeavour to assess their clinical importance. It is beyond argument that they, together with the more superficial oral infections, do exist in relation to certain other diseases which clear up on their removal. But there remains a large residuum of cases in which such treatment is of little or no avail. It is clear from what I have written that it is clinically impossible to gauge the degree of jaw infection; until a very large number of cases checked by radiographic examination have been investigated we are not justified in deciding against this ætiological factor in cases where treatment has presumably failed. And even here we are met with a difficulty in that jaw infection must have existed for some considerable time before radiographic changes become evident. In the early stages of deep jaw infection, then, radiology is useless; we must look to the bacteriologist in some way to fill this unfortunate gap. Of the cases suffering from general diseases supposed to be associated with oral sepsis which I have examined the majority have been over the age of 40—that is, over the age of commencing natural atrophy. That is one definite fact that has struck me, and the second fact that has struck me has been the extraordinary amount of infection that some of these patients must have harboured for years. Many of these patients stated that their general troubles were of recent date, so that for a long period they must have successfully put up a fight against the infection; and it is highly probable that others who do not come into the doctor's hands may ultimately be completely successful. Success would seem to be evident when sclerosis, in large or in small lonely masses, exists without the presence of rarefied areas or of the small dark spots. So long as these evidences of infection exist, however marked be the reparative changes, we must assume that they are kept up by organisms whose virulence, whatever it may originally have been, is now low; but, however low, they must still be regarded as living and capable of supplying the body with their toxins or even of themselves infecting the general circulation. That the mere removal of teeth and closure of the ports of entry of germs is not always sufficient to bring about a cure of the bone has been proved by the Americans. I have myself frequently noted the continued existence of deep infection in edentulous parts of jaws, but I have yet to satisfy myself that a completely edentulous jaw may retain infection for some years, though I have no doubt of its probable existence.

It follows from what I have said in this latter part of my paper that failures do not necessarily prove that oral sepsis was not causally related to the disease attributed to it, the failure being possibly due to the fact that the body is unable to rid itself of the deep jaw infection, or it may possibly be due to the fact that the disease cannot necessarily be cured by removing the cause. The latter reason may account for the frequent refusal of cases of arthritis to clear up. Since beginning the writing of this paper I have had the advantage of reading some records of American work on the subject. The reading has given me no reason to alter my views; it has, indeed, strengthened them. I will refer to one of these papers, by F. F. Molt, of Chicago, in which he deals with the subject of residual sepsis. He states that during the examination of over a thousand edentulous areas from which teeth, giving a history of previous pathological disturbance, had been removed by simple extraction, 41 per cent. showed evidence of residual infection—a very arresting statement. He also states that, on bacterial examination of the granulomata found so often at the apices of infected teeth, the majority show a mixed infection with streptococcus predominating, very few giving negative results. It is after the removal of such teeth areas of residual infection remain, and cultures made from hundreds of such areas show the same type of bacteria found in the granulomata themselves. Although I am not in a position to check Dr. Molt's statements I can see no reason for doubting them.

It is impossible to estimate the proportion of persons in this country who are suffering from jaw infection. One can safely say it is the vast majority, and it is the irony of fate that many who have paid particular attention to their teeth, having their dead teeth treated and saved, are often in the worst state. What is the result of this jaw infection? Does it in many cases remain innocuous, or does it in every case produce some systemic effect which in the minority only assumes so gross a form that it becomes clinically diagnosable? In "Injuries and Diseases of Bones and Joints," by Baetjer and Waters, both experienced radiologists, the following statement appears (p. 223): "Practically all of us when we reach the age of 45 show small exostoses in and around the joints, particularly where the ligaments attach, and yet there may be no clinical manifestation of an arthritic process." One may well ask, What causes these changes?

Conclusion.

Before closing this paper I wish to refer briefly to the possible means of dealing with jaw infection. In America an endeavour has been made to deal with the infected bone about the teeth by curettage, and for it they claim a reasonable amount of success. But it would seem from the evidence of radiograms to be impossible to curettage away satisfactorily the whole of the infected bone, so that much must be left behind untouched, and much that has been damaged—the latter tissue being in a state that may lead to further troubles. Besides this surgical procedure there are at least two other possibilities open after the removal of the ports of infection: (1) By means of vaccines, and (2) by means of ionisation. These are mere suggestions, but they would seem to be measures well worth testing, and if adopted their value can be ascertained only by comparing the results of a long series of cases checked by subsequent radiographic examinations to a similar series in which no treatment has been adopted save that of extraction.

THE LATE DR. W. GIBSON.—The death is recorded from Campbeltown, Argyll, of Dr. William Gibson, who died on Jan. 13th at the great age of 98, having lived during the reigns of five British sovereigns. Graduating M.D. at Edinburgh in 1854 and obtaining the L.R.C.S. Edin. in the same year, he afterwards held several public appointments in the district in which he practised, and retired only a little more than a year ago. Of the years of his life 65 had been devoted to public service.