a few retroperitoneal metastases. Riegel mentions¹⁴ a case in which a large sarcoma nodule was vomited and the diagnosis made. There were in the same case metastases in the skin, but the danger of depending on such co-existing tumors is made clear by von Leube,¹⁵ who found, in one case, sarcoma in the skin and an epithelial tumor in the stomach.

This symptom is probably as little reliable as two others mentioned by Schlesinger—enlargement of the spleen and swelling of the follicles of the tongue, both thought by Kundrat to be common in lymphosarcoma. They may sometimes enable one to correctly recognize gastric sarcoma in a late stage, but so far as the earlier stages are concerned we have no specific symptoms of sarcoma of the stomach, and even when tumor is present it can not be positively differentiated from cancer. It is important to discover the affection of the wall early, and if possible to carry out either radical or palliative operative treatment, without taking time to make a histological diagnosis from the clinical features. If discovered too late for this, the treatment must follow the same general lines as for other tumors of the stomach. The results of operations compare favorably with those for other malignant stomach diseases. In Cantwell's case, a spindle-celled sarcoma involving five inches square of the stomach wall was removed. The patient was free from symptoms for six months after the operation, and then a secondary tumor was discovered.

REFERENCES

1. Klinisches über Magentumoren nicht carcinomatöser Natur (Magensarkom). Schlesinger, Hermann. Festschrift Herrn Hofrath Prof. Dr. L. von Schroetter, Zeitschrift f. Klin. Med. Bd. xxxii, Supplement-Heft. 1897.
2. Westphalen, H.: Ein prim. Sarcom des Magens. St. Petersburger med. Woch., 1893, p. 403.
3. Krüger, Friedrich: Die prim. Bingedewebsgeschwülste des Magendarmkanals. Inaug. Dissert., Berlin, 1894.
4. Hartley: Sarcoma of Stomach; Annals of Surgery, 1896, p. 609.
5. Weiss: Myosarcom des Magens. Jahrb. der K. K. Wiener Krankenanstalten. Bd. iii, 1896.
6. Strauss, Aron: Sarcomatosis der Haut und des Magens. Inaug. Dissert., Würzburg, 1896.
7. Hayem and Lion: Traité de Méd. et de Thérapeutique. Tome iv, 1897.
8. Baldy, J. M.: Removal of Stomach for Sarcoma. THE JOURNAL 1898, 1, p. 523.
9. Brooks, H.: A Case of Primary Multiple Sarcoma of the Stomach, following Gun-shot Wound. Medical News. 1898, 1, p. 617.
10. Cappello, P. and P. E.: A proposito di un raro caso di miomasarcoma cistico dello stomaco. Bull. d. r. Accad. Med. di Roma. 1898-9, xxiv, 321-342.
11. Cantwell, F. V.: Sarcoma of the Stomach. Annals of Surgery, Nov., 1899.
12. Finlayson, Jas.: Case of Sarcoma of the Stomach in a Child aged 3½ years. Brit. Med. Jour., 1899, 2, p. 1535.
13. Welsch, Carl: Ueber Sarkom des Magens. In. Dissert., München, 1899.
14. Riegel, F.: Die Erkrankungen des Magens. ii. Theil. p. 847. Nothnagel's Specielle Path. u. Ther. Bd. xvi, Th. 2.
15. Von Leube, W.: Spec. Diag. der inn. Krankheiten., Bd. i.

VALUE OF THE ANGIOTRIBE IN VAGINAL CELIOTOMY.*

J. H. CARSTENS, M.D.

DETROIT, MICH.

When Battey first suggested the removal of the normal ovaries in cases of otherwise incurable dysmenorrhea, he performed his operations through the vagina; Hegar and Tait at the same time performed a similar operation but by means of abdominal section; the latter especially perfected the operation by the abdominal route for the removal of the uterine appendages in pus cases.

*Presented to the Section on Obstetrics and Diseases of Women, at the Fifty-first Annual Meeting of the American Medical Association, held at Atlantic City, N. J., June 5-8, 1900.

This was the beginning of modern abdominal surgery, and the gynecologists of the past few years became pelvic surgeons, removing tubes and ovaries for all kinds of conditions and symptoms; but the extreme swing of the pendulum has been reached; conservative surgery predominates, and efforts are made to preserve half or at least a part of the generative organs.

The state of perfection to which American surgeons brought celiotomy caused the neglect of the original route suggested by Battey, until, by the effort of the French surgeons, especially, the vaginal route was again brought forward, not only for the relief of cancer, but also for inflammatory and suppurative conditions of the pelvic organs. The extreme French suggestions have in turn reached their limit; and cases are now carefully selected for one route or the other by all able and experienced surgeons.

Battey called his original operation "normal ovarian section," with the idea that the ovaries he removed were normal, but modern investigation has shown that while the ovaries may appear normal macroscopically, sections under the microscope show abnormal conditions. This accounts for the pain, distress and symptoms produced by these structures; we find a good many cases where a woman will complain of a great deal of pain in one ovary, continuing for years and causing a great deal of distress, when by palpitation we can not find anything. Experience has taught us that the removal of that ovary will cure the patient; and the microscope will show that the ovary was diseased. To remove the diseased ovary by abdominal section, with the accompanying dangers and sequelæ, seemed to me poor practice; so I followed the Germans, performing, in selected cases, what they call vaginal celiotomy. But I have encountered one great trouble, and that is to tie the broad ligament and control the hemorrhage, especially in cases of narrow vagina. If the abdomen be entered between the uterus and bladder, or behind them, it is not easy to remove the diseased tube and ovary when they are in the cul-de-sac; and it is very difficult to apply the ligature to the broad ligament and tie it sufficiently tight to control the hemorrhage; so that I have been obliged, in some cases, to apply a clamp and leave it on forty-eight hours. This, of course, frequently causes sloughing of the compressed part and adhesion posteriorly, and prolongs convalescence.

When the angiotribe was first brought forward by Tuffier I was sure here was the instrument which would enable us to frequently perform vaginal operations, where formerly we would have been obliged to do an abdominal section. Take, for instance, the case of a young girl who has fallen from a swing or bicycle and has dislocated the ovary into the cul-de-sac; there it is; you can not get it back; and if you do, it will soon drop again into the abnormal position; with every defecation there is pain; every month a few days are spent in bed, and there is constant misery and suffering.

I have tried all kinds of treatment and the only thing I see to do, as a rule, is to remove the ovary. To do that by abdominal section and leave a scar is certainly not desirable. With a small opening in the cul-de-sac, you can pull down the ovary and tube, apply your angiotribe, crush the tissue, and cut off the diseased parts. It takes only a few minutes, and in a few days the patient is well with nothing to show that she ever underwent an operation, while the danger is virtually nil.

When I went to Europe last year, I thought I would investigate the subject and select whatever instrument I thought best. I examined the Tuffier, Doyen, Heftler and Thumin—of Berlin—and the latter instrument I

have selected as the best. It has a narrow blade of great strength, and instead of a wheel, where your power to crush is limited, it has a small handled lever, which gives you an immense power and is less trouble to apply and hold in exact place than the others; at least it seems so to me. With a little practice you soon get the knack of it and can use it without trouble. I have used this in abdominal hysterectomy without ligatures, and I have used it in vaginal hysterectomy without using a single ligature, so that I consider it perfectly safe to control hemorrhage, but occasionally, if I had a bleeding point that I could not reach and was suspicious of, I would apply an extra ligature. I would not endanger my patient's life by taking any chances, but as a rule I have been able to perform these various operations without the use of a single ligature or clamp, simply by the use of Thuman's angiotribe.

I mention these cases to impress you with the perfect safety of the instrument; its greatest use, however, has been in the removal of small ovarian tumors, of diseased ovaries in the cul-de-sac, and of pus tubes which were between the uterus and rectum. All these could be readily pulled out, the angiotribe be applied to the pedicle, and a strip of gauze introduced, which was allowed to remain for four or five days.

I will not burden you with a report of a number of cases. I simply want to urge a further trial, so that the real status of this instrument may be fixed. I am sure that it will be found to lessen the period of convalescence in pelvic surgery and to save more lives than any instrument hitherto devised.

Therapeutics.

Vulvitis.

- R. Liquoris plumbi subacetatis.....3i
Tinct. hyoseyami.....3ii
Aq. camphoræ, q. s. ad.....3viii
M. et. ft. lctio. Sig. Apply with saturated cloth.—*Waring: Medical Record.*

Superficial Burns.

Rectus recommends the following ointment:

- R. Iodoformi.....gr. 7½-15
Antipyrini
Acidi boricæ, aa.....gr. 75
Vaselini.....gr. 600

M. Wertheimer applies to burns, especially in children, the following liniment:

- R. Thymol.....gr. 1½-2
Aquæ calcis
Olei lini, aa.....gr. 1500

M.

Starr, to allay the pain and prevent the formation of bullæ, applies instantly to a superficial burn the following ointment:

- R. Ferri perchloridi.....gr. 90
Vaselini.....gr. 360

M.

Haas uses an ointment made by dissolving from 150 to 300 grains of aristol in 600 grains of lanolin and the same quantity of vaselin.

Landolt and Gyax employ the following formulæ:

FOR BURNS OF THE FIRST DEGREE.

- R. Cocainæ hydrochloridi.....gr. 22½
Vaselini
Aquæ destil., aa.....gr. 150
Lanolini.....gr. 45

M.

FOR THOSE OF THE SECOND DEGREE.

- R. Cocainæ hydrochloridi.....gr. 22½
Salol.....gr. 75
Vaselini.....gr. 375

M.

FOR THOSE OF THE THIRD DEGREE.

- R. Euphphen.....gr. 22½
Olei olivæ.....gr. 52½
Lanolini.....gr. 225
Vaselini.....gr. 450

M.

This ointment is said to be especially serviceable in burns of the eyelids.—*N. Y. Med. Jour.*

Some Uses of Pilocarpin.

S. Harnsberger notes some uses of this drug not mentioned in recent text-books. He cites three cases illustrating its benign effects in orchitis; it is given in 1/8-grain doses of the hydrochlorate, repeated every two to six hours, with ½-grain doses of codein, as may be necessary to produce copious sweating and relieve pain. The same drugs, in similar combination, are of great service in cholelithiasis and nephrolithiasis. In both hepatic and nephritic colic the pilocarpin is best administered subcutaneously. It deserves a trial in tonic spasm of the diaphragm in certain cases of hiccough; in cases of stricture and obstruction of the intestine when ice-poultices and wet-compresses are ill borne, codein and pilocarpin may be used with benefit. The easiest and quickest way to remove ranula is by means of one-sixth grain of pilocarpin hypodermically.—*Medical Review.*

Topical Use of Quinin in Leucorrhea.

Dr. Hardwicke speaks as follows about the topical application of quinin to the mucous membrane of the cervix uteri and vagina in cases of leucorrhea:

A patient, the mother of six children, who had been a sufferer from the above complaint for some years, having used the various remedies usually prescribed in such cases, but with only temporary benefit, her trouble sooner or later recurring, adopted the use, from prudential motives, of what proved to be quinin pessaries. Since using them not only had her leucorrhea disappeared, but her general health had improved. I have since used quinin topically, in several cases of simple leucorrhea, always with great success—in fact, I do not know of a single instance in which it has failed or in which quininism has been produced. It may be used in the form of douche or pessary. I adopt the latter form as being obviously the better one; the drug has a better chance of closer and more continuous contact with the congested membrane. I prescribe three grains of the hydrobromate in a ½-dram pessary in combination with oleum theobromatis, but the pessus quininae of the "Extra Pharmacopeia," containing the hydrochlorid, answers just as well. One insertion a day is generally sufficient.—*Canadian Pract. and Review.*

Copper Phosphate in Tuberculosis and Chlorosis.

- R. Cupri acetat (neutral).....gr. 1/6
Sodii phosphat. (cryst.).....gr. 5/6
Glycyrrhizæ
Glycerini, aa q. s. ad. pil. i.

M. Sig. two pills to be taken daily.

- R. Cupri phosphat.....gr. xv
Aquæ destil.
Glycerini puræ, aa.....gr. xxxviii

M. For hypodermic injection.—*Le Progrès Méd.*

Senile Pruritis.

Upon the principle that senile pruritis is dependent upon intestinal autointoxication, it is recommended that after purging the patient, a milk diet be ordered and an intestinal antiseptic (2 gm. benzonaphthol daily) be administered. Even in severe and obstinate instances, good results have been obtained within twenty-four hours.—*Parisot: Semaine Méd.*

To Create a Distaste for Drink.

Jergolski claims that eight drops of tincture of strophanthus three times a day will create a distaste for drink.

The following prescriptions have been recommended for alcoholism:

- R. Auri et sodii chloridi.....gr. 1/24
Strychninæ nitratis.....gr. 1/60
Nitroglycerini.....gr. 1/100
Atropinæ sulphatis.....gr. 1/200
Digitalini.....gr. 1/60
Sodii chloridi.....gr. 1/8

M. ft. tab. No. 1. For hypodermic use. —*Dunham.*