movements of the canal by depressing the action of the cord or by blocking the impulses from the splanchnics. As maintained by Craig¹¹ and others, salicylate of physostigmin acts favorably as a remedy for postoperative paresis in clinical cases. In all probability its usefulness is restricted to cases in which the paresis is of central origin, for its physiologic action includes depression of the functions of the spinal cord and powerful stimulation of the stomach and intestines¹²—precisely the requirements for obviating reflex inhibition of the alimentary canal.

We have tested the action of physostigmin salicylate experimentally on animals. A sixtieth of a grain in a freshly-prepared solution was given subcutaneously at various times with relation to the etherization and crushing of the testicles—two hours before etherization, an hour before etherization, during the anesthesia, and subsequently while the food was stagnant in the stomach. In our experience giving physostigmin at any time before the operation or during the anesthesia resulted in no perceptible improvement of the passage of food through the alimentary canal. But when the drug was given afterward during the period of stagnation there resulted a very marked, but only temporary, increase of peristalsis. In the following figures a comparison is presented between the amount of food in the small intestine in two typical cases, one in which the testicles were crushed, the other in which the testicles were crushed and a sixtieth of a grain of physostigmin administered about two hours after the feeding. The figures give in centimeters the aggregate length of the food masses in the small intestine at the times indicated:

	Time in Hours.						
	½	1	2	3	4	5	6
	cm.	cm.	cm.	cm.	cm.	cm.	cm.
Ether, testicles crushed....	0	0	0	0	0	8	15
Ether, testicles crushed, physostigmin at two hours...	0	0	0	32	38	41	37

It is clear from the figures for the third hour that the administration of the drug at the second hour had produced suddenly a very abundant discharge from the stomach into the intestine. Nor was gastric peristalsis alone stimulated; intestinal peristalsis was also remarkably active—lively movements of the food masses in the intestinal coils were seen on all sides. But at the fourth hour and thereafter all was quiet again. As shown by the figures, the discharge from the stomach did not continue to be abundant and the food in the intestinal coils lay almost entirely quiescent; at the end of six hours no food had reached the large intestine. The physostigmin, therefore, although given in an amount sufficient to cause the cat's muscles to twitch, had only a temporary effect. Within two hours this action had apparently ceased. These results raise a question as to the reasonableness of attributing great efficacy to a single dose of physostigmin in clinical cases, when the passage of feces occurred from eight hours to three days after the administration of the drug. It would seem as if physostigmin, because of its relatively brief action, could not properly be held accountable for the results. Be that as it may, however, there is no doubt that, temporarily at least, the salicylate of physostigmin will destroy reflex inhibition of the alimentary canal and cause the normal movements to proceed.

In conclusion it should be stated that the present report of our work is preliminary and that many features

need still to be made clear. The point we wish to emphasize is the distinction between dynamic ileus of spinal origin and dynamic ileus of local origin in the wall of the gut. The various conditions which affect one or the other of these causes of inactivity have not yet been experimentally studied, but on such a study depends the possibility of rational judgment in any particular case.

THE SYMPTOMS AND DIAGNOSIS OF STRANGULATION OF THE INTESTINE.*

JOHN B. ROBERTS, A.M., M.D.

Professor of Surgery in the Philadelphia Polyclinic.
PHILADELPHIA.

The development of the operative surgery of the abdomen has created in the minds of surgeons such an assurance of success that they are liable to neglect the study of diagnosis and to resort too cheerily to exploratory investigation of lesions in this cavity. Physicians, on the other hand, being more or less unfamiliar with the innocuousness of aseptic inspection of the abdominal contents, lose valuable time in fruitless efforts to overcome mechanical impediments to peristalsis. Patients would have a fuller measure of scientific aid in their dire extremity, if surgeons could profit by frequent opportunity to see cases of intestinal obstruction early enough to study initial symptoms. On the other hand, many lives would undoubtedly be saved, if physicians realized the futility of wasting time in the effort to settle diagnostic problems, when the fecal current and perhaps the blood circulation are stopped by mechanical causes, demanding instant manipulative removal.

Such conditions demand the best consideration of philosophical minds, alert to appreciate the importance of both medical and surgical skill. Good "team" work on the part of a physician and a surgeon, or the services of a practitioner who has had both medical and surgical training, will give the suffering patient the best chance for recovery. Some of us, who now practice surgery as a specialty, would often be most unhappy, if we did not possess, through previous experience in internal medicine, some ability to weigh our own surgical opinions and even to review critically the judgments of our medical colleagues.

The symptoms of intestinal obstruction and intestinal strangulation are necessarily similar, because strangulation is usually a complication of obstruction. It is an interference with the vascular and lymphatic circulation in the wall of the bowel and in the mesentery. It may occur as an incident in the history of acute or chronic obstruction. In the former the strangulation may arise coincidently with the obstruction. In both it may arise only after the obstruction has existed for a considerable time. Occasionally, as in some hernias, a portion only of the wall of the intestinal tube may be strangulated with no real occlusion of the caliber of the gut. Yet the influence exerted on the muscular coat of the bowel produces, as a rule, very promptly a symptom-complex practically identical with that of obstruction. These symptoms usually continue until the strangulation is relieved or the patient dies; sometimes, however, local gangrene occurs, an artificial anus is formed and the patient recovers.

Ignorance of the causes of the symptoms formerly

11. American Jour. of Obst. and Dis. Women and Children, 1904, xlix, p. 449.

12. Cushny (A. R.): Pharmacology and Therapeutics, Philadelphia, 1901, 317, 319.

* Read in the Section on Surgery and Anatomy of the American Medical Association, at the Fifty-eighth Annual Session, held at Atlantic City, June, 1907.

termed ileus, or iliac passion, no longer compels the use of such indefinite terms. The Greek verb *ελεῖν*, from which comes *ελεός*, is practically equivalent in a medical sense to the Latin *volvō*. The term ileum, therefore, should be restricted to that lesion of the bowel in which the intestine is rolled, twisted or knotted, so as to produce symptoms of obstruction. Ileus and *volvulus* should be synonymous terms. The word *volvulus* was formerly rather loosely employed, though it is not now.

Inspection of English medical dictionaries will be convincing as to the need for this restriction of the use of the word ileus or its complete elimination from medical nomenclature. A number of mechanical disarrangements of the intestinal tube may cause colic, constipation and vomiting, but there is no good etymologic or other reason for classing them all under the name ileus. As hernia, from *ἔρως*, a sprout, should be applied to a protrusion, so ileus, from *ελεός*, should be applied to twisting, and to twisting only, of the bowel. The word should not be applied to intestinal obstruction, strangulation or colic from any other cause.

It is often difficult to make a differential diagnosis between intestinal obstruction and such conditions as lead colic, flatulent colic, arsenical poisoning, acute enteritis, Asiatic cholera, acute pancreatitis, perforated ulcer of the stomach, thrombosis and embolism of the mesenteric arteries, torsion of the great omentum, inflamed undescended testicle, paralysis of the intestinal muscles, hysteria, acute appendicitis and peritonitis. The examination of the patient should be repeated from hour to hour, in order that no one of the rapidly developing diagnostic points be overlooked. It is only in the earliest stages of the affection that an exact diagnosis between the different forms of obstruction is likely to be feasible. The development of tympanites as a symptom or of peritonitis as a secondary condition often destroys the possibility of an accurate investigation except by exploratory investigation.

Strangulation not only closes the caliber of the bowel, so as to prevent the movement of the fecal current and the gaseous contents, but it also severely irritates the nerves in the mesentery and in the gut and compresses the blood and lymph channels in those structures. The result is that grave arterial anemia and venous hyperemia are produced, and gangrene and perforation may occur. It is a more or less elastic constriction and may be defined as obstruction plus circulatory interference.

An external hernia at the inguinal, femoral or umbilical opening may become strangulated. So may a hernia through one of the more unusual orifices, such as the obturator foramen, the sciatic notch, and the small openings occasionally found in the linea alba, between the navel and the ensiform cartilage. The medical attendant should never neglect to explore these recognized seats of external hernia whenever intestinal obstruction is under consideration. A small femoral hernia may be concealed by enlarged, overlying lymph nodes. In children a scrotal hernia may be translucent and suggest hydrocele. The rectum must always be examined with the finger or proctoscope, to exclude the existence of impaction of feces or stenosis of the lower bowel from tumor. Thomas Bryant called attention years ago to a ballooning, or dilatation, of the rectum, when stenosis of the lumen of the descending or sigmoid colon exists. This symptom may lead to the discovery of an occlusion from carcinoma or other cause too high to be felt with the finger inserted through the anus.

The results of strangulated gut are practically the

same, whatever the anatomic lesion causing the constriction. It makes little difference in the general aspect of the case whether the strangling be due to compression by, entanglement in, or twists, kinks or knottings from omphalo-mesenteric vestiges or peritonitic adhesions; to incarceration in congenital or acquired apertures or fissures; to internal hernias in the foramen of Winslow, the paracecal fossæ the intersigmoid space, the duodeno-jejunal region, or through diaphragmatic openings; or to an external hernia incompletely reduced or not reduced at all. A mere tear in the parietal peritoneum may afford an opportunity for the intestine to slip under it and become strangulated.

The symptoms do not in the main vary much from the characteristic complex when the strangulation arises secondarily to a chronic obstruction of the bowel. Here, however, there is usually discoverable a history of a series of antecedent events pointing to stenosis or obturation.

The diagnosis of probable strangulation, primary as well as secondary, must rest on the sudden onset of great pain, vomiting, absolute stoppage of feces and gas, and intense prostration or collapse. These symptoms may arise immediately after the receipt of a blow on the abdomen, or their advent may follow sudden violent muscular exertion, straining at stool or active purgation from diarrhea or a cathartic.

The pain is generally the first symptom. It is violent, continuous and occasionally increased by paroxysmal colic. It is practically always a part of the symptom-complex, unless gangrene has occurred and the patient has gone into collapse and stupor. It may be localized at or near the site of strangulation and may be relieved somewhat by pressure. In *volvulus* the pain is likely to be more diffused than in internal strangulation by a band or the like. This initial pain may in time be followed by the secondary pain which comes from the development of peritonitis, provided that peritonitis develops ere the death of the patient or his relief by treatment. The pain of peritonitis is not relieved by pressure. It is possible that a slight degree of tenderness on pressure may be present from increased sensitiveness of the bowel, when no peritonitis has as yet developed. This is probably due to the violent contractions of which the muscular tunic of the bowel has been the seat.

The advent of pain is quickly followed by vomiting, due probably to the traction on the peritoneum at the point of strangulation. The vomiting under rare circumstances may precede the access of pain. It usually occurs within the first twenty-four hours and persists until death, though it may cease shortly before that issue. It is accompanied in very many instances by nausea, hiccough and eructations of gas. The vomitus is at first the contents of the stomach; then the intestinal fluids, colored by bile so as to be brownish, yellowish or pea-soup-like, are ejected, and finally fecal vomiting is established. Fecal vomiting need not occur. The seat of the obstruction may be too high in the intestinal canal or the patient may succumb to shock before it has appeared. The time of its appearance may vary from two to five or more days. Death is not uncommon in two or three days, though there are cases that run a more chronic course.

The complete obliteration of the caliber of the gut and the reflex influence exerted by strangulation induce absolute constipation. This obstipation appears as soon as the strangulation occurs and no discharge whatever of feces or flatus takes place. Even when the strangu-

lation affects the ileum the contents of the large bowel are not expelled. There are but very few exceptions to this rule of absolute constipation. Peristalsis below the site of the lesion seems to be inhibited by reflex action. The contents of the rectum and lower colon may be washed out with enemas, but there is as a rule no spontaneous evacuation of intestinal gases or of feces. This constipation may be difficult to differentiate from the paralytic intestinal inertia, which may follow incision of the abdominal muscles and manipulation of the bowels during the operative treatment of various abdominal and pelvic lesions. Such inhibition of peristalsis may occur without the existence of postoperative peritonitis.

Coincident with this more or less characteristic symptomatic triad of strangulation a most profound collapse occurs. This prostration is evidenced by a frequent small pulse, pallor of the skin, low temperature, anxious facies, sunken eyes and the congested internal viscera of shock. Death may take place as a result of this condition, which is probably due to the reflex inhibition of the heart's action and a consequent lowered blood pressure. The immediate cause of the fatal issue is probably anemia of the nervous centers. Similar profound depression of the vital forces occurs in other forms of acute intestinal obstruction, but it is especially noticeable in instances of obstruction with strangulation. The exact form of the strangulation does not seem to exert much difference in the degree of shock. In this form of shock, as in surgical shock from other lesions, the patient remains conscious except in the agony of death.

The severity of symptoms depends on the degree of the strangulation and the amount of bowel and mesentery deprived of its normal circulation. The tighter the strangulation and the greater the extent of bowel involved the more urgent the symptoms. Death from shock, from general exhaustion, caused by the pain and vomiting, or from peritonitis is to be expected early in severe cases. Death is likely to occur before gangrene and perforation take place.

Tympanites occurs only to a moderate degree, as a rule, in strangulation from hernia and bands, but is apt to be marked in volvulus. When, however, peritonitis has developed as a secondary complication the distension becomes general and may be very great. The meteorism is localized at first in the involved loop of bowel and the portion of intestine just above the strangulation. It should be remembered that an extremely distended gut does not give a tympanitic note on percussion, but a low note more like that obtained when the chest is percussed over normal lung.

If no localized tympanites is seen it is probable that the strangulation is incomplete or that the intestinal gases are escaping through minute perforations in the bowel wall. It is the venous stagnation in the bowel from the interference with circulation that causes the localized meteorism due to unabsorbed gases. This localized tympanites differs from the distension that arises from putrefaction of stagnating intestinal contents, and occurs more promptly than the latter.

The locally distended bowel in strangulation changes the shape of the abdomen; hence the least asymmetry should be carefully studied. The bulging of the flanks laterally, the puffing up of the umbilical region in which lies the small intestine, or the distended outline of the colon or of the sigmoid flexure may give a hint, if the surgeon sees the case early and is alert. The strangulated loop is anchored in one spot, and as it becomes distended changes the aspect of the abdominal surface.

The rigid loop may be detected by palpation, because it feels tense and hard. General anesthesia may, by relaxing the abdominal muscle, aid in enabling this fact to be discovered by the examiner's hand.

A study of the urine, feces and blood may perhaps aid in establishing a diagnosis of the existence of obstruction, if not in actually helping the surgeon to a conclusion as to the occurrence of actual strangulation. The discovery of a large amount of indican in the urine during the first two or three days seems to indicate either acute peritonitis or occlusion of the small intestine. Hence when the other symptoms exclude acute peritonitis, intestinal obstruction may tentatively be assumed to exist in the small bowel, if collateral evidence points to interference with the permeability of the intestinal tube. According to Hammer, peritonitis, multiple stenosis in both small and large intestine, and a chronic stenosis in the large bowel may show excessive indicanuria. These possibilities, therefore, modify the value of this symptom. If excessive indicanuria is absent, acute general peritonitis is probably not present. A deficiency in the amount of urine excreted is considered by some writers to be attributable to tightness of the constriction of the caliber of the gut. Others maintain that it simply indicates that, in a case of obstruction, the stenotic lesion is high in the small intestine. Albuminuria and hematuria have little diagnostic significance.

It is unfortunate that little can be learned by inspection of the feces, except occasionally in obstruction in the rectum or the descending and sigmoid colon, when the shape of the fecal masses may be of aid. Blood and pus in the stools, especially the former, have a limited significance. Small amounts of blood are discharged from the rectum in some cases of volvulus, but not in internal strangulation unconnected with volvulus, malignant disease or invagination. In invagination bloody stools are often a marked characteristic.

Leucocytosis is of possible value, when markedly developed within forty-eight to seventy-two hours after the beginning of obstructive symptoms. Its occurrence in various suppurative abdominal conditions without obstruction destroys its value, when found only at a later period.

Fever occurring early in a suspicious case tends to make logical the diagnosis of peritonitis rather than obstruction or strangulation. Its development later may be due to an incidental peritonitis complicating strangulation or any other form of obstruction, but it should be recollected that peritonitis may occur without fever.

The diagnostic value of peritoneal fluids obtained by aspiration, of intubation and inflation of the colon, of auscultation of the abdomen, of the antiperistaltic progress of minute particles from the rectum to the stomach, where they are found by lavage, and of the use of the x-ray to locate masses of dammed up feces made demonstrable by bismuth subnitrate should be tested in appropriate cases.

The diagnosis of obstruction is always a matter of supreme importance, because the patient must be relieved spontaneously or by treatment. Otherwise death will occur. Prompt relief is necessary in all cases, but promptness is absolutely essential in the crisis caused by strangulation. Often the diagnosis can only be made by a process of exclusion. If accurate diagnosis is unattainable, operation must be at once undertaken. Exploratory operation is frequently a confession of ignor-

ance, but such confession is an evidence of sincere concern for the welfare of the patient.

Let it be remembered that there may be more than one point of obstruction or strangulation, especially in cases secondary to former attacks of peritonitis. Do not forget that visible intestinal peristalsis with paroxysmal rigidity and stiffening of definite loops of intestine, after percussion of the abdominal wall, are absent in volvulus (axial rotation) and internal hernia with strangulation, unless these conditions are engrafted on chronic obstruction. This symptom may be of diagnostic value, however, to establish the presence of intestinal obstruction in chronic forms of occlusion, where the muscular coat has become hypertrophied.

Recollect the absence of peristalsis and the possible occurrence of adynamic obstruction in appendicitis, renal and biliary colic, and peritonitis, which are due to inhibition of power in the musculature of the intestine. In these instances pain and sensitiveness may prevent the passage of feces along the intestinal canal and simulate obstruction. There is perhaps a sort of neurotic constipation here, such as may occur in healthy persons who have not access to decent toilet facilities.

When the surgeon has concluded that strangulated intestine is causing the symptoms, he should endeavor to make up his mind as to the location of the trouble and the character of the strangulating process.

The most common site of strangulating lesions, not associated with internal hernia, is the right lower quadrant of the abdomen. The lower portion of the ileum and the cecal region, therefore, should be inspected first. Volvulus, however, is most apt to occur in the sigmoid colon, especially in the aged. Intussusception is not particularly common in the ileocecal region, but it is perhaps most frequently found in some portion of the colon. It is the more frequent disease of the young. Strangulative lesions, being usually acute, are more difficult of differentiation than those other obstructive lesions which are chronic.

The change in contour of the belly, due to tympany, will sometimes aid in settling the site, as has already been stated. In obese patients this determination becomes a matter of increased difficulty. It must be recollected that the sigmoid colon twisted on its axis may fill from distension the abdominal cavity in an astonishing way and extend even to the right hypochondrium. In such cases, however, the left flank also is apt to be convex from distension. The upper lumbar region, on both sides behind, will give on percussion, it is said, a long deep note instead of the normal high and flat note, when the obstruction is in the sigmoid or descending colon. This change in percussion note in the back will be found only on the right side, when the occluding lesion is in the transverse colon or at the splenic flexure. This sign seems to have some value.

The pain is less severe in volvulus than in strangulation by omphalo-mesenteric remains, or bands; and sometimes the pain of volvulus may be referred to the back. In volvulus high up in the small bowel there may occur frequent evacuations from the bowels. This is unlike internal hernia-form strangulation and strangulation from bands, which are accompanied by absolute constipation. Occasionally a little blood may be passed with the feces in cases of volvulus, especially of the sigmoid flexure. Rectal tenesmus is an indication of volvulus being the cause of the obstruction, rather than strangulation from other causes. Fecal vomiting is not common in volvulus, but it is common in strangulation

from entanglement under bands and in openings. Volvulus is more apt to be preceded by habitual constipation, and causes less intense collapse than strangulation from bands.

A previous history of peritonitis would point rather to strangulation by bands, or kinks or knots, than to strangulation from volvulus, were it not a fact that peritonitis may cause narrowing of the mesentery from infiltration and cicatricial changes which tend to make axial rotation (volvulus) occur more readily.

The differential diagnosis is farther obscured at times by the circumstance that volvulus of the small intestines and strangulation by bands and cords may be coexistent.

My final word is: If the diagnosis can not be established with a fair degree of certainty within twenty-four hours, or at most forty-eight hours, open the abdomen aseptically and inspect its contents.

313 South Seventeenth Street.

[THIS SYMPOSIUM ON ILEUS TO BE CONTINUED. THE DISCUSSION WILL FOLLOW THE LAST PAPER.]

THE OCCURRENCE OF TYPHOID BACILLI IN THE STOOLS OF TYPHOID PATIENTS,

WITH A DISCUSSION OF THE PATHOGENESIS OF TYPHOID FEVER.*

DRS. J. H. PRATT, F. W. PEABODY AND A. D. LONG.

BOSTON.

The view is generally held that the intestinal contents in typhoid fever are swarming with typhoid bacilli. The fact that many investigators have often had difficulty in isolating *Bacillus typhosus* from the stools of undoubted cases of typhoid fever has been attributed to imperfect technical methods. Some observations made by one of us seemed to indicate that the reason typhoid bacilli were not obtained from the stools during the height of the disease might be due, not to faulty bacteriologic methods, but to the fact that the typhoid micro-organisms were either absent from the feces or present in extremely small numbers.

The studies here presented extend over two years. About 75 per cent. of the patients had fever at the time of the examination. All were clinically typhoid fever. In only one instance was the agglutination test persistently negative. It is probable that this was a case of infection with *Bacillus faecalis alkalingenes*. Each day the sickest patients to be found in the wards were selected for the examination. For the opportunity of making this study we are indebted to the visiting physicians of the Massachusetts General Hospital.

When Gaffky¹ reported, in 1881, that he had cultivated on artificial media the specific micro-organism of typhoid fever, he made no claim that he could isolate it from the intestinal contents. In 13 cases he obtained the typhoid bacillus from the spleen and in one case from the liver. Pfeiffer,² in 1885, was the first to assert that typhoid bacilli were present in the stools. As he knew nothing about *Bacillus coli* and did not fully

* Read in the Section on Pathology and Physiology of the American Medical Association, at the Fifty-eighth Annual Session, held at Atlantic City, June, 1907.

* From the Laboratory of Clinical Pathology, Harvard University (Director, Prof. R. H. Fitz). This investigation was begun in the Laboratory of Bacteriology. The facilities of the bacteriologic department were available through the courtesy of Prof. H. C. Ernst.

1. Mitt. a. d. Kaiserl. Ges. Gesundheits, 1884, II, 372.

2. Deutsche med. Wochschr., 1885, p. 500.