in the tissues involved before the treatment was commenced. The dangers of infection are such that no one should attempt the operation unless he has an "aseptic conscience," and is the master of an aseptic technic. The presence of infection in the vulva, vagina, or uterus increases very much the danger of dilatation by tents. A purulent discharge from the uterus contraindicates the use of tents, as the dilatation opens up spaces which readily admit infection. The danger of the use of tents in such cases is illustrated by the following:

Mrs. S., suffered from a profuse mucopurulent uterine leucorrhea, the result of an old gonorrheal infection. The uterine cavity had been dilated, curetted, irrigated, packed with gauze, and frequent cauterizing and antiseptic applications had been made, without giving any permanent relief. The cervix was dilated with tents for the purpose of making thorough curettage and for facilitating treatment by gauze packing. She developed cellulitis, which resulted in a pelvic abscess, and which required incision and drainage of the abscess. In this case it was an error to use the tents; rapid instrumental dilatation would have been preferable.

93 E. Eighteenth Street.

PELVIC SUPPURATION.

PROCEDURE AND PLAN OF ATTACK.

Read before the Medical Society of the Missouri Valley, Council Bluffs, Iowa, Sept. 14, 1898.

BY BYRON B. DAVIS, M.D.

OMAHA, NEB.

If we are to draw our conclusions from society discussions, both in this country and abroad, the conviction is forced upon us that there is a greater diversity of opinion in regard to the management of pelvic suppuration than that of any other surgical lesion. Before asepsis had rendered celiotomy so uniformly successful, large pus collections in the pelvis were merely punctured through the vaginal vault; later the abdominal vault was looked upon as the only method deserving of recognition.

The French and Belgian schools next began the practice of doing almost everything by the vaginal route, advocating hysterectomy in all cases of double tubal suppuration. Several Germans and Americans adopted the same line of procedure, while their teachings were energetically opposed by a large number who advocated the abdominal route of attack.

So dogmatic have become the teachings of men practicing the one or the other of these methods that the adherents of the one plan would lead the credulous to believe that to reach a pelvic inflammatory affection from below is due to unsurgical timidity; while the advocates of the other extreme would have us believe that celiotomy in these cases is actuated by almost criminal rashness.

The most conservative discussion on the subject yet held was that opened by Cullingworth at the recent meeting of the British Medical Association in Edinburgh. This discussion was participated in by no less than seventeen eminent surgeons, representing all shades of opinion. The gratifying feature of the discussion is that extremists like Doyen and Jacobs have given up a part of their dogmatism and admit that celiotomy has its place in selected cases; while the English school are willing to concede that there are many cases in which the vaginal route is safest and best. After a careful reading of this discussion, one is forced to the conclusion that safety lies in the practice of neither of the extreme views, but in the careful selection of the method best adapted to each individual case.

No doubt many of us have often been perplexed and in some doubt whether a patient's interests would be best subserved by a radical or so-called "complete" operation from above, or by a safer and less complete operation through the vault of the vagina. A desire to elicit a free discussion on this subject is the purpose of a few remarks today.

In acute attacks of pelvic inflammation, it is often very difficult to decide whether there is greater danger in an immediate operation or in delay. Many cases which seem very serious make symptomatic recoveries, and occasionally I have examined women who had made, so far as could be determined by bimanual examination, anatomic recovery—and even when they had been supposed to have a collection of pus in the pelvis. These experiences have led me to hesitate about advising operation in many of these cases of acute pelvic inflammation.

In only two classes of cases do I uniformly advocate immediate operation: 1, those in which the amount of pus is large and tension great, where there seems to be danger of perforation into the peritoneal cavity, the bladder or the rectum; 2, those cases with profound sepsis, whether the amount of pus is great or small, in which the conviction is forced upon us that unless something is done to stop the septic absorption the patient is bound to succumb.

During the past few months no less than six cases of pelvic suppuration have been met with, in which one or both of these conditions have been present. In each case the patient was in no condition to endure a prolonged operation. All were operated upon through the vagina by free incision and drainage. All made operative recoveries and are now symptomatically well women. Whether any or all may eventually have recurrences it is impossible to say, but in the one case, in which I had an opportunity to make an examination some weeks subsequently to the operation, only the slightest amount of induration remained. Every one of these cases was very weak, with high temperature and rapid, weak pulse, at the time of operation, and to have subjected them to so serious and prolonged an operation as a celiotomy would have surely resulted in a larger percentage of mortality, if indeed any of them would have survived.

It is only just to state that in all but one of these cases convalescence was prolonged over a period greater than in the average celiotomy; but surely safety of the operation is purchased cheaply, even at the price of two or three weeks longer in bed. It is argued by Dr. Joseph Price that this is an operation which has its birth in timidity; is never complete, and leaves the patient in practically as bad a condition as she was before. The experience with these few cases has convinced me that it is better to be accused of timidity than to sacrifice a life on account of not having the courage to perform an operation, because it is supposed to be an indication of timidity.

Of these six cases, the pus was still located in the tubes in four cases, and each tube was opened and drained separately. In the other two cases the pus had broken through the tubes and we had a true peritubal abscess. In three cases the unquestionable cause of the infection was abortion; in the other three it was not determined whether it was abortion or gonorrhea, though the former was probable in all, and in no case could it be positively stated that the infection was gonorrheal. This is, in a small way, evidence in favor of the dictum of A. Martin, who says that very

seldom does a gonorrheal infection require an immediate operation, while infections from abortions frequently require operation in the acute attack.

In all my cases the vaginal fistula closed promptly, but it is only fair to state that a permanent fistula is doubtless to be expected occasionally. Recently I met with such a case, operated by another surgeon, in which a fistula was left. The patient, when she came to me, was in such a condition as not to contraindicate abdominal section, which was done and the woman recovered.

Only twice have I removed adnexa by the vaginal route, except when hysterectomy was done at the same time. In one of these cases no trouble was experienced, in the other so many adhesions were met with that I wished most heartily that the abdominal route had been selected. It gave one a thorough appreciation of the truth that the hand directed by the eye is worth much more than one or two fingers groping in the dark. This assertion, of course, labels me as not a skilful operator by the vaginal route, but I have seen a helpless expression on the face of several skilful vaginal sectionists, when I felt positive that one would have had clear sailing had the abdominal route been chosen. In general there can be no question that celiotomy possesses many points of advantage over vaginal section. Not the least of these is that one can see, as well as feel, what is being done. How frequently, in loosening adhesions, do we find the intestines, if not torn, at least so denuded that it is deemed unsafe to leave them without covering with peritoneum by means of Czerny sutures! In doing such work per vaginam we are almost entirely deprived of an opportunity of resorting to this conservative and life-saving procedure, being often left in doubt as to whether the bowels have been wounded or not. Even when the intestine has not been dangerously injured, it is denuded of peritoneum, and when no opportunity has been given to cover the raw surface, troublesome and painful adhesions are likely to form. Several surgeons are advocating conservative work upon the tubes and ovaries through the vagina. The procedure is to draw the tube and ovary through the vaginal incision, examine, puncture, cauterize, or resect, according to the condition present, and then return the parts to the abdominal cavity. If, as I believe, many of the most painful affections of the pelvis are produced by prolapse of the ovaries, with resulting traumatism by uterus and loaded rectum, this forcible dragging down can not but result in an aggravation of the symptoms, for there is no way of insuring the perfect restoration of such ovaries to their normal position by way of the small and deep opening through the vaginal vault. In chronic inflammations in the pelvis, if there is unquestionably a collection of pus low down, at a point easily accessible from the vagina, the vaginal incision is doubtless safer and in every case to be preferred. In all chronic inflammations, unless the collection is low down, the abdominal incision is in every way better, safer, and more likely to result in symptomatic as well as anatomic cure. As previously stated, in acute inflammations it seems usually better not to operate, except in the conditions mentioned. The question naturally arises: How long is it best to wait after the acute symptoms have subsided before a celiotomy should be undertaken? It is claimed that an acute infection immunizes a patient for a short period—the exact time not yet very definitely ascertained. Believing this to be the case, the

time of election for an operation, if required, would be within a few days after the subsidence of the acute symptoms.

Regarding the technic, the bounds set for this paper will not permit a discussion. One point only I feel constrained to mention. If there is any pelvic tissue it is the worst kind of treatment to resort to irrigations and thus carry infectious matter into the uninfected recesses of the abdominal cavity. It is much better to wipe out carefully and then drain, either through the lower angle of the abdominal incision, or, what is better, through the vagina. It is very easy to make an incision through Douglas' pouch and thus get downhill drainage. Rarely, it may be necessary to make use of through-and-through drainage.

THE MEDICAL PROFESSION.

IS IT GOING INTO VOLUNTARY BANKRUPTCY?

BY LEWIS SCHOOLER, M.D.

DES MOINES, IOWA.

The plea of an overcrowded profession and low fees, as well as too many medical colleges and free clinics, have each in turn received a great deal of attention, with apparently no results, unless upon the theory of suggestion it has increased the evils that are complained of. This is, no doubt, more perceptible to clinicians than to others; and having conducted a clinic for many years, I may possibly be able to delineate a few occurrences that will prove interesting to the profession at large.

Clinical teaching has been so emphasized by the colleges and by the demands of the profession, that undue importance has been given to that feature in all colleges during the last few years, so much so that the institutions that have advertised the greatest number of clinics have generally obtained the largest classes. That this has been a great gain to the profession in general, the writer emphatically denies, and equally emphatically denies the supposed gain to the students. In acceding to the demands of the profession for strictly clinical teaching, prodigious efforts are put forth in order to obtain the necessary clinical material. The friends of the school are either publicly or privately encouraged, if not absolutely solicited, to act as agents of the school, in order that the material may be abundant. When this is the case it necessarily follows that little or no scrutiny is exercised as to the character of the persons who wish to avail themselves of the advantages of Professor —'s clinic, and the distinguished professor, being partially intoxicated by reason of his rapidly increasing popularity, and entirely forgetting that a "free show always draws well," continues to operate upon poor and rich alike.

By way of confession, and that I may not seem to have the other fellow constantly in view, to the exclusion of all retrospective researches, I will admit that I have myself, operated upon at least one case who came to the hospital gorgeously bedecked with diamonds worth many hundreds of dollars, besides being clothed in "purple and fine linen," with a seal-skin sacque over all. Others may or may not have had a similar experience. That this was a wrong to the profession, I confess. That similar wrongs are being perpetrated daily is not doubted at all by any one familiar with clinical work. The real question at issue is: Who is to blame? For one, I am willing to share a portion of it. There are few, if any, schools