Collapse and Treatment.—The operation had just begun, the incision was made, when it was observed that the patient had become intensely cyanosed. In a moment he was pulseless and soon afterward respiration ceased. The head of the table was rapidly lowered and artificial respiration at once resorted to. Realizing the otherwise hopeless situation, the abdomen was rapidly opened and with one hand on the chest wall and the other against the diaphragm I massaged the heart at the rate of from 15 to 20 movements to the minute. The actual time was not recorded in the excitement of the moment, but in approximately 2 minutes feeble but irregular fibrillary contractions were felt. Gradually the impulses became more forcible and more regular, and after an interval of 8 minutes, during which time massage was continued, resuscitation seemed assured. At this time the pulse-rate registered 130.

Result.—The abdominal wound was closed, the operation on the hydrocele completed and the patient returned to the ward. The subsequent history was uneventful.

In a series of forty-six cases, reports of which are collected by Cockovic,¹ in seventeen the results were classified as positive and in twenty-nine as negative. Of the former, in only nine was the effect permanent; in the other patients survived for from five to twenty-seven hours. Thus there was only 20 per cent. of successes.

Perhaps the most important practical consideration is the method of procedure. Shall the heart be massaged directly through a thoracoplastic opening, by the transdiaphragmatic or by the subdiaphragmatic route? Dismissing any theoretical considerations, we find the best results are obtained by the subdiaphragmatic route. In sixteen instances of the latter there were ten positive results, in twenty-four of the thoracic route only seven; and in the six cases by transdiaphragmatic route there were no resuscitations.

The following table² shows the relative value of the various methods:

	Successful.	Partially Successful.	Failures.	Total.
Direct	2	8	18	28
Transdiaphragmatic	0	1	2	3
Subdiaphragmatic	8	5	6	19
Total	10	14	26	50

Because of its greater simplicity and higher percentage of recoveries the subdiaphragmatic route should be adopted as the conventional procedure, except perhaps in children in whom the elasticity of the chest wall enables one to make adequate pressure on the heart without opening either chest or abdomen. How long one is justified in waiting before resorting to massage may be judged from the fact that there have been no recoveries when the syncope has been of more than ten minutes' duration; and the largest measure of success was obtained when the interval did not exceed five minutes. The sooner the massage is begun, therefore, the brighter the prospects. As the proper oxygenation of the blood is indispensable to resuscitation, artificial respiration should at the same time be practiced, or direct insufflation by the Auer-Meltzer method.

In a case reported by Rehn,³ failing to resuscitate the patient by subdiaphragmatic massage and Sylvester's method of artificial respiration, he opened the diaphragm, practiced direct massage, and at the same time established positive pressure with the Brauer apparatus. The result was almost instantaneous on both heart and respiration. In two hours the respirations failed again and further attempts at resuscitation were ineffectual. The secondary collapse was attributed by Rehn to a pneumothorax, which developed when the pleura was torn in opening the diaphragm.

NOMA (CANCERUM ORIS) IN AN ADULT, WITH REPORT OF A CASE

HOWARD D. KING, M.D.

NEW ORLEANS

Noma in the aged is of such unusual infrequency, compared with its occurrence in children, that I am induced to record a case which came under my observation in the summer of 1910. Considering the rarity of the condition as well as the scarcity of the literature dealing with the disease, especially as relates to its presence in the aged, I believe that I am justified in offering a brief account of the nature of noma.

Noma is a rapidly progressive and probably specific infectious process of a gangrenous type, causing extensive tissue destruction and usually having its origin on the mucous membrane of the vulva, gum or cheeks, and sometimes, though seldom, the ear or anus. Noma affecting the region of the cheeks or gums is known as cancerum oris; involvement of the vulva is termed noma pudendi.

Children between the ages of 3 and 8 years are most frequently attacked; the disease occurs more frequently in girls than in boys. This sexual predisposition continues throughout life and is not due to any peculiar susceptibility, but to the fact that the female genitalia offer a more suitable field for infection. Text-books on pediatrics are almost unanimous in declaring the condition to be one essentially of childhood and that the disease is not seen in nurslings or adults. That this impression is somewhat erroneous will be seen from the case reports below.

Between the years 1848 and 1903, an interval of fifty-five years, several cases in the aged have been recorded in medical literature. Tourdes¹ 1848, reported nine authors who had seen noma in persons between 15 and 72 years of age; von Bruns² 1859, from 413 cases collected by him, found only eleven in persons whose ages ranged between 15 and 70; Struch³ 1872, reports a case of his own in a patient aged 30, and calls attention to a case observed by Vogel in a youth of 18; Hildebrand⁴ 1873, collected a series of cases numbering twenty-two in persons over 15 years of age; Brydon⁵ 1882, one case; Herman⁶ 1883, one case; Murillo⁷ 1886, one case; Herff⁸ 1890, one case; Masterman⁹ 1891, two cases; Koster¹⁰ 1892, one case; Stephenson¹¹ 1892, seven cases; Ziegler¹² 1892, one case; Le Count¹³ 1898, one case; Mangubi¹⁴ 1898, one case; Zusch¹⁵ 1901, one case; and Reale¹⁶ 1903, one case.

Poorly nourished, cachectic subjects and those living under vitiated environments are said to be the types most frequently affected. That filth, squalor and tenement life are not alone responsible for the disease

1. Tourdes: Thèse de Strassburg, 1848.
2. Von Bruns, V.: Handbuch der practischen Chirurgie, I, Part 2.
3. Struch: Inaugural Dissertation, Göttingen, 1872.
4. Hildebrand: Quoted by Ziegler, *vide* reference 12.
5. Brydon: Brit. Med. Jour., 1882, p. 838.
6. Herman: Trans. Obst. Soc., Lond., 1883, xxv, 141.
7. Murillo: Noma (*a propósito de un caso en un adulto.*) Rev. med. de Chile, Sant de Chile, 1886-87, p. 312.
8. Herff: Deutsche med. Wehnschr., 1890, xvi, 949.
9. Masterman: Two Cases of Spreading Gangrene Around the Mouth Occurring in Adults, St. Bartholomew's Hosp. Rep., London, 1891, xxvii, 2055.
10. Koster: Ett fall af noma hos 44-årigvinnna, Förh., Götterborg Läk. Sällsk., 1892, p. 74.
11. Stephenson: Cancerum Oris in Adult Life, Brit. Guiana Med. Ann., Demerara, 1892, p. 127.
12. Ziegler: München. med. Wehnschr., 1892, xxxix, 107.
13. Le Count: Tr. Chicago Path. Soc., 1898, iii, 180.
14. Mangubi: Sluchai noma i v. vzroslyvo. Vrach. zapiski, Mosk., 1898, v, 18.
15. Zusch: Ein Fall von Noma mit ausgedehnter Soorbildung in hohen Alter, München. med. Wehnschr., 1901, xvii, 785.
16. Reale: Un caso di noma in adulto, Policlinico, Rome, 1903, ix, 1431.

1. Cockovic: Arch. f. klin. Chir., lxxxviii, No. 4, p. 917.
2. Progressiva Medicine, March, 1911.
3. Rehn: München. med. Wehnschr., Nov. 30, 1909.

has been conclusively demonstrated in more than one instance, especially in the epidemic of noma which occurred in an orphanage at Albany, N. Y., where good practical sanitary, hygienic and dietetic conditions prevailed. It also occurs very frequently as a complication of the exanthemata, notably measles.

In the above-mentioned cases of noma the underlying conditions have been most varied. Thus, we see the disease, according to the case reports, following erysipelas, typhoid fever, amebic dysentery, pleuritis and nephritis. Stephenson¹¹ regards renal disease as an important factor in the causation of cancrum oris in the aged.

The exact status of the disease as to etiology has not been definitely established and is further rendered confusing by the multiplicity of the terms used in the description of the malady. Noma, at all events, is a rare condition, but its assumed rarity was very probably due to the difficulty experienced in recognizing an exact causative organism. At the present time, even, it is by no means certain that the organisms described by investigators of the past have anything to do with the disease. The frequent use of the term "noma" synonymously with "gangrenous stomatitis" is to be regretted, as the latter term is purely a generic one and of no precise significance, nor does it furnish us with any adequate idea of the existing pathologic condition. The following terms have been used and are to-day employed to describe noma: gangrenous stomatitis, gangrena oris, gangrenopsis, stomato-necrosis, necrosis infantilis, buccal anthrax, water canker, cancer aquaticus, and sloughing phagedena. *Wasserkrebs*, a term originally coined by A. Richter in 1828, in his initial treatise on noma, is the name given the disease by the Germans. The French use the phrase *gangrène de la bouche*. The foregoing terms are a source of confusion and of no importance save from a symptomatic standpoint.

Reverting to the etiology of the disease, it was once believed to be the manifestation of some neural disturbance, owing to the almost invariable occurrence of the disease in a certain part of the cheek. Whether the disease is some form of mycotic necrosis, as first claimed by Froriep¹⁷ or of bacterial origin, is still an open question. Lingard¹⁸ found in the tissues long, thread-like growths, the individual threads being composed of small bacilli. He reported also finding a bacillus similar to the above in a calf suffering from what was apparently cancrum oris. Ranke¹⁹ found in noma tissues certain cocci, the specific nature of which have never been recognized. As a result of his findings, Ranke concluded that different forms of gangrene resulting from noma unquestionably occur in children who have a tendency to disease of this character; i. e., without infection from contact.

The endemicity of the malady in public institutions coupled with the seeming choice of the disease for localization on the mucous membrane of the different openings of the body suggest that the causation may be looked for in the invasion from without of microorganisms. Schimmelbusch,²⁰ as a result of his bacteriologic study of one case, found bacilli often in shortened pairs and sometimes in long filaments.

An extremely good digest of the subject of noma is given by Blumer and McFarlane,²¹ who have reported a series of sixteen cases occurring in epidemic form which

they accepted as undoubted noma. As a result of their investigations, they concluded that the disease began as a simple infection and terminated in a mixed infection, and that the mixed infection was not always due to the same organism, but was usually due to a long organism of a leptothrix type. In a most elaborate and highly interesting treatise on noma by Weaver and Tunncliffe²² is incorporated the results of the bacteriologic researches concerning noma as undertaken by Grawitz, Bartels, Guizetti, Babes, Zambilovici, Foote, Nikolaysen, Schmidt, Petruschky, Freymuth, Saft, Krahn, Seiffert, Perthes, Braun, Matzenauer, Elder, Ranke, Rosenberger, Brunning, Rona, Buday, Pollard, Herrman, Passini and Leiner, Walsh, Sailer, Strueh, Schimmelbusch, Longo, Hofmann, and Kuster. In the majority of the cases of the above observers the superficial parts of the slough showed mixed cultures; in the deeper parts, the sections often showed almost a pure growth. The organisms most frequently found were slim, thread-like bacteria, decolorizing rather easily by Gram and not growing on ordinary media.

REPORT OF CASE

History.—June 4, 1910, I first saw Mrs. E. M. F., a housewife, aged 59, whose family history was of no material value. The patient was the mother of eight children; had one miscarriage; menstruated at 14; climacteric reached at the age of 43 without any untoward results. For the last eighteen or twenty years previously the patient had had very poor health, being almost constantly ill with gastric and intestinal disturbances. As the patient tersely stated it: "All my life I have been a sufferer from loose bowels and heartburn." Within the past five years the patient has been afflicted with articular rheumatism. Queries as to previous treatment revealed the fact that during the previous five months the patient, at seven different intervals, had taken large amounts of calomel in broken doses. She declared that her former physician advised her when feeling unwell to take a dose of calomel. A daughter of the patient aptly termed her mother a "calomel fiend."

Examination.—Patient looked at least ten years older than she really was. She was wrinkled and emaciated, and bore all the earmarks of a woman who had had a strenuous struggle with life. Evidences of insufficient nourishment and hard work were most striking; personal hygiene very bad; living environments wretched. On going over the patient very thoroughly I found that I had to deal with a chronic interstitial nephritic condition, complicated with endocarditis.

Treatment and Course of Disease.—As it was apparent that the patient's span of life was drawing to a close, only simple palliative measures were resorted to. Under this mode of treatment the patient maintained her balance for a fortnight, when early on the morning of the fifteenth day of her illness my attention was directed to a slight swelling on the left side of the face. Examination of the mouth revealed an ulcerated patch on the mucous membrane of the gum covering the outer aspect of the mandible. The teeth were carious and disgustingly dirty; the lower teeth were artificial; there was a limited amount of exudation. The odor was most fetid. I prescribed an oral antiseptic and promised to call again the same evening.

On my second visit the swelling had increased and had extended more to the middle line. There had been a slight hemorrhage at this time and the exudation had increased. Within the short space of seven hours the ulcerative process had become a dark and dusky red, which was soon followed by a circumscribed area of dead exudate. There was also a slight sloughing which the patient was trying, in vain, to expectorate. It was at this juncture that Dr. Stephen W. Stafford kindly consented to see the patient with me. As a result of his examination, he concurred in my diagnosis and immediately pronounced the condition cancrum oris. The disease was now spreading rapidly and there was a well-marked line of sloughing tissues following the imprint of a set of

17. Froriep: *Chirurgische Kupfertafeln*, 1844, p. 458.

18. Lingard: *Lancet*, London, 1888, ii.

19. Ranke: *Zur Aetiologie und pathologischen Anatomie des necrotischen Brandes*, Jahrb. f. Kinderh., Leipzig, 1888, xxvii, 309.

20. Schimmelbusch: *Deutsche med. Wochenschr.*, 1889, xv, 516.

21. Blumer and McFarlane: *Am. Jour. Med. Sc.*, xxii, 527.

22. Weaver and Tunncliffe: *Jour. Infect. Dis.*, 1907, iv, 8.

false (lower) teeth. The frenum of the lower lip soon became involved and it seemed that that portion of the mouth between the outer mucous membrane of the alveolar process and the inner side of the cheek was literally rotting away. The odor at this time had become so bad that it appeared to invade the whole household. It was most disgusting and the deodorants employed proved of little value.

On the morning of the third day the frenum of the lower lip began to split from within outward and the whole of the inner aspect of the mucous membrane of the left cheek was one black, greenish and garnet mass, attended with a slight hemorrhage. The tongue was swollen beyond its natural size and deglutition was very difficult. Speech was painful and required great effort.

Later the same day the tissue destruction had progressed to such an extent that the inner angle of the cheek began to break down. The patient's condition at this time was most serious and death was but a question of a few hours. There was profound prostration, feeble pulse and retention of urine. The patient's condition rapidly grew worse, she soon went into a typhoid state and died that night at 11 o'clock. Only twice during her illness did the temperature go beyond 100 F.

June 21, tissue scrapings and oral secretions were submitted for bacteriologic study to Dr. William H. Seeman, who made the following report:

Slide 1.—Leishman (Romanowsky) staining. *Spirochæta dentium* present in considerable numbers. Great quantities of pus cocci. Organism of a leptothrix type plainly discernible. Fusiform bacilli present, but not to any marked extent.

Slide 2.—Weak carbol-fuchsin staining. Innumerable spirochetes easily demonstrable. Bacilli also present which seem to be the early stage of some form of spirillum. In no way does the latter resemble Vincent's spirilla.

Anaerobic culture yielded practically negative results. Pus cocci and non-pathogenic bacilli found. The culture failed to show any thread-like organisms.

It was fortunate that during the above examination Dr. Seeman had at the same time submitted to him secretions from a case of Vincent's angina and he made comparative examinations between the specimens of the two cases. As a result of this comparative bacteriologic study, Dr. Seeman was able to determine very distinctly the morphologic and cultural differences between the causal organism of Vincent's angina and the organism which probably causes noma. The essential difficulty of recognizing a certain and definite causative organism is usually obscured by the great number of bacilli of uncertain character which are always found in nomatous tissue. Moreover, it must be noted that in the mouth of the normally healthy person bacteria are always present in large numbers and the following have been found: *Spirochæta dentium*, *Spirillum sputigenum*, *Bacillus buccalis maximus*, *Iodococcus vaginatus*,²³ *Leptothrix buccalis innominata* and the *Leptothrix buccalis maxima*. In all likelihood, they are non-pathogenic. The existence of dental caries always proves a most favorable nidus for microorganisms. Aside from this, it is interesting to note that the following pathogenic microorganisms may also be present in the mouth under apparently normal conditions: the *Diplococcus pneumoniae*, *Diplococcus lanceolatus*, *Micrococcus septicæmia sputi*, *Streptococcus pyogenes*, *Micrococcus tetragenus*, *Bacillus crassus sputigenes*, and the *Bacillus coli communis*.

From a consideration of the case just recorded it is possible to obtain a clinical picture of the disease. The diagnosis does not present difficulties when the disease is well established. However, noma, when affecting the genitalia, might be mistaken for ulcerating granuloma of the pudenda. The symptoms are similar in some respects

to those encountered in infectious submaxillary angina (Ludwig's). In fact, in many cases it is impossible on clinical grounds alone to distinguish angina from noma. Ludwig's angina differs from noma in that it displays more of a tendency to destroy loose tissues and avoid the denser ones. It often happens that Ludwig's angina terminates in gangrene.

The symptoms produced by noma in the aged do not differ materially from those seen in childhood. It can be readily seen that the predisposing factor in the foregoing case was a frail human economy whose vital powers were lowered by chronic renal and cardiac disease, to which were added insufficient nourishment and vitiated environments. It may be asked if the heavy calomel medication just prior to the last illness may not have played its part in aiding the condition. While it is possible that there may have been a condition bordering on mercurialism, it must be borne in mind, says Reisman,²⁴ that mercury as a causative factor in this condition has been absolutely disproved, quoting Coates²⁵ in support of this view. Dr. Benjamin Rush²⁶ cites an instance in which a colleague of his, Dr. Stewart, was in the habit of giving 20 grains of calomel three or four times a day and continuing this as routine treatment for five or six days without any ill effects. It is to this same Dr. Stewart that modern syphilographers owe the idea of mercury medication by inunction, as he was the first to compel patients to wear socks containing mercurial ointment in the treatment of luetic conditions. Dr. Stafford, who saw the case in consultation, also rejects the idea of calomel medication having anything to do with the production of the disease.

The condition of the patient at all times precluded the possibility of ever doing anything except to institute palliative measures. The drastic treatment so necessary for a cure in these cases was entirely out of the question.

CONCLUSIONS

1. Noma does exist in the aged, but its occurrence is remarkably rare.

2. Noma in the aged does not differ materially from the disease as seen in childhood. The supposed predilection of this malady for the female sex is due alone to the fact that the female genitalia offer a more suitable field for infection.

3. No certain and definite organism has been definitely established as the causative factor; the indictment of the long, thread-like bacillus, so frequently regarded as the cause of the disease, is on purely conjectural grounds. While in the realms of the hypothetical metabolic retrogression brought about through previous illness, vitiated environment and bad hygienic conditions may effect a change in the pathogenicity of the previously harmless microorganisms of the mouth to such an extent as to cause an overwhelming of the tissues of the cheeks and jaws, and thus produce gangrene and a condition of general sepsis.

4. Infectious submaxillary angina may be the forerunner of noma, and once the gangrene arises it would be difficult, indeed, to say whether the condition was due to Ludwig's angina or noma. This, too, despite bacteriologic observations, as it is more than probable that the two conditions could be coexistent. It is highly probable that noma is a process caused by more than one type of organism.

602 Perrin Building.

24. Reisman: Osler's Modern Medicine, ed. 5, 48.

25. Coates North American Med. and Surg. Jour., ii, 1826.

26. Rush, Benjamin: Medical Inquiries and Observations, 1815, iii, 23-4, Philadelphia.

23. *Iodococcus vaginatus* (Miller) is found in unclean mouths; gives a peculiar reaction with iodine, the sheath being stained yellow, the cocci dark blue.