which causes him the least inconvenience, to resort to the operation. Great aseptic care must be used to guard against accidents. With such care I see no reason why the operation for the cure of inguinal hernia should not be practically free from danger and the percentage of relapses be almost *nil*.

GENERAL AND SPECIAL TREATMENT OF TUBERCULOSIS.

Read before the Dodge County, Wisconsin, Medical Society,
at Waupun, Wis.

W. E. OWEN, M.D.

FOX LAKE, WISCONSIN.

It is a source of pleasure to state in the words of an authority, Solis Cohen, that tuberculosis of the lungs is a curable disease. This authority, however, believes that the statement must be considerably modified with reference to the outcome of every case. We all realize from painful experience how utterly inefficient were our past resources in the arrest of these cases. There is no disease within the realm of medicine that gives the physician more prolonged and painful anxiety, because, in the first place, it is a pathologic process, generally occupying a lengthened period of time, whether in the direction of recovery or death; and in the second place, because we can not weigh in the balance the amount of potential energy possessed by the victim and the degree of vitality of the invading bacilli. We are thus surrounded by a cloud of uncertainty.

It may be well to generalize the situation by the statement that tuberculosis is primarily a local manifestation of a degenerative process, which may be limited to a few cells, or lobules of the lungs, or diffused, so to speak, throughout the economy, the characteristic tubercular process being due to some form of lesion whereby the active principle (the bacilli) gains a foothold through that gateway or entrance.

Degeneration of cells is produced by prolonged disregard of natural laws by predecessors, or promoted in the direction of depression, privation or excess. Anything either within or without the economy that is calculated to reduce the physiologic resistance of individual cells prepares the way for this dreaded enemy. We may also state the opposite, that everything within or without the economy calculated to maintain the standard of physiologic resistance are ideal conditions to ward off danger. With normal resistance of individual cells the microbes will come and go harmlessly. After the invasion of the tissues by the bacilli the struggle for existence continues, it being then a case of the survival of the fittest.

I have been deeply impressed of late by the suggestion, that we should examine the lungs of our patients more thoroughly and more frequently, especially those who have chronic dyspeptic symptoms and subject, by least exposure, to catching cold. When there is a persistent yellowish deposit on the tongue, after excluding malaria and rheumatism, we should suspect incipient tuberculosis; particularly when there is the least evidence of a predisposition in the family history. These symptoms, associated with indications of parenchymous change, constitute the so-called pre-phthisical state. It may be defined as a disposition of individual parent cells to produce a progeny of lowered vitality, brought about by conditions entirely outside of the cells; conditions which

produce mal-nutrition of cells. We shall not err very much by saying that in most of these cases the great battle for the supremacy between cell and bacillus has already commenced.

It is evident that to overcome this morbid state of mal-nutrition, our efforts should be directed towards making the environment as physiologic as possible. Our hope of relieving suffering humanity rests upon this principle generally speaking.

Nature has endowed all cells with wonderful recuperative forces, and providing that the parent curse of many generations does not rest upon any biologic cell, nor the surroundings be unfavorable, all living protoplasm by inherent forces will evolve from a lower to a higher state of physiologic vigor. I would suggest that these cases be treated, in all detail, as tubercular, even in the absence of any positive local symptoms. The treatment that promises any degree of satisfaction must be general. We must resort to all measures to improve nutrition, to bring the economy into harmony with its surroundings. We can not call to our aid any one specific, and I doubt if we ever shall; for independent of the invasion of bacilli are the degenerate cells. I feel, however, enthusiastic over the possibilities within reach of all. Our hope of a successful issue rests not upon directly fighting bacilli; but upon fighting degeneration of individual cells; upon increasing potential energy, modifying their surroundings, bringing the cells into harmony and equilibrium with environment. If we succeed in our efforts we may venture to disregard the bacilli, for we shall have succeeded in changing a favorable to an exceedingly unfavorable soil for their development.

A very essential beginning with these cases consists in ridding our patients of all disobedience and anxiety by laying the whole matter intelligently before them. In no other manner can we successfully protect the well and gain the patients harmonious co-operation. Inspire them with confidence and hope. Impress upon them the fact that by prolonged obedience to natural laws, rests their future welfare.

Cheerful surroundings are an essential factor to a successful issue; they aid nutrition reflexly through the mind. Prompt and thorough elimination of the products of normal and morbid tissue metabolism must be maintained. All the emunctories must be kept actively at work.

We will first take up the skin. By warm baths we increase its eliminative function; the superficial capillaries are filled and the lymphatic circulation stimulated so that subsequent inunctions of cod liver oil will be beneficial for nutrition and protect from cold. By cold bathing we stimulate the nervous system, and also train the economy to adapt itself to sudden changes of temperature without producing morbid reactions. Frictions after the cold bath will stimulate the capillary and lymphatic circulation.

The urinary secretion should be under constant observation, particularly its specific gravity. The specific gravity is usually increased. It is an indication for the administration of more fluids. A patient may take with advantage artificial lithia water or sterilized soft water, or better still, distilled water.

Of the alimentary organs, the liver demands the closest attention. Its influence for good or bad results is very great. It is called the graveyard for

blood cells. When there is a hard fight within the economy we have reasonable evidence that many phagocytes are killed in the encounter. The cells break down in the liver, and thereby increase the specific gravity of the bile. The demands upon the liver are frequently beyond its capacity for work. These cells, however, must be removed or the products of their disintegration will cause auto-infection.

These products, leaking, as it were, into the blood, make a very unphysiologic environment for the living phagocytes. Occasional minute doses of calomel, or preferable, as indicated biniodid of mercury, should be administered to reduce the pressure in the bile ducts. Free use of hot water will reduce the specific gravity of the bile. It may be well to stimulate the parenchyma by the usual remedies. A liver that has a normal capillary and biliary circulation will not let poisonous alkaloids leak (percolate) into the blood. Any abnormality in this direction causes passive pressure in the stomach and bowels that will retard digestion and assimilation. We must guard against prolonged retention of intestinal contents. Careful attention to the alimentary canal and its tributaries will result in greater capacity to digest food and prepare it for oxidation and assimilation. We all realize that the demand of the economy for abundant nourishment is very great in all these cases. Occasional flushing of the colon with sterilized water is worthy of attention. We must endeavor to keep up a vigorous circulation by vaso-motor tonics, systematic exercise, and massage.

Special attention should be given the nasal cavities; all obstruction to free admittance of air must be removed. They must also be kept scrupulously clean. We now come to the most important of our consideration of the treatment of tuberculosis of the lungs. It is not what shall we do with the bacilli, for this is a secondary matter, but the all important and vital question of how shall we maintain, restore or increase the capacity of the lungs? No part of the treatment has been more unsatisfactory in the past, and indeed, no part more disregarded. It is a fact that the largest majority of people with acquired tuberculosis of the lungs have not learned one of the greatest lessons in life, that of physiologic breathing.

Collapsed vesicles, hyperemia and lymph stagnation are nature's punishment for a disregard of her endowments. The decree of nature is, that a neglected organ will atrophy and degenerate.

It is an easy matter for plugs of mucus to settle in the vicinity of collapsed vesicles and there decompose. These mucous masses under normal conditions are expectorated. Wherever we have decomposing organic material there we shall find a plentiful supply of micro-organisms. It is a practical question to ask, how shall we open up these vesicles and relieve the hyperemia and lymph stagnation? It is a source of great pleasure to bring to your attention an apparatus, or instrument, called by the manufacturer, a pneumatic dilator, the principles of which are based upon the most sound philosophy.

Dr. Hawley describes it as an apparatus for preparing and administering nebulous medication by a process of forced dilatation, the basis of which is the discovery that the lips will act as a safety valve and that, into any tube, cavity or valve or air cell into which air can be forced by high pressure, the nebulous medicament prepared by this apparatus, can be forced with absolute safety, thereby enabling the

physician to use agents that are capable of preventing or controlling disease.

This discovery makes an important epoch in practical medicine. We should all grasp these principles with ecstasy for they tear open the thick clouds of despair that surrounded the wretched victims of this disease. I extend to it my fondest hopes for success, as it is destined to save many lives, of those otherwise doomed to an untimely grave. The practicability of the apparatus and process is self-evident. The principle is sound logic and not theoretical uncertainty. A full and elaborate illustrated description of the apparatus and process by Hawley has been published in the JOURNAL OF THE AMERICAN MEDICAL ASSOCIATION.

But I have brought the apparatus here for your inspection, it will speak more eloquently for itself than any words of praise from me. As a prophylactic measure it is invaluable. One in health has but to experience the feeling of well-being by a single dilatation in order to appreciate its capabilities. There is absolutely no danger of rupturing either vesicles or tympanum, as the pressure is uniform, the lips giving perfect protection. By it the cardio-vascular and the pulmonary capacities are greatly increased, and to use the words of Solis Cohen, "resulting in increased function, increased elimination of waste products of function; with increased buoyancy of spirits which is no mean factor in treatment; in a word, heightened vitality."

That the capillaries throughout the body are wonderfully influenced is evident from the fact that the cold hands and feet become warmer. Williams also believes that dilatation of the alveoli in the neighborhood of the various lesions produces an important localizing effect and thus tends to prevent the spread of tuberculosis by secondary infection, from a caseous center or from a secreting cavity with inflamed walls, while the emptying of the vessels and of the capillaries of the region through alveolar dilatation leads to the absorption of caseous and cretaceous tubercular deposit. There is no more ideal treatment to prevent hemorrhage or control it after its occurrence, as it relieves the engorged vessels and removes congestion.

The mechanical principle of forced dilatation with air itself, you clearly see is invaluable; with these dilators, however, we can force into the lungs, the finest medical nebula imaginable. Ordinary inhalers for medicated air may be discarded, as we can not reasonably expect the medicament to get past the reserve air left after the most extreme expiration. They are regulated by the patient's will and consequently not satisfactory. By forced dilatation with nebulous medicament we overcome obstructions which offer an impassable barrier to ordinary inhalations. We can easily imagine the rapid diffusion of fine medicated nebula with the reserve air; and thereby attenuating, at least, the micro-organisms. We will demonstrate the explosive effect desired in order to remove the decomposing pulp of mucus. One bottle contains medicament for cough and all manner of bronchial irritation. The other contains the local specific for these cases—free iodine and chlorid of gold with menthol enough to mask the taste of the iodine. We can hardly realize the extent of the absorbing surface of the bronchi.

In addition to local effect, we also get systematic effects from absorption. For enlarged lymphatic

glands and lymph stagnation, the iodides are held in high esteem by general consent.

Above all else, however, we must endeavor to get as much pure air as possible into all these cases, as normal oxygen is the best excitant of the recuperative forces latent in all cells.

I have thought it waste of time to enter into consideration of the numerous remedies as we are all very familiar with their merits.

SOME REMARKS ON TEST MEALS.

Read by title in the Section on Practice of Medicine at the Forty-sixth Annual Meeting of the American Medical Association, at Baltimore, Md., May 7-10, 1895.

BY HENRY SALZER, M.D.

CLINICAL PROFESSOR OF DISEASES OF THE STOMACH, BALTIMORE MEDICAL COLLEGE, BALTIMORE, MD.

Test meals are given for the purpose of testing the digestive and motor power of the stomach, and thus judging of the condition of the mucous membrane and its glandular elements, as well as of its muscular action and its size, for an enlarged, or rather dilated, stomach depends chiefly upon these conditions.

The test meal now almost universally employed is Ewald's test breakfast, consisting of 30 to 50 grams of bread and 300 c.c. of warm water, or tea without sugar or milk.

One hour after this meal a soft, elastic tube is introduced into the stomach, and the entire contents of the stomach brought up, either by attaching an aspirator apparatus to the tube, or by pressing with the hand at the epigastrium, and encouraging the patient to assist this pressure by contraction of the abdominal muscles as it is done for defecation.

Other test meals have been suggested:

Leube-Riegel's consists of 300 c.c. of soup, 60 to 120 grams of meat and 50 grams of bread with 250 c.c. of water, and either the contents are aspirated or pressed out four hours later, when we expect the height of digestion, as we do one hour after the bread and water meal is taken, when it ought to be entirely empty, if both digestive and motor power are good.

Klemperer advises milk and bread, and two hours later, aspiration. The same time is recommended after the test meal of Germain See, which consists of 60 grams of scraped meat and 30 to 50 grams of bread.

Jaworsky's test meal is very complicated and requires a complete cleansing of the stomach by lavage, a meal of the whites of one or two boiled eggs and 100 c.c. of water, and aspiration every quarter hour for one and one-fourth hours, when the aspirated water becomes clear and acidulated and possesses digestive properties.

Boas advises as a test supper for atony a light meal late at night and lavage next morning, when the normal stomach is empty, while the atonic is not.

It is admitted that every one of these tests is useful, and that every investigator may reach good results with the method he prefers. The food obtained is in a state of advanced digestion, and permits the corresponding conclusions by inspection of the whole contents, by examination of the residue of the filter, and by chemical examination of the filtrate, which represents a true stomach juice at the height of action.

The Ewald breakfast gives these factors as well as the others; its use is very simple, and the masses we have to deal with are very clean and easily distinguished. Yet the disadvantage connected with this,

as well as with other test meals heretofore used, is the necessity of obtaining the entire contents of the stomach before one can judge if the muscular functions are normal. In health the entire residue after one hour should not exceed 20 to 50 c.c.

A further disadvantage of Ewald's test meal is, that it demonstrates the functions of the stomach in regard to only a very limited amount of work. For instance, Honigmann as well as myself, have found that with a more complicated test meal, *i. e.*, Leube's, the results with the same person showed marked differences. No impaired condition was found with the Ewald test meal, yet when a Leube test meal was used, the result was strikingly different.

Ewald and Boas advise in doubtful cases, the control test with a Leube-Riegel meal. These two factors, the necessity of a controlling test meal, and especially the importance of gaining by aspiration or pressure the entire contents of the stomach, is a great annoyance in private practice. It is very often the case that the patient is available only for a limited time and that repeated sittings can not be had to accustom him to the use of the tube; furthermore, by expression or aspiration parts of the mucosa are easily drawn or forced into the fenestræ or eyes of the tube, especially with patients not used to the manipulation, a factor which will be very unpleasant where peptic ulcer may exist, and where you may be especially anxious to gain an idea of an existing hyperchlorhydrie or at least hyperacidity. This objection can be made equally against the Einhorn bucket, which is very apt to bring up pieces of the mucosa, and is certainly not swallowed and removed more easily than a very soft, thin, elastic tube.

The following simple modification of the present test meals does away completely with all these inconveniences, and has been employed by me exclusively for the past three years to my greatest satisfaction.

I make the patient take early in the morning 30 to 50 grams of cold roast, free from all fatty and skinny parts and cut into small pieces one-eighth of an inch wide and about half an inch long; further 250 c.c. of warm milk, or milk and water, and 30 to 50 grams well-toasted wheat bread (dark-brown toast) or Zwieback. Four hours after this meal, 30 to 50 grams of stale wheat bread without crust or toasted particles, and 300 c.c. of warm water, or tea without sugar or milk. One hour later I introduce a thin, soft, rubber tube with two eyes near the lower end; this tube is inserted to the depth of from 50 to 60 cm. to reach the bottom of the stomach. In the case of tall persons, and of those in whom a splashing sound may be detected below the navel I prefer to insert it 60 cm. The tube is allowed to remain a few seconds, until some retching takes place, when it is slowly and carefully withdrawn, and in the case of great distress even without waiting for any contents of the stomach to appear at the upper end; if without any effort some flows out and a larger quantity can be obtained, so much the better. As soon as the stomach end of the tube leaves the mouth it is held upward and over a porcelain dish; by blowing gently into the upper end a sufficient quantity of the stomach contents (generally from 3 to 6 c.c.) is obtained to enable one to make all required tests. You see at a glance whether residues of the first meal are present or not. If there is only a milky, chyme-like addition to the bread and water, and no undigested casein, meat or toast, you know that the *motor* but not the