Klauber, E. EDEMA OF THE OPTIC DISK IN BRAIN INJURY. [Klin. Mbl. f. Aughkl. Apr., May, 1918.]

Klauber reports upon two groups of cases of injury to the skull comprising 72 cases each. The complication in the first group consisted in injury to the bones of the skull and of these 2.8 per cent. of the cases had papillary edema, two of these cases proving fatal. In the other group the dura mater was punctured. 32 per cent. of these died, 30.6 per cent. had papillary edema. The cause of the edema is believed to be in an endocranial stoppage of the lymph and it occurs in infected wounds of the brain chiefly in wounds of the back of the head. Klauber considers it "in a certain measure a prolapse of the brain into the viscera of the eye." He examined 6 cases histologically. Only a smaller number of these showed infiltration of small cells into the papilla. There was edema in the optic nerve trunk decreasing from before backward. Klauber does not consider the subarachnoidal dropsy in the ampulla and the descending perineuritis and interstitial neuritis in the hinder parts of the optic nerve, which are connected with the terminal meningitis, as essential to the appearance of the papillary edema.

Lewis, F. P. RETINAL HEMORRHAGE. [J. A. M. A., June 15, 1918.]

Dr. Lewis reports a case of retinal hemorrhage. A woman had been suffering from recurrent hemorrhages from the retina and had been under the advice and observation of competent specialists for eight months. Elimination by the kidneys and skin was the chief treatment used. She had no arteriosclerosis; her blood pressure was only 120 mm. of mercury, but she had marked leukocytosis. Her age was only 46 years. Five years before she had an abscess of the antrum which had apparently recovered, though when it was subsequently opened a quantity of sterile pus was removed. Pyorrhea was very evident, and the roentgenogram showed apical abscesses of several teeth. The infected teeth were removed, and cultures made. These showed that the active organism was the *Streptococcus hemolyticus*. The removal of the teeth was followed by cessation of the hemorrhages. In another case of recurrent hemorrhage in a man, aged 70, with a blood pressure of 200 mm., the same cause was suspected, and the same organism found. He was treated by autogenous vaccines, and the hemorrhages ceased. Lewis says that retinal hemorrhage is much more frequent than generally supposed and is not dependent on blood pressure. "It may be associated with, but is not directly caused by, arteriosclerosis, albuminuria, diabetes or other diseases involving disturbed metabolism or focal infections. In any of these conditions, protein toxins may be given off. These are easily absorbed into the circulation, more especially when they arise from focal infections that are proximal to the eye. The organism which is most commonly present is the streptococcus, and the form frequently found is that of the *Streptococcus hemolyticus*. The hemorrhage thus

produced is not the result of force from behind, but is due to softening of the tissues of the capillary or arterial walls by the local action of this toxin, thereby producing lysis." This, the author says, is a new conception as regards a cause of retinal hemorrhage, but it establishes a definite etiology of the condition and that is of value. The differential character of the toxins given off by the different strains of bacteria has not yet been studied, but it is known, however, that several varieties produce hemorrhage, and Wells considers it quite probable that of the chemical agencies causing hemorrhage bacterial products are the most important, and Lewis thinks that changes in blood pressure are practically negligible as causes of retinal hemorrhage. This adds also to our understanding of hemorrhages in other parts of the body. In cerebral hemorrhages it is more likely, from whatever source it comes, a protein poison is responsible, weakening the endothelium and causing rupture.

Wallis, G. F. C. OPTIC NERVE, CHIASM, AND SPHENOID BONE. [Practitioner, Jan., 1917.]

The author opposes the orthodox teaching, still largely repeated in the texts, that the optic chiasma occupies the optic groove of the sphenoid bone. As this question has a bearing on many pathological conditions, he has restudied the anatomy of this region in a number of post-mortem subjects. In only one of the eleven bodies examined was the chiasma approaching the more generally accepted position. In this subject, a female aged 93, more than one-half of the chiasma rested upon the optic sulcus and the olivary eminence. The author concludes that while the chiasma does occasionally rest on the optic sulcus, it is nearly always posterior to it. As to the antero-posterior diameter of the chiasma the smallest dimension in the subject examined was 7 mm. and the greatest 11 mm. Measuring the length of the intracranial portion of the optic nerve he found the shortest dimension to be 7 mm. and the greatest 12 mm. The diameter of the intracranial portion of the optic nerve is commonly about 4.5 mm., but may reach 7 mm. The length of the optic canal is usually extremely small (optic foramen), but sometimes the optic nerve is encased in a cylindrical canal of considerable length.

These observations, though based only on a few cases, show the bearing of sphenoidal sinusitis on affections of the optic nerves and chiasma with the consequent field involvement. In case the chiasma happens to be situated almost wholly in front of the pituitary body when that organ is diseased, it may possibly explain those cases where the symptom of bitemporal hemianopsia is absent.

Koepppe, L. NIGHT-BLINDNESS. [Mün. med. Woch., April 9, 1918.]

Hemeralopia, by reason of its present importance in military medicine, is restudied by this author. In contrast to earlier views which at-