growth may be removed if you will circumscribe it with sufficient amount of quinine and urea solution concentration of one-fourth of 1% infiltration. If any nerves are met with they can be blocked off with 1% novocaine solution. The dissection can be done as well or even better under local anesthesia, because the time element is no essential with reference to the patient's safety.

Dr. Urban Maces, New Orleans, La.—I would hesitate to perform the amputation of a carcinomatous breast under local anesthesia. Having been drilled in the clinic of Dr. Matas, of New Orleans, I have adopted the neuro-regional methods in preference to the pure infiltration anesthesia, and have performed an endless variety of operations with this method. Only recently I have had the opportunity of doing a double decortication of the kidneys upon two patients with paravertebral anesthesia.

In hernia operations we have adopted the neuro-regional technique suggested some years ago by Cushing and Bloodgood. After studying a series of cases where pure neuro-regional methods were used and a like number of cases where infiltration only was used, I found that wound healing was materially influenced in the latter. In some instances almost 60% of the infiltration cases showed infection to a greater or less extent.

It is our custom to use 0.5% solution of novocaine with adrenalin, 5 m. to the first and three minims to each succeeding ounce.

Dr. Brookshire.—Please state the amount of Novocaine. Do you use it in any definite quantity?

Dr. Cole (closing).—I purposely left out nearly everything except the simple photographs because you are all familiar with the injection methods. To a large extent, it has become a routine with me in acute surgical abdomens with pus. In acute gall bladders, appendicitis, etc., undoubtedly a two-step operation may mean the saving of the patient.

Not knowing that Dr. DuBose had read a paper similar to mine, I was certainly not familiar with the fact that the doctor had reported these cases. However, I do not claim originality, but simply reported one or two cases that had not been published in this country before.

For blocking, a 1% or 2% solution, limiting the amount to 5 to 10 cc. is best. I use 3 drops of adrenalin to each ounce of anesthetic solution for infiltration. It is astonishing the amount of solution that can be used. I have used as high as 26 to 28 grains, where the incision was open, and a great deal of the fluid was released.

The preliminary use of scopolamin, 1/150 grain, and of morphine before operation is often essential.

Fortunately the aged patients are very good subjects for local anesthesia; and there can be no question that the procedure is much more valuable in the old adult than in the young.

I published an article upon this subject in the Florida State Reports in which I called attention especially to the different methods.—intra-neural injection, blocking, or combination of local anesthesia with the nerve blocking.

Dr. DuBose.—I would call it twilight sleep, plus nerve-blocking.

A NEW TREATMENT FOR CHANCROIDS

BY GEO. R. LIVERMORE, M.D.,

Professor Genito-Urinary Surgery, University of Tennessee; Genito-Urinary Surgeon, Baptist Memorial and City Hospitals, Memphis, Tenn.

The many remedies in use for the treatment of chancroids is conclusive evidence of the inadequacy of any one of them. It is now a generally accepted fact that chancroid is caused by the strepto bacillus of Ducrey, but that contamination with other bacteria is practically always present. In order to effect a cure it is essential to eradicate all infection. The actual cautery, nitric acid, etc., will often do this successfully, but if they fail a large amount of necrotic tissue is left that is both a protection and a fertile field for bacterial growth. A drug that will destroy the infection and not injure the tissue is ideal and the one that I wish to bring to the attention of the profession fulfills these requirements. This drug is Silvol, a new protein silver compound containing about 20% silver. It occurs in the form of dark brown lustrous scales, is hygroscopic and readily soluble in water.

The method employed in treating chancroids with Silvol, consists in cleansing the chancroid thoroughly with warm sterile water and then covering the entire ulcerated area with Silvol crystals. A gauze dressing and bandage are then applied and not removed till the following day. If upon removal of the dressing, the chancroid has not lost its virulent appearance, a second application is resorted to. When Silvol is applied to a chancroid there is a marked stinging sensation, but it is usually not severe and never lasts very long.

Although Silvol is not a specific and it is not suggested as the one sure shot, it is by far the best single remedy that I have used for the treatment of chancroids.

Exchange Building.

AUTHORS' ABSTRACTS.

(See page 748)