formation in the connective tissue between the muscle bundles. A number of authorities conclude that the bone is a product of detached pieces of periosteum, others that it rises from the connective tissue cells between the muscle bundles. Clinically the condition is not difficult to recognize, especially with the aid of the x-ray. However, when the new shadow rests directly on the shaft of the neighboring bone it will be difficult to differentiate the ossifying myositis from a condition called

Fig. 12.—Microphotograph of a section of an adamantine epithelioma, involving chiefly the alveolar border of the lower jaw. In this section the basal columnar cell and its various morphologic changes toward the center of the alveolus are well shown. (Sent me by Dr. Steensland.)

Fig. 13.—Microphotograph of a section of an adamantine epithelioma of the gum.

traumatic exostosis or ossifying periostitis. Blood cysts have been observed only in the ossifying myositis. Extensive operations are not necessary. One should remove as much of the new bone production as possible without destroying function. Slight local recurrences of the bone formation are to be expected. Fortunately the amount is never great, gives little or no discomfort, and second operations are rarely necessary. I

do not think there is any relation between the hemorrhagic cysts observed in myositis ossificans and the cysts of the long pipe bones.

As to the question of the origin of the giant cells, recent investigation would indicate that the periosteal or medullary giant-cell tumor is probably an angioma or an angiosarcoma, and that the giant cells are due to budding of the endothelial cells of the vessels in these very vascular tumors. The most interesting publication on this subject is by Friedländer.* Further investigation, however, should be made of this most interesting tumor.

EXPERIMENTALLY PRODUCED GENUINE EPITHELIAL METAPLASIA IN THE STOMACH,

AND THE RELATIONS OF EPITHELIAL METAPLASIA TO CARCINOMA, AS DEMONSTRATED BY CASES REPORTED IN THE LITERATURE.

GUSTAV FÜTTERER, M.D.

CHICAGO.

As the observations on epithelial metaplasia to be reported in this paper were made in a series of experiments on the stomach of rabbits, undertaken for the purpose of studying the relations between mechanical irritation and development of carcinoma, I think it necessary to state this fact, merely to show under what circumstances the findings were made. My views concerning the rôle which mechanical irritations play in the etiology of carcinoma, and how those views have been formed, have been fully explained in former writings, and I can, therefore, refer the reader to them, in order to prevent repetitions, which would make this article too extensive.¹

Some observations of importance have been made which I do not report, because they need corroboration, but they certainly encourage further efforts along the same lines.

In a series of experiments on the stomachs of 94 rabbits, a genuine epithelial metaplasia was found in five cases, only two of which will be described in detail, as a description of the others would cause an unnecessary repetition, for while the findings differ somewhat in all the cases, they are the same in principle. It may also be mentioned that I have specimens of epithelial metaplasia from a former experiment, bringing the total of cases to six.

In each case a piece of mucous membrane of the stomach was resected and hypodermic injections of a solution of pyrogallie acid were made, to reduce the percentage of hemoglobin, and thus prevent the healing of the defect, the edges of which were then exposed to friction by solid food. At first, as in former experiments, circular resections of mucous membrane around the pylorus were made, but the feeding of the animals was not carried out according to orders given, and they all died, postmortem examination revealing very much dilated stomachs filled with solid food, while only liquid food was to be given during the first four days. Not being able to get better care of the animals at the time, I had, in order to reduce the mortality, to resect pieces of mucous membrane near or further away from the pylorus, and later, when Dr W. H. English, to whom I am very much indebted, kindly took it on himself to supervise the keeping of the animals, my results became very much better.

* Archiv f. klin. Chir., 1902, vol. lxxvii, p. 202.

1. Fütterer: Aet. d. Carc., Bergmann, Wiesbaden, 1901, and THE JOURNAL A. M. A., March 15, 1902.

Fig 1.—Case 1. Hematoxylin-eosin stain. Bausch and Lomb, obj. 3 inches, magn. 30. V. Vegetable matter. P. 1, 2, 3, Peg of squamous epithelial cells. Rad. Root of the peg. Muc. Mucosa. Musc. m. Muscularis mucosæ. Msc. Muscularis (hypertrophic). R. Rest of muscularis, not invaded by the peg. S. Serosa.

The microphotographs have been made by Dr. W. H. Knapp of Chicago, to whom I am very much indebted for the excellent work.

Fig. 2.—Case 1. Hematoxylin-eosin stain. Hensch and Lomb, ocular 2, obj. $\frac{1}{2}$ inch, magn. 250. In the middle of the field is found the part marked Rad. (Fig. 1), with higher magnification. P. Peg of squamous epithelial cells. Cyl. Dilated glands, with very high cylindrical epithelium.

METHOD OF OPERATING.

1. The skin over the region of the stomach was shaved.
2. Ether narcosis.
3. The skin was cut with a knife.
4. The muscles were held with pincers, and the abdomen was cut open with scissors.
5. Sterilized gauze was packed around the wound, the stomach was pulled out almost entirely, and gauze was packed all around it.
6. The stomach was opened with scissors on the anterior walls in the pyloric portion.
7. The contents of the stomach were removed with a teaspoon.
8. A part of the posterior wall of the stomach was pushed through the wound with the fingers.
9. The mucosa of this part was cut with scissors and a grooved director was introduced into the submucosa, loosening the mucosa plus the muscularis mucosa from its bed, and then the loosened piece was cut off with scissors.
10. Two catgut sutures, one uniting the mucosa and one the rest of the wall of the stomach, closed the anterior wound of the organ.

11. The stomach was allowed to fall back into the abdominal cavity, the gauze strips were removed, and the abdomen was closed by sutures.

12. The animals operated on received regular injections of 0.14 pyrogallic acid in solution hypodermically at different intervals, as they were needed to destroy hemoglobin. Hemoglobin examinations were made once a week, and all the animals were weighed.

On the whole, I must say that the results, so far as the production of certain forms of ulcers was concerned, were not entirely satisfactory, but I believe that the experience gathered will enable me in the near future to be more successful in this direction. The work to be reported here consists of a series of experi-

globin, 85 per cent. Operation Jan. 8, 1904; died Feb. 1, 1904. January 23, hemoglobin, 60 per cent; weight, 2½ pounds. January 30, hemoglobin, 50 per cent.; weight 2¾ pounds. One hypodermic injection of .14 pyrogallic acid in solution had been made.

In this case, as in Case 2, an infection of the right hand prevented me from performing the postmortem myself. When the stomach was handed to me, shortly after it had been removed from the animal's body, thickening and hardening of the part of the posterior wall of the stomach where the resection had been made, and where a regeneration of the mucosa had occurred, were recognized. In the midst of the regenerated portion could be seen a very small indentation and microscopic examination revealed here the presence of a peg of squamous epithelial cells, as is represented in Figure 1, with low power,

Fig. 3.—Case 2. Hematoxylin-eosin stain. Bausch and Lomb, obj. 3 inches, magn. 30. Muc. Mucosa with elongated glands and round-cell infiltration. Musc. m. Muscularis mucosae. Msc. Muscularis (hypertrophic). S. Serosa. P. Peg of squamous epithelial cells.

ments conducted between Dec. 6, 1903, and April 2, 1904.

REPORT OF THE FIVE CASES OF METAPLASIA.

The five cases of metaplasia recorded in our table have their principal features in regard to origin, the formation of pegs of squamous epithelium and their growth into the wall of the stomach in common, while there are some differences in the development and differentiation of the layers constituting the pegs.

CASE 1.—(Operation No. 32.)

Rabbit of white color; female, weighing 3½ pounds; hemo-

and in Figure 2, showing its origin from the mucosa, with high power.

Microscopic Findings.—With low power an epithelial peg was seen, the upper end of which reached but slightly above the mucous membrane at both edges of the indentation, before mentioned. The epithelial peg had three branches, one in the upper layer of the muscularis propria, and the two others deep in the outer layers of the muscularis propria, through which the main part of the peg had almost penetrated and reached the serosa. The peg was isolated, on all sides surrounded by mucous membrane, and no similar formations were found anywhere. Only in one place was the peg firmly connected with

Fig. 4.—Case 2. Hematoxylin-eosin stain. Bausch and Lomb, ocular 2, obj. $\frac{1}{2}$ inch, magn. 325. Gl. Glands of the mucosa. P. Peg of squamous epithelial cells. Cyl. Metaplasia. The cylindrical epithelium of the glands of the mucosa, changing into squamous epithelium. The nuclei contain a great deal of chromatin and the body of the cells becomes much longer. The glands themselves are dilated. It is of importance to notice that the epithelium lining the glands and undergoing metamorphosis shows perfectly regular arrangement, as under normal conditions, and it can easily be recognized that these cells are the regular columnar cells of the glands and not implanted skin cells, which would fill the lumen of the glands.

Fig. 5.—Case 3. This case was only mentioned, but not described in the text, because it was so similar to Case 1 that it was thought unnecessary. A section from this case was selected for microphotography, because it was thinner and better stained than those of the other case, which, however, showed the same findings. Str. c. Stratum corneum. Str. l. Stratum lucidum (only indicated). Str. gr. Stratum granulosum. D. Change of direction of the nuclei. Str. m. Stratum malpighii. Musc. m. Muscularis mucosae. Str. p. Formation of a papilla.

the mucous membrane of the stomach, and this place has been marked rad., i. e., radix, presumably the root of the peg. The indentation was filled with vegetable matter, the muscularis propria was very much hypertrophied, and the peg itself had, with its ramifications, invaginated the hypertrophic muscularis mucosæ, pushing it almost clear through the muscularis propria, without breaking through it anywhere. The structure of the epithelial peg, which was not perfectly solid, resembled that of the skin in many ways, but there were some important microscopic differences to which attention will be called later.

Figure 5 shows well the resemblance of the structure of the walls of the peg to the layers of the normal skin. The specimen photographed has been taken from Case 3, not because Case 1 did not show the same structures, but because the sections happened to be thinner, and on account of this and their staining were more suited for microphotography.

Figure 5 shows: 1. A stratum malpighii with a dense layer of oval nuclei containing a great deal of chromatin, many karyokinetic figures and long cells, more or less at right angles to the longitudinal axis of the epithelial peg. Upward, the nuclei become more round and more pale, containing less chromatin, and then rather suddenly they appear more drawn out, more cylindrical, and as if drawn by a different current, they change their position, their longitudinal axis, rather suddenly, becoming parallel to the longitudinal axis of the peg itself, thus forming a well-differentiated stratum granulosum. In those three cases I have not been able to find prickle cells, which may be due to their absence or to the thickness of the celloidin sections, and to the fact that the cells of the stratum malpighii lie much more closely together than they do in the normal skin. 2. A papillary body is well developed. 3. The stratum granulosum shows a deeper hematoxylin stain than the upper parts of the stratum malpighii, and between this layer and the stratum corneum there are lighter portions, which seem to indicate a stratum lucidum, although I have not been able to convince myself that such a stratum really exists. 4. There is a sufficiently well differentiated stratum corneum, with considerable keratohyalin and with Gram's method small patches of hornification could be demonstrated in Case 1 and Case 3. In this case, as in the others, some striped muscular tissue was found in the parts of the muscularis of the stomach near to the epithelial peg.

After what has been said, it is clear that the epithelial peg described represents a formation of epidermis in the stomach, but the question arises whether this is really an epithelial metaplasia, or rather a development from an aberrant germinal deposit or an ingrowth of epidermis or an implantation.

As to a development from a germinal deposit, by looking at the question in general, it would seem as if such an origin can never be excluded, because if, for instance, a single squamous epithelial cell is found in a field that should only be lined with cylindrical cells, then this one cell may represent an embryologic deposit, and things would become more complicated, if more than one of such cells were present.

In this case the epithelial peg was found in the center of the regeneration filling the artificial defect. If there had been a germinal deposit, it would have been removed at the operation, and as the peg was absolutely isolated, no ingrowth from a possible germinal deposit at the edges of the artificial defect can have come into play. It may be said here that in Case 2, Figure 4 shows clearly the development of metaplasia from the columnar cells of the gland, and I consider those findings alone as conclusive. An ingrowth of epidermis is absolutely to be excluded, as there was no fistula in any of the cases, as the metaplasias were found in about the middle of the stomach, and as each one was perfectly isolated.

There remains the possibility of an implantation during the operation.

Against the possibility of an implantation the following reasons can be advanced:

1. The possibility of a successful transplantation of detached normal epithelial cells has never been demonstrated. I myself have made use of every occasion for many years to transplant normal epithelial cells, under most favorable conditions, in hundreds of cases, and neither I nor anyone else has ever observed a further development of the cells transplanted. If these were implantations, they would be the first successful cases of implantation of detached epithelial cells on record, and that would be a matter of great importance in view of the sudden downward growth of the transplanted cells.

2. The description of the method of operation employed in our experiments seems sufficient to show that an implantation of epidermis cells can not have taken place, and if it should have occurred it would have been under the most unfavorable circumstances. The cells would have been transferred to a bleeding surface, and been subjected to the mechanical influences of the food, the contractions of the stomach and digestive influences of the gastric juice. If cells transplanted under most favorable conditions do not thrive, how can they develop under such most unfavorable conditions? I have often tried to transplant epidermis cells into the submucosa of the stomach, but have never been successful.

3. Sixty-nine of the animals operated on died during the first week after the operation, and as no microscopic examinations of their stomachs were made, we have five metaplasias out of 25 cases, which would be a very high percentage, indeed, for implantation.

4. It has been mentioned that the development of the layers forming the pegs differed very much, and that would not be the case in implantations.

5. Figure 2 and Figure 4 show plainly the development of the pegs from the epithelium of the glands which, particularly in Figure 5, shows its normal arrangement along the walls of the gland, without any disorder in position. If an implantation had occurred, the implanted cells could never show such an arrangement, but they would fill the lumen of the glands. I have specimens of an implantation into the glands of the mucosa of the stomach from a carcinoma of the esophagus, and that gives entirely different findings.

6. Hornification is only found in very small spots. An implantation would very probably show more of it.

7. But very few prickle cells have been observed. Many specimens were examined before a few were found. An implantation would have furnished more of them.

8. In the unstained specimens no pigment was seen.

9. If an implantation of skin had occurred, the vigorous down-growth from the very start of the development of the pegs, occurring in every case, would very much need an explanation.

CASE 2.—(Operation No. 27.)

Rabbit; female, brown, weight and hemoglobin at time of operation not recorded. Operation Jan. 7, 1904; died Feb. 19, 1904.

	Per cent.	Weight.
January 23 hemoglobin.....	45	3¼ pounds.
January 30. hemoglobin.....	38	3¼ pounds.
February 6. hemoglobin.....	45	3¼ pounds.
February 13. hemoglobin.....	38	3¼ pounds.

During the last six days no weight has been recorded, and it can not, therefore, be stated whether any loss of weight occurred during this period.

When the freshly removed stomach was handed to me I saw

a strand 7 mm. long that had been cut off from its continuation. The strand was round, had a diameter of 4 mm., was of rather firm consistency, and was filled with a whitish mass. At a point corresponding to the insertion of the strand into the wall of the stomach, toward the middle of the organ, on its posterior side, a prominent and hardened portion of the mucosa was found on the inside of the organ, which represented a healed defect, Figure 3. The strand was cut off close to the stomach, for imbedding and cross-sectioning, but was lost, and, of course, I have no idea where its continuation in the animal may have ended.

The examination of the stomach specimens revealed another metaplasia very similar to the one described, but much more developed. A large epithelial peg had pressed the muscularis mucosæ through the greater part of the muscularis propria, and the former and the outer layers of the latter, plus serosa, had, no doubt, formed the walls of the hollow strand that was filled with squamous epithelial cells. Figure 3 shows the peg on its way through the hypertrophic muscularis, and Figure 4 demonstrates beautifully the metaplasia of the columnar epithelial cells of the glands, and I consider those findings alone sufficient proof against an implantation.

Microscopic Findings.—The development of the epidermoidal layers does not show nearly the same differentiation as in Case 1, yet a papillary body, a stratum malpighii, and a stratum corneum are sufficiently well developed.

Case 3 is very much like Case 1, and Cases 4 and 5 are more like Case 2.

In Case 4 a strand was again found coming from the stomach, and here I saw it end between two lobes of the liver. The strand ended blind, and at its end fine adhesions connected it with the liver capsule. The walls of the strand were formed by muscularis mucosæ, the outer layers of the muscularis propria and the serosa, to its very end, and the epithelial cells had not broken through them anywhere.

Concerning my conception of the changes found in all five cases, I would say:

1. All five cases represent genuine epithelial metaplasias, as an ingrowth, an implantation, and a development from a germinal deposit can be excluded. There was no fistula in any of the cases.

2. In all the cases the metaplasia has at once been followed by a down-growth.

3. This down-growth has occurred under considerable growth pressure, as shown by the invagination of the muscularis mucosæ through the greater part of the hypertrophic muscularis propria, and the invagination of the rest of that layer for quite a distance outside of the organ.

4. The metaplasia has been followed by what we may call a practically unlimited proliferation of the squamous cells.

5. Such a down-growth and such an unlimited proliferation of cells are two principal characteristics of carcinoma.

6. We can here use the term metaplasia only for the metamorphosis of the columnar to the squamous cells, for there is no more metaplasia going on in the pegs, the products of metaplasia. I would just speak of atypical pegs in an abnormal position.

7. The epithelial cells have nowhere broken through the muscularis mucosa, and the most characteristic peculiarity of carcinoma, the ingrowth into connective tissue, is absent.

8. What would have become of the strands, filled to the very end with squamous epithelial cells, if the animals had lived longer, can not be said and only further experiments can show it.

9. These are the first cases of epithelial metaplasia, produced by experiment, on record, if we except a metaplasia in the bladder, which Lubarsch mentions, with-

out describing it. *It is to be remembered that structures considered to be characteristic of the ectoderm have been produced from entodermal tissues, a fact which would make it impossible to uphold a sharp division of the germinal layers.*

LITERATURE ON EPITHELIAL METAPLASIA.

In the human being a great many cases of epithelial metaplasia have been described and their careful reading offers a great deal of material for reflection. Particularly the close relation of epithelial metaplasia to carcinoma is strongly emphasized by the observations recorded.

Eye.—In the eye, Michel¹² and Leber¹³ have seen metaplasia of the cylindrical epithelium of the conjunctiva, in connection with chronic conjunctivitis, and in the nose and its accessory cavities, and in the ear metaplasias are found with relative frequency, but they are mostly of a superficial character. In reference to the metaplasias of the nose, the writings of Suchanek, Schuchard, Habermann, Bungner, Kahn and Zerniko will be mentioned.

Nose.—Suchanek¹⁴ found transitional epithelium and squamous epithelium in the secretions from acute rhinitis and influenza-rhinitis, which, according to his opinion, did not come from the bottom of the nasal cavity, or from the introitus narium, but from different parts of the regio respiratoria, and even from the regio olfactoria. In all of these parts, according to Suchanek, patches of epithelial metaplasias are sometimes found in connection with chronic catarrh, and irritations as they occur with different industries.

Accessory Sinuses.—Schuchard¹⁵ points out the remarkable peculiarity of the mucous membrane of the antrum of Highmore to form squamous epithelium under the influence of catarrhal conditions. He also mentions that a hydrops of the antrum of Highmore can lead to the formation of steatomata or butter cysts, and that both probably owe their origin to metaplastic changes of the epithelium.

Habermann¹⁶ describes a cholesteatoma of the right frontal sinus. The patient had a fistula under the right eyebrow, from which pus had been discharged for about nine months, and Habermann believes that an ingrowth of epidermis through this fistula into the frontal sinus had occurred.

Bungner¹⁷ saw a benign, hard papilloma in the upper portion of the nasal cavity. The papilloma showed hornification and had developed from psoriasis nasi of the mucous membrane of the septum. There was also ozena, which had existed since puberty.

Kahn¹⁸ mentions a hard papilloma which was covered with an immense layer of stratified squamous epithelium. The surrounding mucous membrane had not been examined.

Zerniko¹⁹ examined a fibroma of the nasopharyngeal cavity, which was covered partly with cylindrical epithelium, partly with squamous epithelium, and between both kinds a transitional epithelium was present.

Ear.—In the ear Voltolini¹⁰ found large squamous epithelia in the contents of the drum cavity, in 14 cases. Lucae believes that purulent inflammation of the middle ear may, through the formation of granulations, lead to a formation and desquamation of epidermis.

Knapp¹¹ and Habermann¹² report cases of cholesteatoma. The case of the latter author was accompanied by a purulent inflammation of the middle ear and perforation of the drum, so that there was probably an ingrowth through the perforation rather than a metaplasia.

Zeroni¹³ saw cholesteatoma in a polypus of the ear, and Kessel and Stendener¹⁴ have described large epithelial pegs in polyp of the ear, without reporting anything possible about their origin. The publications of Troeltsch,¹⁵ Bezold,¹⁶ Siebenmann¹⁷ and Schwartz¹⁸ have not been at my disposal.

There are a number of communications to be found in literature relating to metaplastic conditions in the larynx, in the trachea and in the bronchial tubes and lungs.

Larynx.—Virchow,¹⁹ describing epidermoidal changes of the parts of the larynx which are lined with squamous epithelium, called this condition pachydermia laryngis, and Posner and Fraenkel²⁰ also published similar observations.

Heymann²¹ thinks that epithelial metaplasia of the larynx occurs in places which, during the functions of the larynx, are most exposed to mechanical irritation, and Haycroft and Carlier²² think that the transitional and stratified squamous epithelium present in the adult trachea are produced from a fully differentiated ciliated membrane through friction.

Kanthak²³ says: "We see, therefore, first, that after chronic irritation a general epidermoidal metamorphosis may be found over the whole larynx. Second, that only some parts of the mucous membrane of the larynx may be affected. Third, that transitional forms from cylindrical squamous epithelium, as described by Heymann,²¹ are only the initial stages of metaplasia. Fourth, that ulceration favors such metamorphosis, and, fifth, that the most pegs and more formation of papillæ are found in the older and more developed cases, and that in these cases the pegs may either be regular or irregular and atypical. In the latter case it has come to the development of a pachydermia diffusa." Referring to the length of the epithelial pegs, Kanthak²³ says: "The epidermoidal metamorphosis can go so far that a genuine pachydermia may be observed, with the development of strong papillæ and epithelial pegs, which often grow downward just as we are used to seeing it in the skin."

Bergengrün²⁴ describes a case of verruca dura laryngis, in which, following a chronic catarrh, he saw several white nodular tumors on the inside of the posterior wall of the larynx, the surface of which showed epidermoidal character, with hornification.

Trachea and Bronchi.—In a primary carcinoma of the trachea, which had its seat at the bifurcation, the microscopic examination showed that this primary growth was squamous epithelioma, while the neighboring mucous membrane and its mucous gland were lined with an unchanged cylindrical epithelium. There was a metastasis in the left suprarenal capsule, which also consisted of squamous epithelium. Virchow,²⁵ in speaking of the carcinomata of the bronchi, says: "They are pure bronchial carcinomata, and they mostly have the peculiarity to belong to the canceroids, as they have

REPORT ON EXPERIMENTS PERFORMED BETWEEN DECEMBER 6, 1903, AND APRIL 2, 1904.

No.	Date of Operation	Animal	Color	Sex	Weight, lbs.	Hemoglobin Per cent.	Date of Death	Postmortem	Last Wgt. Recorded in lbs.	Last Hemoglobin Recorded, Per cent.	Macroscopic Findings	Microscopic Findings.
1	Dec. 6, 1903.	rabbit.	not given.	...	3 $\frac{1}{2}$	82	Dec. 7.	1	Dec. 6, 3 $\frac{1}{2}$	Dec. 6, 82	Dilated stomach filled with food	
2	Dec. 1903.	rabbit.	not given.	...	2 $\frac{3}{4}$	78	Dec. 8.	1	Dec. 6, 2 $\frac{3}{4}$	Dec. 6, 78	Dilated stomach filled with food	
3	Dec. 20, 1903.	rabbit.	black.	f.	3 $\frac{3}{4}$	82	Dec. 21.	1	Dec. 20, 3 $\frac{3}{4}$	Dec. 20, 82	Dilated stomach filled with food	
4	Dec. 20, 1903.	rabbit.	black.	f.	4 $\frac{1}{2}$	89	Dec. 21.	1	Dec. 20, 4 $\frac{1}{2}$	Dec. 20, 89	Dilated stomach filled with food	
5	Dec. 20, 1903.	rabbit.	brown.	m.	5	86	Dec. 21.	1	Dec. 21, 5	Dec. 21, 86	Dilated stomach filled with food	
6	Dec. 20, 1903.	rabbit.	white.	f.	4 $\frac{3}{4}$	82	Dec. 20.	1	Dec. 20, 4 $\frac{3}{4}$	Dec. 20, 82	Dilated stomach filled with food	
7	Dec. 25, 1903.	B. hare	brown.	f.	5	85	Dec. 25.	1	Dec. 25, 5	Dec. 25, 85	Dilated stomach filled with food	
8	Dec. 25, 1903.	B. hare	brown.	f.	4 $\frac{1}{2}$	86	Dec. 25.	1	Dec. 25, 4 $\frac{1}{2}$	Dec. 25, 86	Dilated stomach filled with food	
9	Dec. 25, 1903.	rabbit.	brown.	f.	4 $\frac{3}{4}$	97	Dec. 26.	1	Dec. 25, 4 $\frac{3}{4}$	Dec. 25, 97		
10	Dec. 25, 1903.	rabbit.	brown.	m.	4 $\frac{3}{4}$	83	Dec. 26.	1	Dec. 25, 4 $\frac{3}{4}$	Dec. 25, 83		
11	Dec. 27, 1903.	rabbit.	brown.	f.	4 $\frac{3}{4}$	86	Dec. 28.	1	Dec. 27, 4 $\frac{3}{4}$	Dec. 27, 86		
12	Dec. 27, 1903.	rabbit.	brown.	f.	6 $\frac{1}{2}$	79	Dec. 28.	1	Dec. 27, 6 $\frac{1}{2}$	Dec. 27, 79		
13	Dec. 27, 1903.	rabbit.	brown.	m.	6	92	Dec. 28.	1	Dec. 27, 6	Dec. 27, 92		
14	Dec. 27, 1903.	rabbit.	brown.	m.	4 $\frac{3}{4}$	82	Dec. 29.	1	Dec. 27, 4 $\frac{3}{4}$	Dec. 27, 82		
15	Dec. 27, 1903.	rabbit.	brown.	f.	4 $\frac{3}{4}$	82	Jan. 1.	1	Dec. 30, 4 $\frac{3}{4}$	Dec. 30, 82		
16	Dec. 27, 1903.	rabbit.	brown and white.	m.	4 $\frac{3}{4}$	86	Jan. 1.	1	Dec. 27, 4 $\frac{3}{4}$	Dec. 27, 86		
17	Dec. 27, 1903.	rabbit.	black and white.	f.	5 $\frac{1}{2}$	83	Feb. 10.	1	Feb. 6, 5 $\frac{1}{2}$	Feb. 6, 83	Ulcer	Purul. infil't edges of ulcer.
18	Dec. 27, 1903.	rabbit.	black and white.	m.	4 $\frac{3}{4}$	79	Mar. 29.	1	Mar. 28, 4 $\frac{3}{4}$	Mar. 28, 65		
19	Dec. 27, 1903.	rabbit.	brown and white.	f.	4 $\frac{3}{4}$	83	Feb. 2.	1	Jan. 30, 3 $\frac{3}{4}$	Jan. 30, 55		
20	Dec. 31, 1903.	rabbit.	white.	f.	3 $\frac{3}{4}$	70	Dec. 31.	1	Dec. 31, 2 $\frac{3}{4}$	Dec. 31, 70		
21	Jan. 1, 1904.	rabbit.	white.	f.	4 $\frac{1}{4}$	85	Mar. 16.	1	Mar. 12, 2 $\frac{1}{2}$	Mar. 12, 52	Healed defect.	Metaplasia
22	Jan. 1, 1904.	rabbit.	brown.	m.	5 $\frac{1}{4}$	70	Jan. 6.	1	Jan. 1, 2 $\frac{1}{2}$	Jan. 1, 52	Ulcer	
23	Jan. 1, 1904.	rabbit.	brown.	m.	4	84	Feb. 8.	1	Feb. 6, 2 $\frac{3}{4}$	Feb. 6, 7		
24	Jan. 1, 1904.	rabbit.	brown.	f.	3 $\frac{3}{4}$	84	Jan. 2.	1	Jan. 1, 3 $\frac{3}{4}$	Jan. 1, 84		
25	Jan. 1, 1904.	rabbit.	brown.	f.	2 $\frac{3}{4}$	78	Jan. 3.	1	Jan. 3, 2 $\frac{3}{4}$	Jan. 3, 78		
26	Jan. 1, 1904.	rabbit.	white.	m.	4 $\frac{1}{2}$	83	Jan. 8.	1	Jan. 7, 4 $\frac{1}{2}$	Jan. 7, 83		
27	Jan. 1, 1904.	rabbit.	brown.	f.	5	90	Feb. 19.	1	Feb. 13, 3 $\frac{3}{4}$	Feb. 13, 38	Healed defect.	Metaplasia
28	Jan. 1, 1904.	rabbit.	brown.	f.	5	90	Apr. 20.	1	Apr. 10, 3	Apr. 10, 55	Ulcer	
29	Jan. 1, 1904.	rabbit.	brown.	f.	5	76	Jan. 8.	1	Jan. 8, 5	Jan. 8, 8	Ulcer	
30	Jan. 1, 1904.	rabbit.	brown and white.	m.	3 $\frac{1}{4}$	83	Jan. 14.	1	Jan. 8, 3 $\frac{1}{4}$	Jan. 8, 83	Ulcer with perforation peritonitis	
31	Jan. 8, 1904.	rabbit.	white.	f.	4 $\frac{1}{4}$	80	Feb. 15.	1	Feb. 6, 4 $\frac{1}{4}$	Feb. 6, 60		
32	Jan. 8, 1904.	rabbit.	white.	f.	3 $\frac{3}{4}$	85	Feb. 1.	1	Jan. 30, 2 $\frac{3}{4}$	Jan. 30, 50	(*)	Metaplasia
33	Jan. 8, 1904.	rabbit.	white.	f.	3 $\frac{3}{4}$	85	Jan. 21.	1	Jan. 8, 3 $\frac{3}{4}$	Jan. 8, 85	Ulcer	
34	Jan. 8, 1904.	rabbit.	brown and white.	f.	4 $\frac{1}{2}$	85	Jan. 13.	1	Jan. 8, 4 $\frac{1}{2}$	Jan. 8, 85	Two ulcers.	
35	Jan. 8, 1904.	rabbit.	gray and white.	m.	4 $\frac{1}{2}$	90	Jan. 9.	1	Jan. 8, 4 $\frac{1}{2}$	Jan. 8, 90		
36	Jan. 10, 1904.	B. hare	brown.	m.	5 $\frac{1}{4}$	81	Jan. 11.	1	Jan. 10, 5 $\frac{1}{4}$	Jan. 10, 81		
37	Jan. 10, 1904.	B. hare	brown.	f.	5 $\frac{1}{4}$	80	Jan. 10.	1	Jan. 10, 5 $\frac{1}{4}$	Jan. 10, 80		
38	Jan. 10, 1904.	B. hare	brown.	f.	3 $\frac{3}{4}$	76	Jan. 10.	1	Jan. 10, 3 $\frac{3}{4}$	Jan. 10, 76		
39	Jan. 10, 1904.	B. hare	brown.	m.	4 $\frac{1}{2}$	76	Jan. 11.	1	Jan. 10, 4 $\frac{1}{2}$	Jan. 10, 76		
40	Jan. 10, 1904.	B. hare	brown.	f.	4 $\frac{1}{2}$	72	Jan. 11.	1	Jan. 10, 4 $\frac{1}{2}$	Jan. 10, 72		
41	Jan. 10, 1904.	rabbit.	gray and white.	f.	3 $\frac{3}{4}$	69	Jan. 17.	1	Jan. 10, 3 $\frac{3}{4}$	Jan. 10, 69	Ulcer with hemorrhage.	Purulent infiltration of
42	Jan. 10, 1904.	rabbit.	white.	f.	3 $\frac{3}{4}$	77	Jan. 13.	1	Jan. 10, 3 $\frac{3}{4}$	Jan. 10, 77	Ulcer	edges.
43	Jan. 10, 1904.	B. hare	brown.	f.	4	75	Jan. 13.	1	Jan. 10, 4	Jan. 10, 75	Ulcer	
44	Jan. 10, 1904.	rabbit.	gray.	f.	3	68	Jan. 11.	1	Jan. 10, 3	Jan. 10, 68		
45	Jan. 10, 1904.	rabbit.	black.	f.	3	74	Jan. 25.	1	Jan. 23, 2 $\frac{1}{4}$	Jan. 23, 72	Ulcer	
46	Jan. 10, 1904.	rabbit.	gray and white.	f.	3	75	Jan. 10.	1	Jan. 10, 3	Jan. 10, 75		
47	Jan. 10, 1904.	rabbit.	white.	f.	3	87	Feb. 9.	1	Jan. 6, 2 $\frac{3}{4}$	Jan. 6, 60	Ulcer	Purulent infiltration of
48	Jan. 10, 1904.	rabbit.	white.	f.	3 $\frac{1}{4}$	68	Jan. 10.	1	Jan. 10, 3 $\frac{1}{4}$	Jan. 10, 68		edges.
49	Jan. 10, 1904.	rabbit.	white.	f.	3 $\frac{1}{4}$	86	Jan. 13.	1	Jan. 12, 3 $\frac{1}{4}$	Jan. 12, 86		
50	Jan. 12, 1904.	B. hare	white.	f.	3 $\frac{1}{2}$	82	Jan. 13.	1	Jan. 12, 3 $\frac{1}{2}$	Jan. 12, 82		
51	Jan. 12, 1904.	B. hare	brown.	f.	3 $\frac{1}{2}$	79	Jan. 13.	1	Jan. 12, 3 $\frac{1}{2}$	Jan. 12, 79		
52	Jan. 12, 1904.	B. hare	brown.	f.	3 $\frac{1}{2}$	84	Jan. 13.	1	Jan. 12, 3 $\frac{1}{2}$	Jan. 12, 84		
53	Jan. 12, 1904.	B. hare	black and white.	f.	4	68	Jan. 14.	1	Jan. 12, 4	Jan. 12, 68		
54	Jan. 12, 1904.	B. hare	white.	f.	4	85	Jan. 14.	1	Jan. 12, 4	Jan. 12, 85		
55	Jan. 17, 1904.	B. hare	gray.	f.	6	86	Jan. 18.	1	Jan. 17, 6	Jan. 17, 86		
56	Jan. 17, 1904.	B. hare	brown.	f.	5 $\frac{1}{2}$	82	Jan. 18.	1	Jan. 17, 5 $\frac{1}{2}$	Jan. 17, 82		
57	Jan. 17, 1904.	B. hare	brown.	f.	4 $\frac{1}{2}$	78	Jan. 18.	1	Jan. 17, 4 $\frac{1}{2}$	Jan. 17, 78		
58	Jan. 17, 1904.	B. hare	brown.	f.	6 $\frac{1}{4}$	85	Jan. 18.	1	Jan. 17, 6 $\frac{1}{4}$	Jan. 17, 85		
59	Jan. 17, 1904.	B. hare	brown.	f.	5 $\frac{3}{4}$	81	Feb. 10.	1	Feb. 6, 3 $\frac{3}{4}$	Feb. 6, 50	Ulcer	
60	Jan. 17, 1904.	B. hare	brown.	f.	5	82	Jan. 18.	1	Jan. 17, 5	Jan. 17, 82		
61	Jan. 17, 1904.	B. hare	brown.	f.	6	62	Jan. 18.	1	Jan. 17, 6	Jan. 17, 62		
62	Jan. 17, 1904.	B. hare	brown.	m.	4 $\frac{1}{2}$	76	Jan. 18.	1	Jan. 17, 4 $\frac{1}{2}$	Jan. 17, 76		
63	Jan. 17, 1904.	B. hare	brown.	m.	5 $\frac{1}{4}$	84	Jan. 18.	1	Jan. 17, 5 $\frac{1}{4}$	Jan. 17, 84		
64	Jan. 17, 1904.	B. hare	brown.	f.	4 $\frac{3}{4}$	88	Jan. 18.	1	Jan. 17, 4 $\frac{3}{4}$	Jan. 17, 88	Ulcer	
65	Jan. 17, 1904.	B. hare	brown.	m.	3 $\frac{3}{4}$	76	Jan. 18.	1	Jan. 17, 3 $\frac{3}{4}$	Jan. 17, 76		
66	Jan. 17, 1904.	B. hare	brown.	f.	3 $\frac{3}{4}$	80	Jan. 18.	1	Jan. 17, 3 $\frac{3}{4}$	Jan. 17, 80		
67	Feb. 21, 1904.	rabbit.	brown.	f.	5	83	May 12.	1	Apr. 10, 3 $\frac{3}{4}$	Apr. 10, 55		Metaplasia
68	Feb. 21, 1904.	rabbit.	gray.	f.	4 $\frac{1}{2}$	90	Feb. 29.	1	Feb. 23, 3 $\frac{1}{2}$	Feb. 23, 25	Ulcer with perforation.	
69	Feb. 21, 1904.	rabbit.	brown.	m.	4 $\frac{1}{2}$	75	Feb. 21.	1	Feb. 21, 4 $\frac{1}{2}$	Feb. 21, 75		
70	Feb. 21, 1904.	rabbit.	brown.	m.	4 $\frac{1}{2}$	88	Mar. 21.	1	Mar. 19, 2 $\frac{3}{4}$	Mar. 19, 41		
71	Feb. 21, 1904.	rabbit.	brown.	m.	4 $\frac{1}{2}$	85	Mar. 30.	1	Mar. 28, 4 $\frac{1}{2}$	Mar. 28, 73		
72	Feb. 21, 1904.	rabbit.	white.	f.	4 $\frac{1}{2}$	80	Feb. 22.	1	Feb. 21, 4 $\frac{1}{2}$	Feb. 21, 80		
73	Feb. 21, 1904.	rabbit.	black.	f.	3 $\frac{3}{4}$	70	Mar. 16.	1	Mar. 12, 2 $\frac{3}{4}$	Mar. 12, 45		
74	Feb. 21, 1904.	rabbit.	gray and white.	f.	70	70	Feb. 21.	1	Feb. 22, 70	Feb. 22, 70		
75	Feb. 21, 1904.	rabbit.	brown and white.	f.	4 $\frac{1}{2}$	65	Feb. 22.	1	Feb. 22, 4 $\frac{1}{2}$	Feb. 22, 65		
76	Feb. 21, 1904.	rabbit.	black and white.	f.	3 $\frac{3}{4}$	65	Feb. 22.	1	Feb. 22, 3 $\frac{3}{4}$	Feb. 22, 65		
77	Feb. 21, 1904.	rabbit.	white and yellow.	m.	4 $\frac{1}{2}$	65	Feb. 21.	1	Feb. 22, 4 $\frac{1}{2}$	Feb. 22, 65		
78	Feb. 21, 1904.	rabbit.	white and yellow.	f.	4 $\frac{1}{2}$	70	Feb. 24.	1	Feb. 22, 4 $\frac{1}{2}$	Feb. 22, 70		
79	Feb. 27, 1904.	rabbit.	brown and gray.	m.	4	85	Mar. 13.	1	Mar. 6, 4	Mar. 6, 81		
80	Feb. 27, 1904.	rabbit.	gray and brown.	f.	4 $\frac{1}{2}$	90	May 9.	1	Apr. 10, 3 $\frac{3}{4}$	Apr. 10, 73	One	Metaplasia
81	Feb. 27, 1904.	rabbit.	brown.	f.	72	72	Feb. 27.	1	Feb. 27, 72	Feb. 27, 72		
82	Feb. 27, 1904.	rabbit.	brown.	m.	3 $\frac{3}{4}$	90	Mar. 5.	1	Feb. 27, 3 $\frac{3}{4}$	Feb. 27, 90	Ulcer with perforation.	
83	Feb. 27, 1904.	rabbit.	brown.	f.	4 $\frac{1}{4}$	93	Mar. 23.	1	Mar. 19, 3 $\frac{3}{4}$	Mar. 19, 50	Ulcer	
84	Feb. 27, 1904.	rabbit.	white.	f.	3 $\frac{1}{2}$	92	Apr. 9.	1	Apr. 2, 2 $\frac{3}{4}$	Apr. 2, 46	Ulcer	
85	Mar. 6, 1904.	rabbit.	brown and white.	f.	5 $\frac{1}{2}$	70	Mar. 8.	1	Mar. 6, 5 $\frac{1}{2}$	Mar. 6, 70		
86	Mar. 6, 1904.	rabbit.	brown.	m.	3 $\frac{1}{2}$	95	Mar. 9.	1	Mar. 6, 3 $\frac{1}{2}$	Mar. 6, 95		
87	Mar. 6, 1904.	rabbit.	gray-brown.	m.	3 $\frac{1}{4}$	78	Mar. 8.	1	Mar. 6, 3 $\frac{1}{4}$	Mar. 6, 78		
88	Mar. 6, 1904.	rabbit.	gray and white.	m.	3 $\frac{1}{4}$	85	Mar. 12.	1	Mar. 6, 3 $\frac{1}{4}$	Mar. 6, 85		
89	Mar. 6, 1904.	rabbit.	brown.	f.	4 $\frac{1}{2}$	70	Mar. 20.	1	Mar. 12, 3 $\frac{1}{2}$	Mar. 12, 40	Ulcer	
90	Mar. 6, 1904.	rabbit.	brown.	m.	3 $\frac{3}{4}$	90	Mar. 6.	1	Mar. 6, 3 $\frac{3}{4}$	Mar. 6, 90		
91	Apr. 2, 1904.	rabbit.	black and white.	f.	4 $\frac{1}{4}$	75	Apr. 4.	1	Apr. 2, 4 $\frac{1}{4}$	Apr. 2, 75		
92	Apr. 2, 1904.	rabbit.	?	?	5 $\frac{1}{4}$	80	Apr. 4.	1	Apr. 2, 5 $\frac{1}{4}$	Apr. 2, 80		
93	Apr. 2, 1904.	rabbit.	?	?	3 $\frac{3}{4}$	65	May 18.	1	Apr. 10, 3 $\frac{3}{4}$	Apr. 10, 65		

* Healed defect; but in its place a thickened and hardened mucosa. † Death from ether.

Number of cases, 94; deaths, 94; death during the first week after operation, 69; postmortems, 35; ulcers, 20; healed defects, 3; hemorrhages, 1; perforations, 3; peritonitis, 1; microscopic examinations, 8; metaplasias, 5; purulent infiltration of the edges of ulcers, 3; purulent infiltration of the stomach was made, we have 5 metaplasias out of 25 cases.

a strongly developed squamous epithelium, which does not occur in the bronchi; they show about the same structure as the carcinomata of the vocal ligament or those of the lips." Virchow observed a pure bronchial carcinoma which had developed from a bronchus of the first order, and which had grown into the lumen, filling and obstructing it completely. It was interesting to see that a kind of dissemination had occurred in the lower bronchi, where a number of small eruptions had occurred.

In the case described by Friedländer,²⁶ a canceroid, according to his opinion, had developed from tubercular ulcers of the bronchus, the metaplastic and hornified squamous epithelium of which had grown into a lung cavity. Siegart²⁷ found a papilloma in the lower air passages which was covered with squamous epithelia, with prickles; it contained epithelial pegs and epithelial pearls, and was surrounded by a ciliated cylindrical epithelium. Pässler²⁸ also describes a primary bronchial carcinoma, with hornification, which, according to him, had developed from a germinal deposit. A squamous epithelioma of the bronchus of the upper lobe of the left lung at its junction with the main bronchus is reported by Ernst.²⁹ There were multiple metastases of similar structure in the dura mater of the cerebrum and cerebellum, and there was also, in the case of Bergengrün,³⁰ a metastasis in the left suprarenal capsule. Cases of squamous epithelioma of the lung have been reported by: Siegart,³⁰ 1 case; Borst,³¹ 2 cases, one of which was in a cavity; Ferls,³² 1 case; Grünwald,³³ 1 case, according to him originating from the alveolar epithelium and Wolf,³⁴ 1 case, developing from the lung and the bronchi, and another one growing in a cavity.

The writings of Ohloff,³⁵ Teschendorff,³⁶ Schmidt,³⁷ Griffin,³⁸ Fuchs,³⁹ Japha,⁴⁰ Wechselmann,⁴¹ Tillmann,⁴² Stumpf⁴³ and Werner,⁴⁴ in which more such cases have been reported, have not been at my disposal.

Pharynx.—In reference to hornification of epithelium in Waldeyer's adenoid ring of the pharynx, Siebenmann⁴⁵ says: "The tendency to proliferation and hornification of the epithelium is found in a pronounced way not only in the adenoid ring itself, but also in the mucous membrane in the vicinity. It develops either in a benign typical way or atypical, as carcinoma, which is often observed in this locality."

Tonsils.—On the tonsils, Wex⁴⁶ found hornified epithelium covering the gland of a child 5 years old.

Intestinal Tract.—In the stomach, physiologic hornification is found in some low mammalia, edentata, who have no teeth or only rudimentary teeth, and have to break up hard calcareous material in the stomach.

Leydig found physiologic hornification of the mucosa of the stomach in echidna, halmaturus and bradypus, the last belonging to the placentalia, and Posner⁴⁷ found hornification in the pars cardiaca and pars pylorica of man.

Borst⁴⁸ has observed squamous epithelioma in the stomach, the microscopic structure of which is shown in his work on tumors, but a description I have not been able to find.

Bohm⁴⁹ found squamous epithelium and squamous epithelioma in the rectum, but he himself is of the opinion that an ingrowth had taken place. Those are all the reports concerning metaplasia in the gastrointestinal tract which I have been able to find in literature, and we can, therefore, say that so far no pathologic epithelial metaplasias have been seen here.

Gall Bladder.—In the gall bladder, Lubarsch⁵¹ found an epithelial metaplasia in the form of an excrescence of the size of a lentil. The proliferation, which showed the picture of condyoma acuminatum, and was surrounded by hornifying and hornified squamous epithelium. There were present a chronic cholecystitis, four gallstones and a hydrops of the gall bladder.

Mönkeberg⁵² found a canceroid combined with adenocarcinoma of the gall bladder, and Deetz⁵³ reports four cases of squamous epithelioma of the gall bladder, which all had made metastases, consisting of squamous epithelium.

Pollak⁵⁴ saw two cases of squamous epithelioma of the gall bladder, and Nehrkorn⁵⁵ one, with hornified metastasis in the lymphatic glands.

Hausemann⁵⁶ illustrates a case without giving a description, and the reports of Rhein,⁵⁷ Weber⁵⁸ and Ohloff⁵⁹ were not accessible to me.

Dr. Arthur Dean Bevan has observed a case of adenocarcinoma with metaplasia. The case has not yet been published, but Dr. Bevan had the kindness to acquaint me with its details, and to permit me to mention it.

A farmer, aged 52, married, was admitted to the Presbyterian Hospital of Chicago, on Dec. 15, 1903, and dismissed on Jan. 3, 1904. There were symptoms of gallstones with gastric disturbances, dizziness and vomiting; attacks of violent pain in the gall bladder. The patient had lost twenty pounds during the last month preceding the operation, which was performed by Dr. Bevan on Dec. 23, 1903. The gall bladder was surrounded by adhesions and contained pus and gallstones. Scrapings from the gall bladder showed microscopically partly the picture of an adenocarcinoma and partly that of a squamous epithelioma.

Urinary Tract.—Rokitansky⁶⁰ was the first to describe cholesteatomatous changes in the urinary tract, but, as the edition of his text-book in which he speaks about this condition has not been at my disposal, I cite him from Glaeser: "Epidermoidal formations are found as thick and lamellated leaves of white, glossy layers of epidermis cells, which are desquamated. They occur in small patches, or cover large portions or even the surface of the whole apparatus, down into the urethra, as a proliferation occurring on the chronic inflammatory papillary mucous membrane." According to Orth,⁶¹ satin-like, glossy desquamations, consisting of hornified epithelial cells, form on an infiltrated mucous membrane, and this change may occur with tuberculosis, with cystic pyelitis, or alone, as he observed it in the case where there also was a xerosis of the conjunctiva.

In the pelvis of both kidneys strong thickening of the epithelium and a satin-like, glossy mass was found, with numerous brownish-yellow concretions.

Epstein⁶² also saw such changes in the pelvis of the right kidney, with formation of papillae in the mucous membrane.

Marchand observed a case which was described in a dissertation by Liebenow.⁶³ A perineal section of a boy of 14 years had led to the formation of a fistula, and epidermoidal changes of the epithelium of the urinary passages developed, which Marchand at first believed to be due to an ingrowth through the fistula, while later he gave up this belief, thinking the change was due to metaplasia. There was also a cholesteatoma on the lower surface of the diaphragm which probably represented a metastasis.

Kischensky⁶⁴ describes a primary squamous epithelioma of the calices, with a tendency to hornification and petrification. From the pelvis down to the bladder hornification of the upper epithelial layers was found, and the calices were filled with a new formation which had also involved the right kidney. Metastases were found in the liver, in the right knee joint, and also in the epiphysis and diaphysis of the femur.

Beselin⁶⁵ reports cholesteatomatous desquamation in the pelvis, with primary tuberculosis of the kidney on the same side, and Halle, epidermoidal changes in the pelvis, the ureters and in the bladder, with chronic cystopyelonephritis, while Braatz saw epidermoidal changes in the pelvis of an extirpated kidney, with the stratified squamous epithelium, prickly cells, basal cylinder cells and a stratum corneum. The patient recovered after operation, and it seems that epidermoidal desquamation had caused the attacks of colic which had existed before the operation.

Pollak⁶⁶ describes a case which was observed by Lubarsch; in the pelvis, and up into the calices, were found glossy, gray-white patches consisting of hornified squamous epithelium, due, according to Pollak, to genuine metaplasia.

Metaplasia in the urinary tract has also been observed by Dr. William F. Metcalf and Dr. Homer E. Safford⁶⁷ of Detroit, who describe an epithelial metaplasia in the pelvis, with a stone in the lower part of the ureter of the same side, and at the seat of the stone an adenocarcinoma was found. This is surely of great importance, if we consider the rarity of carcinoma in the urethra. The case has not been published yet, but Dr. Safford had the kindness to send me his specimens and the description, in order to enable me to become acquainted with the details of the case.

In the bladder, cases have been described by Marchand⁶⁷ of squamous epithelioma with canceroid plaques, prickly cells and horn pearls; Halle,⁶⁸ chronic hypertrophic interstitial cystitis and epidermoidal changes confined to the mucosa of the bladder, and by Thompson,⁶⁹ Cabot⁷⁰ and Pollak.⁶⁶

Thompson saw an epithelioma of the bladder; Cabot, a case of pachydermia vesicae, with chronic cystitis, and Pollak, patches of epithelial metaplasia with chronic cystitis.

In the urethra, Thiersch⁷¹ found squamous epithelioma in the pars membranacea, and the anterior part of the pars prostatica, and Bumm⁷² writes about metaplastic changes in the urethra.

Neelsen⁷³ and Finger⁷⁴ found metamorphosis of the cylindrical epithelium to squamous epithelium in chronic gonorrhea, and Pollak,⁶⁶ patches of metaplasia in an inflamed and ulcerated urethra, and also a papillary growth having the structure of a condyloacuminatum. This growth had the size of a pea, and a hornified surface.

The reports of Thiersch, Bumm, Klebs,⁷⁵ Abbaran,⁷⁶ Kaufmann⁷⁷ and Lowensohn⁷⁸ were not accessible.

Genitalia.—In the testicle, Virchow⁷⁹ found cholesteatomatous masses in a fibrocystoid, and Lotzbeck,⁸⁰ cited by Beselin, a cholesteatoma of the size of a cherry-stone.

In the uterus, Velt⁸¹ and Ruge⁸² are convinced that a metaplasia of the cylindrical epithelium of the mucosa of the cervix and that of the erosions to squamous epithelium occurs, and Ruge brings illustrations demonstrating such metamorphosis of the epithelium.

Zeller⁸³ claims to have found squamous epithelium in the uterus in 63 cases, while Ruge and Gebhard⁸⁴ consider metaplasias in the uterus as a rare occurrence if we exclude cases where they occur on the surface of polypi of the cervix, which protrude through the external orifice, and also metaplasias in old inversion of the uterus. In a number of polypi the deepest seated of which protruded through the external orifice, Küstner found squamous epithelium. The combination of glandular carcinoma and horny carcinoma has been described by Emanuel,⁸⁵ who brings good illustrations to prove that the epithelia have been changed to squamous epithelium. In the case of Hering,⁸⁶ of a canceroid of the uterus, the size of an egg, the rest of the mucosa of the organ was covered with squamous epithelium, and Gebhard published a case of hornified squamous epithelioma of the uterus in a woman 66 years of age, who had been treated locally with crude pyroligneous acid, and who wore a pessary for prolapsus of the vagina. Gellhorn⁸⁷ and Lahmann⁸⁸ each report a case of hornifying carcinoma, and E. Fraenkel and G. Wiener⁸⁹ saw cells in glandular carcinoma of the corpus uteri which very much resembled squamous epithelium. They formed nodules consisting of solid strands.

Fleischlen⁹⁰ observed a case, and says: "All microscopic pictures prove the origin of the squamous epithelioma of the corpus uteri from the superficial epithelium. The cylindrical epithelium here has been changed into stratified squamous epithelium, which lines the whole inner surface of the uterus."

Pollak⁵⁴ Case 1, psammocarcinoma, with hornifying epithelium and prickly cells; Case 2, typical prickly cells and hornification.

Hofmeier⁹¹ reports four cases: an adenocarcinoma with typical squamous epithelioma, a squamous epithelioma of the uterus, a squamous epithelioma and glandular carcinoma of the uterus, and a glandular carcinoma and squamous epithelioma in a woman of 50 who was a virgin.

In a case of Zahlotsky⁹² a microscopic examination of masses scraped from the uterus led to the diagnosis of a squamous epithelioma, and Emanuel⁸⁵ describes a hornified carcinoma of the uterus, the whole cavity of the uterus being lined with a stratified squamous epithelium.

Löhlein, cited by Fleischlen, describes a squamous epithelioma, and von Rosthorns⁹³ article on hornification of the mucous membrane of the uterus has not been at my disposal.

Mammary Gland.—In the mammary gland, Kürsteiner⁹⁴ observed an adenoma of the gland, with cylindrical and stratified and partially hornified epithelium.

Pollak⁶⁶ found squamous epithelial cells, prickly cells, concentric arrangement of the cells and hornification in a metastasis from a cylindrical cell carcinoma of the stomach in the lung. He says: "This case is of particular interest, as the suspicion of the development of epidermis cells from an aberrant germinal deposit can be excluded, the primary growth only consisting of cylindrical epithelium, as all the cells of the primary carcinoma originated from cells which embryologically have nothing in common with epidermis cells. If, then, metaplasia is found in those metastases, it proves that cylindrical epithelium can be transformed to squamous epithelium under the influence of certain external conditions."

It may be that Pollak's opinion is the correct one, but he has not brought sufficient proof, or at least he does not say that he has

examined the whole primary growth, and he, therefore, can not exclude a metaplasia having occurred there, as we have seen it in other cases reported.

Cullen⁷⁷ reports a number of cases of squamous cell carcinoma of the uterus and cervix.

We have already mentioned a number of cases of combination of cylinder cell carcinoma with squamous epithelioma reported in literature, and there is no reason to suppose that such a combination could not occur in the stomach as well as in other organs, particularly since Borst has reported his squamous epithelioma of the stomach. To the reports, which I have mentioned above briefly, I only wish to add the conclusions to which Eichholz⁹⁸ has come in his experimental work. They are: "1. Stratified squamous epithelium can become so similar to epidermis that the two can not be differentiated. 2. In epidermoidal changes of transitional epithelium a metaplasia can not always be excluded with certainty, although in most cases, as in my experiments, it is to be supposed that an ingrowth has taken place. 3. Cylindrical epithelium can never form epidermis. This is proven by my experiments and by my examinations of human material. If an epithelial metamorphosis occurs in an organ, with cylindrical epithelium, then both, the cases reported in literature and the results of my own experiments, ingrowth of squamous epithelium or a development from an aberrant germinal deposit, furnish the explanation."

If Eichholz had given the cases reported in literature a more unbiased consideration, he would not have come to such radical conclusions. He would find it hard, indeed, to mention only one instance, to explain the occurrence of squamous epithelioma in the gall bladder, in cases where no fistula exists, without admitting a genuine metaplasia to be at least highly probable.

As regards the cases reported in literature, many of them must no doubt be classed as genuine forms of metaplasia, while others may have to be eliminated, but the process of sifting, in order to be of real value, must be done with great care.

If we review the cases recorded, adding our own, we can probably differentiate between three different forms of metaplasia, namely:

1. A more superficial form in the nose and its accessory cavities, and in the ear, where the massing of the desquamated cells often leads to the formation of cholesteatoma.

2. Metaplasias with considerable thickening and hornification, such as occur in the urinary tract and in the uterus, often in the course of time, followed by the formation of a squamous epithelioma.

3. Metaplasia which leads to squamous epithelioma so early that it is hard or impossible to find metaplastic changes in the mucous membrane surrounding the metaplastic tumor.

4. My own cases of metaplasia show strong downward growth at once, and it looks as if they belonged to the former class, but I should like to have it distinctly understood that they do not represent carcinoma yet, and that only further experiments can show what eventually becomes of them.

As to the causes of epithelial metaplasia, we have seen that hydropic conditions of the mucosa of the antrum of Highmore may lead to metaplastic changes. Catarrhal conditions, rhinitis and influenza rhinitis are pointed out as causes by Suchanek and others. Gonorrhea may lead to metaplasia in the urethra; tuberculous ulcers in the air passages, stones in the gall bladder and concretions and stones in the urinary tract play

an important part, but the importance of mechanical factors is very clearly indicated by the fact that some animals who have to break up hard food in the stomach have a physiologic metaplasia with hornification reaching down to the pylorus.

In my own cases I think it very probable that friction has been the cause of the metaplasia.

In closing this paper, I wish to state that the frequent allusions to the close relation between metaplasia and carcinoma found in the literature here reported, and also in the report of my own work, are intended only to point out the actual relations as they exist, that the reader may receive a fair idea of them. The cases which have been reported make it very clear that epithelial metaplasia is very often followed by the development of a carcinoma, but what the real relations are between my metaplasias and carcinoma is a matter that can only be made clear by further experimental work.

34 Washington Street.

LITERATURE.

1. Fütterer: Aet. d. Carc., Bergmann, Wiesbaden, 1901, and THE JOURNAL A. M. A., March 15, 1902.
2. Lehrbuch d. Augenheilkunde.
3. Graefe's Arch. f. Ophthalm., vol. xxix.
4. Monatsschr. f. Ohrenheilk., 1891, vol. iv.
5. Volkmann's Samml. klin. Vort., No. 340, 1889.
6. Zeitschr. f. Heilk., vol. xxi, No. 6, 1900.
7. Langenbeck's Arch., vol. xxxix, 1889.
8. Wiener klin. Woch., 1890, p. 953.
9. Virch. Arch., vol. cxviii, 1892. Virch. Arch., vol. viii, p. 371.
10. Monatsschr. f. Ohrenheilk., 1868, No. 9, cited from Wendt. Arch. f. Heilk., Wagner, vol. xiv. Arch. f. Ohrenheilk., vol. vii, 1873.
11. Arch. f. Augen. u. Ohrenheilk., vol. iv, p. 162.
12. Arch. f. Ohrenheilk., vols. xxvii and xxix, p. 42.
13. Arch. f. Ohrenheilk. vol. xlii, 1897.
14. Arch. f. Ohrenheilk., vol. iv.
15. Lehrb. d. Ohrenheilk., 5. Aufl., p. 425.
16. Zeitschr. f. Ohrenheilk., vol. xx.
17. Zeitschr. f. Ohrenheilk. Steinbrügge: Zeitschr. f. Ohrenheilk., vol. viii, p. 224.
18. Schwartz: Chir. Krankh. d. Ohres., p. 221.
19. Berl. klin. Woch., No. 19.
20. Deutsche med. Woch., 1889, p. 30.
21. Virch. Arch., vol. cxviii.
22. Quarterly Journal of Micr. S., New Series, No. 120, vol. xxx, part 4, 1890.
23. Virch. Arch., 118, 119 and 120.
24. Virch. Arch., vol. cxviii, 1889. Reiche: Centralbl. f. Allgem. Path., 1893.
25. Berl. klin. Woch., 1891, No. 50, p. 1185.
26. Fortschr. d. Med., vol. iii, No. 10, 1885.
27. Virch. Arch., vol. cxix, 1892.
28. Virch. Arch., vol. clxv, 1896.
29. Ziegler's Beitr., vol. xx, p. 155.
30. Virch. Arch., vol. cxxxiv.
31. Die Lehre von d. Geschwülsten, vol. ii, p. 899.
32. Virch. Arch., vol. lvi, p. 437.
33. Münch. med. Woch., 1889, No. 32.
34. Fortsch. d. Med., 1895.
35. Ueber Epithelmetaplasie an der Schleimhaut von Gallenblase u. Trachea. Diss., Greifswald, 1891.
36. Ein Fall von multiplen Papillomen d. Larynx u. d. Trachea. Diss., Königsberg, 1893.
37. D. Krankh. d. ob. Luftwege, Berlin, 1896.
38. Gac. Clinique de Torino, 1875.
39. Beitr. z. Casulistik d. prim. Lungencarcinoms. Diss., Leipzig, 1890.
40. Ueber prim: Lungenkrebs. Diss., Berlin, 1892.
41. Ein Fall v. prim. Lungenkrebs.
42. Tillmann: 3 Fälle v. prim. Lungencarcinom. Diss., Halle, 1889.
43. Stumpf: Diss., Glessen, 1891.
44. Werner: Diss., Freiburg, 1891.
45. Watsuji: Zeitschr. f. Krebsforschung, B. 1, H. 5, 1904. Has come too late to be considered in this article. I would call attention to it on account of the cases reported by Watsuji, and also for additional literature.
46. Ueber Verhörung des Epithels im Gebiet des Waldeyerschen adenoiden Schlundrings. Arch. f. Laryng. u. Rhin., vol. ii.
47. Beitr. zur. norm u. Path. Hist. d. Rachenröhre, Zeitschr. f. Ohrenheilk., 1899.
48. Virch. Arch., vol. cviii.
49. Die Lehre von den Geschwülsten, vol. ii, 1902, p. 665, 1889, Fig. 180.
50. Virch. Arch., No. 140, p. 524, 1895.
51. Arbeiten aus d. Path. Anat. Abtheil. des Königl. Hyg. Instituts zu Posen, 1901.
52. Virch. Arch., 164, 1902.
53. Virch. Arch., 164, 1902.
54. Arbeiten aus d. Path. An. Abth. d. Kgl. Hyg. Inst. zu Posen, 1901.
55. Virch. Arch., vol. cliv.
56. Die Diagnose d. bösartigen Geschwülste.
57. Cited from Deetz.
58. Diss., Würzburg, 1891.
59. Diss., Greifswald, 1891.
60. Lehrb. d. Path. Anat., Wien., 1861.

61. Lebrb. d. Spec. Path. Anat., vol. ii, p. 208.
62. Deutsch. Arch. f. klin. Med., 1882, vol. xxxi, pp. 63-67.
63. Ausgedehnte Epidermis bekleidung d. Schleimh. d. Harnwege. Diss., Marburg, 1891.
64. Kischensky: Ziegler's Beitr., vol. xxx, No. 2.
65. Beselin: Virch. Arch., vol. xcix.
66. Not yet published.
67. Langenbeck's Arch., vol. xxii, p. 676, 1878.
68. Ann. des mal. des organes genitourinaires, 1896.
69. Lectures on some important points connected with the urinary organs. Philadelphia, 1884.
70. Amer. Jour. of the Med. Sc., 1891.
71. Der Epithelkrebs. Cited from Marchand.
72. Transactions of the Path. Soc. of London, vol. xl.
73. Arch. f. Derm. u. Syph., 1887, p. 177, 1889.
74. Arch. f. Derm. u. Syph., 1887.
75. Handb. d. Path. Anat., p. 68.
76. Les tumeurs de la vessie; Paris, Steinheil, 1892, p. 132.
77. Deutsch. Chir., Lief. 50 a.
78. St. Petersburg. Med. Zeitschr., 1862, vol. ii, p. 225.
79. Virch. Arch., vol. viii, p. 401.
80. Cited from Beselin.
81. Zeitschr. f. Geb. u. Gyn., vol. ii.
82. Zeitschr. f. Geb. u. Gyn., vol. viii, 1882.
83. Zeitschr. f. Geb. u. Gyn., vol. ii, 1884.
84. Zeitschr. f. Geb. u. Gyn., vol. xxiv, 1892.
85. Zeitschr. f. Geb. u. Gyn., vol. xlv, No. 3, 1901.
86. Zeitschr. f. Heilk., vol. viii, Wien, 1887.
87. Zeitschr. f. Geb. u. Gyn., vol. xxxvi, p. 430.
88. Arch. f. Gyn., vol. lxii, p. 439.
89. Fraenkel and Wiener: Beitr. z. Geb. u. Gyn., vol. ii, p. 351, 1899.
90. Flaischlen: Zeitschr. f. Geb. u. Gyn., vol. xxxii, 1895.
91. Zeitschr. f. Geb. u. Gyn., vol. xxxii.
92. Nouv. Arch. d'Obstr. et de Gyn., vol. xxxii, 1895.
93. Emanuel: Zeitschr. f. Geb. u. Gyn., vol. xxxii, 1895.
94. Festschr. d. Deutsch. Ges. f. Gyn., Wien.
95. Virch. Arch., vol. cxxxvi, 1894.
96. Metaplasie in Carcinoma metastasis. Arb. aus d. Path. Abth. d. Koenigl. Hyg. Inst. zu Posen, 1891.
97. Cancer of the Uterus, New York, 1900.
98. Experimental work: Eichholz: "Experimentelle Untersuchungen ueber Epithelmetaplasie." Arch. f. klin. Chir., vol. lxxv, No. 4, 1902, p. 1002.

LITHEMIC NASOPHARYNGITIS DUE TO SYSTEMIC DISTURBANCE.*

J. A. STUCKY, M.D.

LEXINGTON, KY.

Inasmuch as closer and more accurate observation has been made, new facts brought out, old theories exploded, and better results obtained, since the reading and discussion of this topic at the Western Section of the American Rhinological, Otological and Laryngological Society held in St. Louis two years ago, I do not feel that it is necessary to offer any apology for bringing the subject again before you. I desire to emphasize in the very beginning that it is full and free discussion that we are after.

In the treatment of all disease, whether due to pathologic condition or functional disturbance, the fact as to the cause of the existing condition is what concerns us most. What is the matter and why does it exist, and what will relieve or palliate most effectually the condition that confronts us, are the perplexing questions that we must face.

By the term lithemic nasopharyngitis is meant that type of disease having a close etiologic relationship to rheumatism, or the so-called lithemic diathesis, and has its origin in autointoxication caused by the absorption of by-products from the intestinal tract, or in the faulty metabolism occurring in the liver.

It may be defined as an acute congestive rather than an inflammatory process.

In the former paper, I believed the cause to be acid urates which had not been eliminated by the proper channels, or the absorption of products from the alimentary tract, the exact nature of which was unknown. In a record of one hundred cases where the urine has been carefully examined, and indican in excess found, as soon as that form of treatment was followed which

eliminated this, the patient was relieved, and as this was found more frequently than the excess of acid urates, I conclude it is the most potent excitant of the nasopharyngeal irritation. Hence more stress is put on its elimination than on uric acid. The local manifestations of this so-called diathesis may not be confined to the pharynx, but may exhibit itself in affections of the laryngeal, nasal, auditory, and gastrointestinal tract, without any special organic or systemic symptoms.

The attack causes primarily no lesion, and there is no distinct pathologic alteration, this being a local manifestation of a systemic condition. These cases were almost invariably neurotic, good livers, leading more of a mental than an active physical life. They complain of always taking cold, sneezing, have a stuffy, full feeling in the head, one nostril always stopped up, copious and frequent discharge from the nose, greatly annoyed by accumulation behind the palate and crackling noise in the ear. In addition to these nasal symptoms, attention will be called to a stiffness in the throat, with the sensation of a foreign body, like a piece of bread-crust or fish-bone in the tonsil. Altogether they are a miserable, snorting, snuffling, handkerchief using, irritable, and generally uncomfortable set who blame all their ills on a "catarrh."

Interrogation reveals the fact that more or less constipation exists and some laxative is frequently needed. Little water is drunk. Meat and sweets are freely used as well as salads, acids, coffee and tea. The skin is abnormally florid or muddy, in the former case suggesting too free use of alcoholics, in the latter so-called "biliousness"; in both cases the liver is most unmercifully suspected if not damned. A case with such a history leads me to suspect at once indicanuria with probably an excess of uric acid.

My observation is that where there is an abundance of indican in the urine in cases such as I have described there is no satisfactory and permanent relief until the treatment thoroughly eliminates this and the insufficient functioning of the large and small intestine be remedied and improper diet corrected. It is not so much a question of diet as of regulating the intestinal tract. In aggravated cases, I always have the gastric contents analyzed after giving a suitable test meal, in order to know how much, if at all, the stomach is at fault, for if gastric digestion is not complete, and the contents of the stomach are discharged into the duodenum in a hyperacid condition, the secretion of this organ is not sufficient to neutralize the acidity and, as a result, intestinal digestion is inhibited. The result is a putrefactive fermentation of the mass, developing disagreeable flatulence, dyspnea and intestinal distress. Many of the cases have more nasopharyngeal annoyance, in the form of sneezing, running at the nose, either immediately or several hours after eating. In the former case we suspect the trouble to be reflex gastric irritation; in the latter intestinal toxemia, due to indigestion.

It has been demonstrated that ingestion of citric, melic, or the fruit acids during peptic digestion inhibits secretion of hydrochloric acid and the peptic juices and increases fermentation of the carbohydrates, and the elimination of fruit in the diet, especially with meals, has been most beneficial. Where the patient craves an acid, and many of these patients do, dilute mineral acid drinks, preferably hydrochloric, are given with meals. When the cause of suboxidation is found and removed the victory is won, and in order to bring this about treatment must be directed toward elimination, atten-

* Read at the Fifty-fifth Annual Session of the American Medical Association, in the Section on Laryngology and Otology, and approved for publication by the Executive Committee: Drs. G. Hudson Makuen, George L. Richards and John F. Barnhill.