A CASE OF ACUTE, NON-TRAUMATIC, MULTIPLE OSTEO-MYELITIS,

PRODUCED BY THE STREPTOCOCCUS, OCCURRING IN A MALE ADULT.*

JOHN T. BOTTOMLEY, M.D.
BOSTON.

NARRATION OF CASE.

A. H. C., 43 years old, a theatrical manager, was born in France, lived most of his life in England, and had been five years in this country. His family history was negative. His previous history up to 1892 was not noteworthy. Though he had led an irregular life, with much excitement, travel and exposure, he had always been strong and robust. Venereal infection was denied; a moderate use of alcohol admitted. About 20 years ago he broke his left tibia at the junction of the lower and middle thirds, but it had given him no bother since.

Ten years ago he had an attack of "left-sided pleurisy" which lasted about six weeks. Though he soon recovered his normal weight and strength, he has since then had more or less pain in his left chest, a harassing cough, an almost constant bloody expectoration and the lower lobe of his left lung has been flat to percussion. These symptoms have always been exaggerated after prolonged physical exertion or mental worry. Occasionally, but only for a period of a few weeks at most, he would be free from them. Such a period occurred last spring, but the symptoms, returning in June, continued till his death. Several specialists, both here and abroad, had examined him, but a definite diagnosis had never been made. The uniformly bloody, prune-juice expectoration had been carefully examined again and again for tubercle bacilli, streptothrix, actinomyces and for neoplastic elements, but always with negative results. He had undergone, as a diagnostic measure, most thorough anti-syphilitic treatment without relief or change in his condition. His symptoms, subjective and objective, continued the same year after year. Repeated punctures at different times showed only a thickened pleura. No fluid, purulent or otherwise, was ever obtained, nor was a cavity in his lung ever demonstrated. A distinguished New York specialist who saw him a number of years ago had an opportunity of re-examining him last fall and found the physical signs unchanged, which rendered very improbable the diagnosis of a slow-growing malignant tumor of the lung which had once been advanced.

Aug. 1, 1902, while making a railroad journey, very sudden and severe pain began both in his left arm just below the shoulder and in his left leg about the middle. Though he had considerable fever (103), the severe pain was his only subjective symptom for several days; then he had a chill, a diarrhea started in and he began to feel "sick all over." A physician was called who reports that at that time (six days after the acute onset) there was "a swelling with deep pain on pressure over the inner surface of his left upper arm, at the junction of the middle and upper thirds. There was also deep induration over the outer aspect of the middle third of the left leg. Both arm and leg were swollen and the swellings were so tender that a thorough examination was impossible. There was no glandular involvements. The urine was negative; there was no temperature." Anti-rheumatics internally with local cooling lotions seemed to afford some relief for a day or two, but the pain then became more intense and the patient started a run of fever which varied between 100 and 102, but followed no definite type. Sweating now began, slight at first, but gradually increasing till it became a feature of his illness. A white count, made about the sixteenth day, showed a leucocytosis of 17,000. The patient became more and more ill; the temperature assumed a higher level and fluctuation finally appeared in the swellings on the arm and leg. A surgeon

who saw the case in consultation at this stage opened abscesses under local anesthesia, the patient's critical condition contraindicating a general anesthetic. The humerus and ulna were found to be partially denuded of periosteum, but connection with the bone marrow could be established." During the following month several operations were done on the arm and leg to secure proper drainage. The symptoms, however, continued and the patient's condition grew gradually worse. His temperature varied from 101 to 103, while his pulse was about 125. A month after the onset of the disease his right forearm and right leg began to swell and to become tender and painful. An abscess over the upper part of the ulna was opened and drained; bare bone was once more found. The leg was not opened. The pus from all the abscesses showed the streptococcus in pure culture and the same organism was found in the expectoration. About the sixth week of the illness the state of the left arm became so bad that amputation was advised, but it was not done, partly because of the patient's unwillingness and partly because of his condition, which by this time had become so critical that even simple incision of an abscess under local anesthesia threw him into marked shock. For the latter reason a sequestrum which had formed in the left humerus was not removed. Numerous punctures into the flat area of the chest were made, but no pus was found. Yet a specialist in chest diseases who saw the patient in the sixth week of his illness diagnosed the chest condition as a pulmonary abscess.

Oct. 17, 1902, in the eleventh week of his illness, the patient came under my personal observation. He was emaciated and evidently septic. His temperature was running about 101.4, his pulse about 120. His respirations were only slightly accelerated. His chief subjective symptoms were his cough, severe pain in the left arm and right leg and moderate pain in his left leg and right forearm. His urine showed a slight trace of albumin, but was not particularly abnormal. He had an intermittent diarrhea which annoyed and weakened him greatly.

The whole left upper arm was swollen, thickened and edematous. On all its aspects were the wounds of his previous operations, some partially healed, some wholly open. From all came a free, foul-smelling, thick, purulent discharge; through all the examining finger could feel bare, roughened bone. The shoulder joint was involved in the process. The left forearm and hand were also swollen, but only as a result of the impeded return circulation.

On the external aspect of the left leg in its upper third was a partially healed wound about four inches in length. A sinus at its center led down to the fibula denuded of its periosteum over a considerable area. The discharge was not profuse. The upper half of the leg was only moderately swollen.

There was some general swelling of the whole right leg, but it was more noticeable over the lower half, especially just above the ankle joint, where there was marked edema and the tenderness was exquisite. The skin over this area was not reddened.

The right forearm was swollen from elbow to wrist, mostly posteriorly and on the ulnar side. Over the upper posterior surface of the ulna was a wound two inches long which, like the others, led down to bare bone and gave exit to a profuse discharge of thick pus. The elbow joint was apparently not involved, but was fixed at an angle of 45 as a result of its being held in that position for some weeks.

The rest of the story can be briefly told. At the first visit simple incisions were made in the right leg to relieve the tension and let out the pus. The previous incisions on the other members were lengthened to provide more effective drainage. These procedures afforded so little relief that, after consultation with Dr. John C. Munro, radical measures were decided on, notwithstanding the critical condition of the patient. In two operations, one following ten days after the other, under general anesthesia, a very foul sequestrum six inches long was removed from the left humerus (the well-marked involucrum adding much to the difficulty of the operation), one

* Read at the Fifty-fourth Annual Session of the American Medical Association, in the Section on Surgery and Anatomy, and approved for publication by the Executive Committee: Drs. A. J. Ochsner, J. E. Moore and De Forest Willard.

about four inches long from the left fibula and another about two inches long from the right ulna; the right leg was freely opened and large areas of bare bone found on both tibia and fibula; the tibia was trephined and, pus having been found in the medullary cavity, a gutter was made with a chisel down the whole front of the cortex. The medullary cavity of the right fibula was opened with rongeur forceps and drained. It was not so extensively involved as the tibia. The right ulna was treated likewise and free drainage provided.

Necrosis of bone was very extensive in the left humerus and right tibia. There was a large quantity of foul, thick pus in the soft parts of the lower right leg. The periosteum was but little thickened over the upper half of the tibia, but was retracted from a large portion of the cortex of the lower half.

These operations brought the temperature down to normal for the first time in the course of the disease and there it remained till the end, save when backing up of the pus on two occasions caused a temporary rise to 101. The pulse also fell to the neighborhood of 100 and for two weeks following the operations the patient was very comfortable and spent part of each day in a wheelchair. But, unfortunately, recurrent attacks of diarrhea, the several operations necessary to maintain proper drainage, with possible advancing amyloid disease of his internal organs, so weakened him that he died early in January, 1903, on the one hundred and fifty-fifth day of his illness. Without doubt he had septicemia. An autopsy could not be obtained.

His fever was at its highest during the third and fourth weeks, touching 104.8 on the twenty-ninth day. It touched normal for the first time on the ninety-seventh day. The average temperature was about 101.5; the average pulse about 115; the highest rate reached was 140, coincident with the highest point of temperature. Except for a period of two weeks following the free opening of the bones, the patient was very restless and nervous, particularly at night. On and off a severe diarrhea gave him much trouble and was most difficult to control. He took very large quantities of nourishment throughout his whole illness and had very little gastric disturbance. Profuse sweats were common; chills were absent. During the course of his illness he developed a marked pilosis—a condition not uncommon in these cases. His cough was very troublesome and the expectoration always blood-stained, sometimes with dark, but more often with bright blood.

The immediate cause of his death was cardiac failure, of which he had several attacks during the last two weeks of his illness.

COMMENTS ON THE CASE.

With regard to the diagnosis in the case there is but little to say. The long-standing affection of his lung, whatever may have been its nature, afforded to the streptococcus a focus of development in the body, a point of entrance to the circulation and a consequent opportunity to produce the unusual condition already described.

After the paper of Lexer¹ and that of Nichols, presented at the present meeting, there is at present nothing to be added to the bacteriology and pathology of osteomyelitis. This clinical contribution to the subject has reason for its being in these three factors: (a) the case showed multiple foci; (b) it occurred in an adult 43 years old, and (c) the infecting organism was the streptococcus.

Acute multiple osteo-myelitis, while not a very rare disease, is still by no means common. Funke² reported to the Surgical Congress in 1895 644 cases of osteomyelitis; of these only 37 were examples of the multiple type. Neumark³ reports 30 cases, five of which were multiple; Volkmann and Roser⁴ consider the multiple form rare, and Wenzel reports a single case in a girl of 13. Haaga⁵ says that one-fifth of all the cases are multiple. But, apart from the question of the rarity of

this form of the disease itself, the fact that the present case occurred in an adult far removes it from the ordinary run of osteo-myelitis cases. Only 56 of Funke's 644 cases were found in patients over 25 years old. Of the 37 cases with multiple foci, all—without exception—occurred in children. In fact, he says that the multiple form occurs only in children. Four cases mentioned by Volkmann⁶ were all in patients under 14 years old.

In a fairly thorough search of literature I can find but one similar case of acute multiple osteo-myelitis occurring in an adult. H. F. Devel, in the proceedings of the Warsaw Society in December, 1885, reports the case of a soldier, 24 years old, who, in the course of a severe cold, developed osteo-myelitis foci in a rib and tibia synchronously; these were followed in the next three months by foci in six other bones. The infecting organism was not mentioned. Even this case comes on the border-line between youth and manhood.

Moreover, the case is noteworthy from the bacteriologic standpoint also. Lexer¹ in his exhaustive article (1896-1897) collected only 20 cases of osteo-myelitis in which the streptococcus was the infecting organism; 10 of these had been previously reported to the French Surgical Congress by Lannelongue.⁷ Lexer remarks that the streptococcus, like the staphylococcus, shows a marked tendency to develop in the bones of the young. Lannelongue's cases varied in age from 10 days to 13 years. Of Lexer's 10 cases 8 were under 14 years of age.

In the examination of the pus from 73 cases of osteomyelitis Lannelongue and Achard⁸ found the streptococcus alone only 7 times.

Lexer has demonstrated that the clinical course of cases of streptococcus osteo-myelitis may not be unlike that of a staphylococcus infection. He divided streptococcus infections into three groups, in the first of which—containing those cases in which phlegmon was very marked in the affected bone and sequestration was found—the present case falls. Clinically it does not differ in any important item from the cases caused by the staphylococcus. Before Lexer wrote his paper Lannelongue had recognized a difference both clinically and anatomically between the streptococcus and the staphylococcus infections, and the symptomatology of the present case illustrates the fact (mentioned by Lannelongue) that both the fever and the pain in the streptococcus cases are apt to disappear on the third or fourth day. The improvement, however, is only temporary. Both symptoms return again.

It seems to be generally agreed that the pus in streptococcus cases is thinner and the necrosis of bone and soft parts less than in the cases of staphylococcus infection.

Klemm⁹ says that a widespread phlegmon of the medulla is rare in streptococcus cases and that edema of the soft parts is of wide extent. Because a single case can not be maintained as the standard, the fact that the pus in the present case was thick and that the medulla was widely involved is of no particular importance. Besides, secondary infection was present at the time I first saw the case.

But let the clinical or the pathologic or the bacteriologic aspects and differences be as important as they may, the fact remains that the important and imperative thing, the factor that means most to us and to the patient, is an early diagnosis. An early operation must follow an early diagnosis. The old principle of "Where there is pus, open and drain" holds here. Pus is in the medullary cavity and it is the medullary cavity

in the acute cases that must be opened and drained, that results may be achieved. Doing less than this is doing nothing. Temporizing, when a diagnosis has once been made, is criminal.

REFERENCES.

1. Archiv für klin. Chir., Bd. lvil.
2. Verhandl. der deutsch. Gesellschaft für Chir., xxiv. Kong., 1895.
3. Archiv für Kinderheilkunde, No. 22, 1897.
4. Quoted by Wenzel in Deutsch. Zeit für Chir., 1872, p. 410.
5. Beit. zur klin. Chir., 1880.
6. Canstatt's Jahresbericht, 1869.
7. Revue de Chir., 1895, p. 895.
8. Pl. Maucalre in Traité de Chir. (Dentu et Delbert), 1896, vol. xi, p. 520.
9. Quoted by Bloodgood in Prog. Med., December, 1899.

DISCUSSION

ON THE PAPERS OF DRs. BOTTOMLEY AND NICHOLS.*

DR. DE FOREST WILLARD, Philadelphia—It is not only the ordinary practitioner who is dilatory in diagnosis, but there are many surgeons who are slow in making up their minds and the golden opportunity is lost. Every case of osteomyelitis is infective, and we know that it needs as prompt attention as does a case of acute appendicitis. It should be recognized in the first forty-eight hours; it should be operated on in the first forty-eight hours. It should be operated on as soon as the diagnosis is made, whether it be of a streptococcus infection or a staphylococcus infection, and therein I would differ with Dr. Bottomley. In regard to the infection: I believe that the streptococcus infection is particularly serious, but in either case cutting inside the medullary cavity of the bone, freeing it from pressure and from infective elements is the only method which promises relief from total, or large destruction of tissue. The ordinary practitioner, and still less the surgeon, ought not to allow these cases to go on to destruction, suppuration and death. These cases are practically dying from sepsis when they come into our hands, and they have been treated for what? For that bane of progressive medicine, "rheumatism." If we could strike the term rheumatism out of the surgical world, it would be a blessing. One of the most potent destroyers of men, women and children is that cloak of ignorance, inattention and neglect, "rheumatism." How many cases we see dying day by day with an abscess containing ounces of pus, and the physician still treating it for rheumatism, or for typhoid fever. This is not an uncommon thing—every surgeon sees it over and over again. What is most needed is early, correct diagnosis. The treatment is plainly indicated. Where we have the evidences of rapid infection, with increasing pain, and deep-seated tenderness, there is only one rational treatment, that is to operate at once, relieve pressure and establish thorough drainage from the medullary cavity of the bone. The opening can not be made too quickly or too freely.

DR. J. CLARK STEWART, Minneapolis—I think it is most opportune that these papers should have been read to-day, because we have been hearing so much of the various forms of abdominal surgery, particularly that of ulcer of the stomach, gall-bladder disease and appendicitis, while it is rare to hear papers on osteomyelitis. I overheard a remark while in my seat, which I hope the gentleman who made it will pardon my repeating. When Dr. Nichols took the stage he said, "What, another paper on osteomyelitis; they must have a great deal of osteomyelitis in Boston," which reminded me of Dr. Oschner's story in regard to gall-bladder disease in Rochester, and I think that Dr. Nichols might answer very much in the same way that Dr. Mayo did: "We diagnose our cases of osteomyelitis in Boston." I think, without doubt, that osteomyelitis occurs in the practice of every man here, and possibly oftener than he recognizes, and there is no question that a very large percentage of the cases are primarily diagnosed erroneously. The diagnosis of these cases as rheumatism, which Dr. Willard mentioned, is probably the commonest error made, this being especially true of the subacute and chronic cases. Among the acute cases the diagnosis of typhoid is often made, and this error is a very serious one in causing delayed treatment. It

has been my lot to see several cases treated in a general hospital as typhoid until the abscess presented at some point, and this, of course, has not the excuse of the same diagnostic error earlier in the history of the case when sepsis may so mask the local symptoms as to make semblance to typhoid quite striking. The leucocytes ought to clear up all these obscure cases, as there is a decided leucocytosis in all acute cases of osteomyelitis. Early diagnosis of osteomyelitis is unquestionably the important point, and with the facilities at our command ought to be made in most cases without great difficulty. In the acute and chronic cases the difficulty of diagnosis is very greater, and frequently there is no diagnosis made until a lesion is discovered at the bottom of the superficial cavity. Fortunately, in these forms early operation is of so great importance as in the acute cases. In regard to Nichols' paper there were two or three points in which I am particularly interested. One was the instruction that in operations you should cut in the medullary cavity of the bone without interfering with the marrow. This seems to me dangerous as a general rule, for while I have no doubt in many cases operated on early, one is able to say where the marrow is infected, still judging from my own experience there are cases with multiple foci scattered throughout the bone where it would be extremely difficult without removing the marrow itself to determine the limit of the infection. As to Cushing's operation for the removal of the whole bone seems to me that this is unnecessarily severe and a misuse, for there are a great many cases of partial osteomyelitis in which a much less extensive operation will secure perfectly satisfactory results. In many cases you can remove the sequestrum, clean out the cavity and get primary organization of a blood clot, and in other cases by the method of packing, the final results are very satisfactory. Personally, I have had very little experience with the operation, and the results which he reports in cases of necrosis are certainly very brilliant, but I doubt whether it should displace the other methods which have stood the test of time.

DR. RICHARD H. GIBBONS, Scranton, Pa.—Those of us who have been taught by William T. Detmold, James P. Thomas M. Marco of New York, and D. Hayes Agnew, Grosses and our distinguished ex-chairman and teacher, Willard, need not fail to make a diagnosis of osteomyelitis early, or in much earlier times. When I first began practicing medicine, thirty years ago, Detmold taught that pain of character in bone was due to one of two causes, either inflammation, or inflammation that would result in abscess. If one will get pitting on pressure over the tibia, whether it is inflammation or not, or how extensive it may be, you will get but slight pitting if due to non-syphilitic inflammation of bone. Long since, I have been applying the expected bone the test which the dentist applies to the questionable tooth. As Dr. Murphy has said, you will not get pitting on pressure, unless it is long-continued, but you will get pitting on percussive pressure. However slight it may be, you will get a painful response by this method, even in the first hours of attack. The thing to do before opening up or trephining a flamed bone is to locate the seat of the infection, by tapping over its supposed site. I have had considerable experience in this kind of work in the anthracite coal region of Pennsylvania, where many of the boys engaged in and about the mines become infected, and I have chiseled out the bone throughout its length, and I have never had a bad effect, though I chiseled the bone all the way from the head of the tibia to the ankle joint. In one case I trephined and then entered the entire bone, down into the malleolus. This was in an old man, and he recovered. These cases ought to be treated well by operation, promptly performed, unless there is a local infection in connection with the trouble.

DR. J. B. MURPHY, Chicago—I wish to congratulate gentlemen on the practical points in the papers, and I wish to speak particularly on the second paper; I desire to still more forcibly emphasize the point of early diagnosis in acute

* Because of its length Dr. Nichol's paper will be published later.

Mayo, on Umbilical Hernia.

H. S. Cleveland

Figure 1.

A.S.Cleveland.

Mayo, on Umbilical Hernia.

Figure 2.

AS Cleaveland.

Mayo on Umbilical Hernia.

Figure 3.

tive osteomyelitis. What do we mean by early diagnosis in acute infective osteomyelitis? In what way is the physician of importance to his patients in a case of acute osteomyelitis? He is of importance to his patient in only one way, and that is when he makes a diagnosis inside of the first forty-eight hours. He is only of value to the patient when the case is diagnosed and operated on inside of the first forty-eight hours. Can it be done? Yes. The history of a case of osteomyelitis is so classic, so typical and so uniform that one can make the diagnosis ten hours after infection has occurred, or ten years, by interrogating an individual who was present at the time of the infection. What is the order of symptoms? The acute pain is usually preceded by a chill, followed by high temperature. The pain occurs usually in the extremities of the long bones, and in the majority of cases in children is not associated during the first forty-eight hours with effusion into the joint close to the focus of the disease, differing in this particular from rheumatism, which it is so uniformly erroneously diagnosed. The pain and sepsis increase hourly, but there is no local sensitiveness. Now then, what takes place in the third twenty-four hours after the infection? It is purely a symptomatic expression of pathology. During this period there occurs septic intoxication, acute absorption, high temperature, delirium, and all that goes with it. And with the advance of pathologic changes, with the increase of infectious products, with increasing pressure within the medulla of the bone from the bacterial invasion, we have destruction of the bone tissue, both by the toxic and biotic action of the infective flora, acting under great pressure. Rapid absorption continues until there is destruction of a vessel in one of the Haversian canals and the escape of the pus into the subperiosteal space. In the early stages of a case of acute osteomyelitis, there are no signs of inflammation on superficial pressure over the bone. I mean, during the first forty-eight hours, while the infective material is confined and restricted by the bony wall. In the third day of the infection, we usually have an escape of the pus from the center to the subperiosteal space, and sensitiveness to superficial pressure. But there is a sign that is an important one from a practical standpoint in the early stages; deep and continued pressure elicits pain and frequently indicates the position of the infection, even while the infection is still entirely within the bony wall. This will aid in making the diagnosis of the location of the focus of the infection. When the pus elevates the periosteum, there is rapid necrosis as the circulation is cut off from without and within. In forty-eight hours one may have complete destruction of the entire shaft of the bone. Therefore, if we are to be of service and beneficial to the patient, we must relieve that infection focus from the condition which is so dangerous in any tissue of the body, namely, infection under pressure, by making an incision down into the bone and allow the infective material to escape and thus avoid necrosis. We have just as perfect a guide to the source of infection as we find in many other cases of infection. One should drill a hole through the bone into the seat of the infection and allow the pus to escape from the bone, and thus relieve the intoxication and avoid necrosis.

If the general surgeon will get that window in the bone inside of the first forty-eight hours, he will save the patient from destruction of the bone, from the possibility of involvement of the joint, from the danger of fatal sepsis, and put him in condition for the line of demarcation to form, so that he will have but one small sequestrum. Then the case goes on and gets well, and the dead bone is easily removed through this window, after the separation of the sequestrum, usually in six weeks. Chills, high temperature, local aching and no sensitiveness in the first forty-eight hours, except on deep and continued pressure for one or two minutes, when the pain becomes excruciating, are very strongly in favor of acute osteomyelitis and demands as prompt surgical intervention as acute infectious peritonitis.

Dr. EDWARD H. NICHOLS, Boston—One or two points brought out in the discussion deserve some mention.

As regards the question of diagnosis: during the past five years I have had an opportunity to see many of these cases during the acute stage, at the Boston City Hospital, both on

my own service and on that of the other surgeons, who have kindly allowed me to see their cases, and I quite agree with all that Dr. Murphy says in regard to the desirability of early diagnosis, if we are to expect the best results. I do not agree, however, that the early diagnosis always is easily made. In some of the cases the typhoid symptoms come on so early that the patient is too stupid to notice the pain, and the cases may be treated as typhoid fever by excellent diagnosticians. In other cases the joint pain and swelling may be so extreme as to entirely obscure the bone affection, although monarticular pain, especially in children, always should arouse at least a suspicion of osteomyelitis.

In regard to the reparative power of the marrow: the integrity of the cortical bone is dependent on the integrity, both of the periosteum and of the endosteum. Extensive stripping of the periosteum usually produces considerable necrosis of the cortex; if the endosteum also is destroyed marked necrosis of the cortex is certain. Thence if one trephines into infected marrow and cures out the remnant of undestroyed endosteum he destroys the membrane which, in some cases at least, might regenerate the infected marrow, and maintain the integrity of the inner layer of the cortex. No one would, of course, remove the periosteum because he found an abscess between it and the shaft, and the same thing applies to the removal of the endosteum.

In regard to the removal of the necrotic shaft. It is not always necessary to remove the entire diaphysis. It may be sufficient to remove one half, or whatever portion of the diaphysis is necrotic. The removal of the entire diameter of the necrotic shaft is, however, much preferable to removing merely a shell of apparently necrotic cortex. When a mere shell of cortex only is removed it will be found, if the cases are followed for a number of years, that recurrences take place in a large majority of cases, and permanent cures are not common. The cases where removal of a superficial shell gives a permanent cure are usually cases of periostitis with little involvement of the marrow.

FURTHER EXPERIENCE WITH THE VERTICAL OVERLAPPING OPERATION FOR THE RADICAL CURE OF UMBILICAL HERNIA.*

WILLIAM J. MAYO, A.M. M.D.
SURGEON TO ST. MARY'S HOSPITAL.
ROCHESTER, MINN.

The results of operations for the radical cure of umbilical hernia in adults have, heretofore, not been encouraging. These patients are usually obese, with attenuated abdominal muscles, and the thin, rigid character of the ring does not offer mechanical conditions advantageous to lasting union.

The principles of closure have been the same as after an ordinary abdominal section, the object being to split the fascia at the ring margins laterally, until the recti muscles are reached, and then, by a series of buried sutures, to reconstruct the abdominal wall in layers. The defect in this method is the natural separation of the recti muscles at the level of the umbilicus. Below this point the muscular structures are practically in contact, but above there is from one-fourth to three-fourths of an inch separation. To bring the recti together in this locality amounts to a muscle transplantation. In a small hernia this muscular approximation is not difficult, and is the operation of choice, but in the larger openings an inch or more in diameter, it is impracticable, and, so far as I have been able to judge, the attenuated muscles, when secured, are

* Read at the Fifty-fourth Annual Session of the American Medical Association, in the Section on Surgery and Anatomy, and approved for publication by the Executive Committee: Drs. A. J. Ochsner, J. E. Moore and De Forest Willard.