round cells are scattered diffusely throughout the capsule. Extending from the capsule are trabeculae, projecting into the substance of the tissue, dividing the parenchyma into innumerable alveolar areas. The structure of these trabeculae resembles that of the capsule, except that the smaller divisions show degenerative

Fig. 5.—(High power drawing.) A, Tumor cells; B, blood vessel containing tumor cells; C, extravasation of blood into stroma; D, stroma; E, young blood vessel among tumor cells.

changes, due probably to pressure caused by growth of the parenchymatous tissue. The nuclei in the trabeculae are less deeply stained and the intercellular tissue of a lighter pink color. The blood vessels in these trabeculae

Fig. 6.—Meningeal involvement—dura invaded by tumor cells. (Photomicrograph.)

are interesting. Some of the vessel walls are composed of a single layer of endothelial cells, which are in direct contact with the fibrous structure of the trabeculae; these vessels are large and contain red corpuscles. Other

vessels are present which have exceedingly thick walls, very narrow channels, and contain but few corpuscles. In many of the blood vessels are large heavily stained cells, similar to those found in the acini of the tumor, and which are to be described later.

The parenchyma is arranged in round, oval or irregular-shaped areas, ranging in size from small microscopic divisions, to the size of a pea. These dark-staining masses with thickly beset cells, present a marked contrast with the paler enveloping trabeculae. The cells are large, round or epithelioid in shape and closely arranged. Their protoplasm in general is translucent, containing many irregularly arranged chromatic granules. Young blood vessels containing red-blood corpuscles are seen in the parenchyma; the walls of these are made up of tumor cells. Areas of degeneration, varying in size from small microscopic areas to .5 cm., are noted in many of the alveoli, especially the larger ones. These areas are of a hyaline-like character, stain pink, and present a homogeneous ground substance in which are imbedded transparent degenerated cells. They correspond to the areas described in the gross section as yellow-colored spots. The faint reticulum contains much blood.

A portion of the dura excised from where the membranes were involved in the tumor, showed, on microscopic examination, invasion by the same tumor cells described in the foregoing (Fig. 6).

The age of the patient, the rapid growth of the neoplasm, its histologic structure and the progressive course of the disease since the operation, are all in conformity with the diagnosis of sarcoma.

Results.—Three months after the operation the patient was found greatly debilitated, and presented an exaggeration of previous symptoms. Control of the bladder was entirely lost and involuntary bowel movements frequent, especially when the patient was asleep. He was unable to walk or even to support himself on his lower extremities. The pains in the limbs and back continued as before the operation, and emaciation was marked.

I am indebted to Dr. E. R. LeCount for valuable suggestions and criticisms during the preparation of this work, and to Drs. B. W. Sippy and A. D. Bevan for the clinical record of the case reported.¹⁹

RECENT ADVANCES IN ACTINOTHERAPY.

WILLIAM S. GOTTHEIL, M.D.

Dermatologist to the City Hospital and to Lebanon and Beth-Israel Hospitals.

NEW YORK CITY.

Young as is the treatment of disease by the various forms of radiant energy, there is already confusion in its terminology, and it will not be out of place to begin this review with the statement that the term "actinotherapy" is now the accepted designation for the method that employs the violet and ultraviolet spectral rays. The x-ray treatment is known as "radiotherapy," while "phototherapy" is employed for all kinds of light treatment collectively. There is need for insistence on exactitude in the employment of these terms, and even some of those who especially occupy themselves with phototherapeutics are open to criticism in this regard. Thus Freund entitles the authoritative work that he has just published "*Grundriss der Gesamten Radiotherapie*," including therein not only the Röntgen treatment, but also

19. I would mention, in addition to the authorities already cited, the following: Thorburn, W.; *Med. Chronicle*, 1889, x, 177-180; Fisher, W. W.; *Tr. Prov. M. and S. Assn.*, Lond., 1841, 2, x, 203; Oliver, J.; *Brain*, Lond., 1886-7, ix, 550-552.

that by high-frequency currents, Becquerel rays, visible and ultraviolet light rays, and even heat rays. The latter two agents he groups under the subheading of phototherapy. If the masters are not clear in their terminology the pupils will certainly go astray.

The scope of this retrospect is therefore restricted to the consideration of the progress that has recently been made in the therapeutic use of the ether vibrations that are appreciable to the unaided eye, together with the ultraviolet and chemically demonstrable ones that lie just beyond them. It practically includes only treatment with concentrated sunlight and with the various forms of the electric arc. It does not include treatment with the ordinary incandescent bulbs; and as there is a very prevalent misconception on that subject, it may not be out of place to refer to it here. The sun and the voltaic arc are the only efficient sources of actinotherapeutic energy that we possess. The incandescent bulb light has extremely little actinic power and penetration; nor can any practicable increase in its candle power or number give us a light that is comparable with even a very small arc in these qualities. The so-called light baths that are equipped with incandescents are practically sweat baths, and nothing more. The assertion that any actinic effect is gotten from them is entirely unsubstantiated, and I shall not consider them in this review.

A number of forms of apparatus, announced as improvements on the original Finsen type, have been introduced during the last year or two. Their aim throughout has been the very desirable one of simplifying and cheapening the apparatus, for it can not be denied that the Danish machine is complicated and troublesome to run, as well as expensive to install and use. I have had an opportunity to see and examine a number of the newer models, including the dermo-lamp, the Lortet-Genoud and the London lamp, and with the others I am familiar from the drawings and descriptions. They all run on small amperages and can be used on the ordinary incandescent wiring, and insofar possess very apparent practical advantages. They all give a superficial reaction; but they can not be expected to have any penetrative power. No attempt has been made to prove that their feeble light penetrates even the epidermis. In experiments recorded last year,¹ I proved that light of sufficient intensity from an arc could be made to penetrate not only the skin, but all the internal organs, and even the skin of the back after traversing the entire thickness of the body and then to influence a photographic plate. I used, however, a 50-ampere arc, giving a 20,000 to 30,000 candlepower light. Power is required for penetration; and without penetration, we can not expect to affect even the deeper layers of the skin, much less the internal organs.

This is a legitimate criticism of the new machines running on from 2 to 10 amperes of current. We can not hope to treat the deeper dermatoses with them, and it is entirely useless to attempt to affect a tubercular or carcinomatous focus in one of the internal organs. It is precisely in these fields that the future of the method lies. The few superficial dermatoses that are amenable to the action of the low-power instruments can well be treated by other means. In the latest publication from his institute,² Finsen is especially emphatic on this point. Leredde³ agrees with him, and says justly

that the intensity of surface reaction is no measure of the deep effect. The Lortet-Genoud apparatus gives a bulla in twenty minutes, but the light does not penetrate the tissues. It takes fifteen minutes for it to penetrate a rabbit's ear and affect a sensitive plate. Thinner living animal tissue than that it would be hard to find. With the powerful apparatus that I employ, a muscular human forearm can be penetrated in about the same time, and the chest or abdomen in two or three times the length of time required by the Lortet-Genoud apparatus to penetrate the rabbit's ear. The conclusions that I came to were that light from a source of sufficient actinic power and in sufficient concentration can be made to penetrate the entire thickness of the human body, including both layers of the skin, and that all the internal organs are accessible to its influence; and also that, since no organ or tissue can be more than half the thickness of the body from a cutaneous surface, and most of the internal organs are much less than that distance, the time required for efficient actinic penetration to any depth would be only a fraction of that required in the experiments; and conversely, if the time employed were equal, the chemotaxic effect would be much greater.

Both Finsen and Leredde conclude, and I thoroughly agree with them, that we must abandon the newer weak forms of apparatus, and return to those that concentrate large amounts of light. Of these only two are available. The original Copenhagen model is now purchasable in Europe, but it is hardly suited for any but institution use. It is extremely bulky, must be suspended from the ceiling, and requires an entire room. It is true that four patients can be treated at once; but as almost all the light comes from the crater of the arc, and since that crater is continually changing its position, it is evident that each of the four tubes radiating from the light will receive on the average only one-fourth of the light. This is the chief reason for the daily sessions of an hour or more for long periods of time that are required at the institute.

The instrument that I employ, while by no means small, is movable on castors and is suitable for the office. The full actinic force of the arc is directed on the area to be treated, and the only limit to the concentration of the circle of light is the efficiency of the cooling device in cutting out the heat rays.

Some legitimate objections to the earlier forms of the apparatus have lately been removed. The use of ordinary glass in the condensers and waterbath, which impedes the passage of the ultraviolet rays, has been abandoned; a Jena glass that is almost as transparent as rock crystal to them is now employed. Such, at least, is the assertion of the manufacturers. I can only state from careful comparative spectroscopic tests that it permits very much more of the ultraviolet to pass through than does the ordinary material. I also use a new carbon containing a large amount of iron and manganese in the place of the ordinary electrode. It gives a markedly violet flame that is entirely different from the usual light.

Of other improvements in the apparatus there have been but few. Broca⁴ has improved the compressor which is necessarily employed with the low amperage lamps to get any penetration at all. I have not found it necessary to use anything of the kind. De Beurmann⁵ applies a 1/1000 adrenalin solution to the part to expel the blood. He claims that it can be kept anemic for an

1. Medical Record, April 18, 1902.

2. Mitteilungen aus Finsen's Med. Lysinstitut, III, 1903.

3. Annales de Derm. et de Syph., December, 1902.

4. Annales de Derm. et de Syph., April, 1902.

5. Ibid., July, 1902.

hour in this way. Piffard⁶ has lately demonstrated an improvement on this method, using cataphoresis to drive the adrenalin into the tissues. The iron electrodes that Bang announced last year have appeared. They seem to work well for short periods of time with the smaller machines.

In the present state of our knowledge of the subject, I am prepared to say that it is quite useless to attempt actinotherapeutic work with anything less than 20 amperes and 55 volts; and that double that force is desirable if the internal organs are to be treated.

Some attention has recently been paid to the employment of sunlight in the place of the arc. Its applicability is limited, of course, since it can only be used during certain hours on bright days; but the absence of cost and the simplicity of the apparatus required are in its favor and I think will be extensively employed. The conditions are somewhat different from those with the arc. In the latter case we are dealing with rays diverging from a point, and it is desirable to approach the collecting lens as closely as possible to the source of light; and expense limits the size of the lenses in the machine. In sunlight we have practically parallel rays, and the amount of light collected is limited only by the size of the lens. Some ultraviolet light passes through all glass, and it is therefore possible to make the sunlight collecting lens of ordinary glass so large as to compensate for its deficient permeability, and make it equal in actinic power to a smaller rock crystal or glass lens. These considerations have led Finsen to employ ordinary glass and not rock crystal in his sunlight apparatus. The arc contains more violet and ultraviolet rays than does sunlight, on account of the atmospheric absorption; and the electric light can be used on any day and at any time of the day or night.

It would lead me too far to go into the consideration of all the various methods and apparatuses now employed, and I shall, therefore, proceed at once to a very brief consideration of the theory of actinotherapeutic action. The spectral rays that we employ are certainly bactericide. Velyaminoff⁷ has experimented very extensively with the most varied pathogenic and non-pathogenic organisms, and states that the micro-organisms are directly and rapidly destroyed by the light; and Nagelschmidt⁸ agrees with him. Sack,⁹ on the other hand, is more inclined to attribute the results obtained to the photochemical and inflammatory effect. There is also an undoubted effect on the vasomotor nerves. Capillary dilatation, as Bie¹⁰ has shown, is an invariable effect; and this is not a temporary matter, but persists for a long time, even for months, after the other effects of actinic action have passed. In fact, the latest investigations would tend to show that the rays cause a distinct endovasculitis after a time, and that, while slight action may cause vascular dilatation, the ultimate result is an inflammation of low grade with consequent permanent atrophy of the vascular tissues. We must conclude that the action of the light rays is a complicated one. It is, in part, directly bactericide and in part photochemical on the vasomotor nerves and the vascular structures. In addition to these, there are remoter effects in the stimulation of the tissue oxidation processes and new connective tissue formation.

In one respect the light action differs markedly from that of the x-ray, and that is a very important one from a practical point of view. The reactive tissue changes that it occasions are always moderate and controllable, and they appear promptly. There is never any delayed, excessive or uncontrollable effect. The condition of the hands of many of the Röntgen operators, and their insistence on their patients signing a release before beginning treatment, is sufficient evidence of the reality of the danger. A very considerable experience, now extending over several years, leads me to be positive on this point, and this with the most powerful currents available. The reaction can be pushed to vesiculation in a single session, if desired, and redness and swelling always occur. But all these phenomena disappear in a few hours or a day or two, and require, at the utmost, the application of a dusting powder. There are never any remote effects. Actinotherapy is perfectly safe.

Turning now to that portion of the subject that interests us most—the results obtained from the method—we are confronted with the facts that it has so far been employed by dermatologists almost exclusively, and that many of these used only the smaller and inefficient apparatuses to which attention has already been drawn. The report material is large, but not all of it is valuable. In lupus vulgaris, tuberculosis verrucosa cutis, tubercular ulcer, postmortem tubercle, etc., it is firmly established, not only as the most eligible, effective and innocuous treatment, but even as a test. Lesions that do not react and improve under it are not tubercular. Finsen's cases now number about 1,000, and almost all of them have been cured. His conclusions have been confirmed by many recent investigators, of whom I shall mention only a few. Klabowsky¹¹ reports 37 cases, including many old and intractable ones, that all did well. Gaston, Baudouin and Chatin¹² report 78 cases, one-third of them being completely and permanently cured; most of the others came irregularly or disappeared. There are similar experiences from a long list of investigators, among whom I shall merely mention Petersen,¹³ von Ziemssen,¹⁴ Velyaminoff,¹⁵ Schmidt,¹⁶ Sack,¹⁷ Danlos,¹⁸ Jessner,¹⁹ and Hopkins.¹⁸

I have treated a number of cases myself with most excellent results, but I have not as yet published anything about them. I have had sufficient personal experience, however, to entirely confirm the conclusions of the European investigators. The affections are cured with the smallest possible amount of tissue absorption; the cicatrix is often hardly apparent. This is a point of extreme importance, since the face is the part most often affected.

Lupus erythematosus is a new field for the actinotherapeutic method. An article will shortly appear embodying my own results. Out of four private cases I have had three cures, and the fourth one was very extensive and of many years' standing. The patient lived in a foreign country, and was only under treatment for a short time. Ravogli¹⁹ has treated a case of nine years' standing, in which the nose and both cheeks were involved. Finsen's latest report includes 31 cases, with 11 complete recoveries; and 19 cases were still under treat-

6. Medical Record, March 7, 1903.

7. THE JOURNAL A. M. A., April 12, 1902.

8. Archiv für Derm. und Syph., Nos. 2 and 3, 1903.

9. Dermatologisches Centralblatt, September, 1902.

10. Verhand. des Vereins f. Innere Med., Wiesbaden, April 15-18, 1903.

11. Wratch: Dermatologisches Centralblatt, July, 1902.

12. Wratch: Monatshefte f. prak. Derm., April 15, 1902.

13. Monatshefte für prak. Derm., June 15, 1902.

14. Ronssky Wratch: THE JOURNAL A. M. A., April 12, 1902.

15. Archiv für Derm. und Syph., March, 1902.

16. Münchener med. Woch., Nos. 13 and 14, 1902.

17. Annales de Derm. et de Syph., June, 1902.

18. THE JOURNAL A. M. A., Sept. 13, 1902.

19. Cincinnati Lancet-Clinic, Nov. 9, 1901.

ment at the time the account was written. Leredde and Pautrier had 23 cases; 3 were complete failures; 3 were improved; 6 had "segmentary cures" of the patches treated, and 11 were entirely cured. These authors have added some interesting details concerning their cases, which are worthy of note as emphasizing the intractability of the disease and enabling us to estimate the value of their results. They had had altogether not less than 802 galvanocautic séances, 382 scarifications, 462 high-frequency current sessions, without the cure of a single patch. Many of the authors previously mentioned record good results in the disease; a recapitulation is unnecessary. My own opinion is that at least one-third and possibly one-half the cases of this obstinate affection are curable by actinotherapy.

I have treated several cases of alopecia areata, and have on the whole been satisfied with the results. I believe, however, that it is very evident that under that designation are included at least two distinct affections, one of which is circumscribed, limited to the scalp chiefly, and benign in that it tends ultimately to spontaneous recovery, while the other is progressive, affects the entire body, and is malignant in the sense that treatment seems to have but little effect on its course. All my cases were of the former variety, but I can testify to a surprisingly rapid return of the hair in the patches treated. In some cases there was a good growth of lanugo hair after half a dozen sessions. Whether these cases are parasitic or not, our knowledge of the effects of light on the tissues would lead us to expect results; and I regard these cases as especially suited for this method of treatment. A number of other investigators give favorable reports, among whom Sack,²⁰ Schmidt,¹⁷ Hyde and Montgomery,²¹ and Leredde⁴ may be mentioned.

Actinotherapy has been lately employed in the various forms of birthmark, nevus telangiectasis, etc. We are practically helpless in these cases when they are at all extensive; and, as the face is often affected, any method that promises relief is welcome. The endovasculitis excited by the light finally leads to cicatricial contraction and obliteration of the redundant vessels. A number of the writers above mentioned report successful cases, but I have not yet had occasion to try it in a suitable one. The difficulty has been that all my cases have been infants or very young children, whom it has been impossible to keep quiet while the light has been applied. I have tried in two cases to put the child to sleep with the aid of the breast or the nursing bottle; but this is an expedient that can not be employed with the regularity and for the length of time that is required. I have, therefore, been compelled to advise the postponement of treatment until the children are older. There then would be no difficulty in applying it, and I should certainly recommend its trial in adults, more especially in extensive cases of port-wine mark and non-hypertrophic telangiectasis.

A year ago Leredde, Bie, Blacker and Sequiera, Sjögren, Helm, and others reported cures of various cases of carcinoma of the skin, epithelioma, rodent ulcer, and Paget's disease with the light. That the more recent literature of the subject is small is probably due to the fact that radiotherapy has been so extensively employed in this class of cases, and that many experimenters have been using the smaller low-amperage apparatuses, to the inefficiency of which I have already called

attention. Leredde⁴ and Velyaminoff¹⁶ have employed the more powerful machines with results at least as good as those obtained from the x-ray. I have no results to report as yet. My cases have been either extensive ones in hospital practice, in which, for reasons already stated, the treatment could not be employed, or private ones in which the circumstances precluded the use of a method that takes time. Further and extensive experimentation is required. The treatment is rational, in view of what we know of the action of the light, but actual trial only can decide as to its efficacy.

I have had one case of sarcoma cutis under the treatment for some time. It is quite extensive, however, and only a small portion of the growth can be treated at a time. The results have been fairly good. Individual nodules and tumors show distinct retrogressive changes under the influence of the light. The method is only applicable, of course, to the cases showing a single or only a few tumors. The disseminated cases, in which large areas or the entire cutaneous surface is affected, can not be treated in this way.

Rosacea, especially when the stage of permanent vascularization has been reached, would seem to be a good field for the employment of actinotherapy for the same reasons as in nevus. Finsen reports 15 of these cases, with 4 cures, 5 marked improvements, 1 still under treatment, and the remaining 5 as having stopped treatment after a short time. Bang,²² and a number of others, report good results. I have used it in a number of cases of acne rosacea, but found it too slow to employ alone. It was generally employed in connection with curetting and local applications. I shall endeavor to use it exclusively in the next bad case of acne pustulosa that I meet.

In the parasitic skin diseases, ringworm, favus, sycosis and pityriasis versicolor, Von Ziemssen²³ and Leredde²⁴ have done well with the method. The same authorities report good results in psoriasis and rhinophyma. I have not been able to do anything with it in psoriasis; and we possess other simple and effective means of treating the disease. A number of investigators have used the method in various eczematous affections, in burns and wounds, in tropical ulcer, etc., but neither their number nor their results warrant their inclusion in this report.

As regards the employment of actinotherapy for the internal organs, the only reports as yet are those of Freudenthal²⁵ and Williams.²⁶ The former has found that the light is a most important factor in combating the symptoms of hay fever, and a valuable adjuvant in the treatment of pulmonary tuberculosis. The author takes a very conservative position in the matter, as is entirely proper in view of the very limited amount of work that has as yet been done in this field.

I can only say that the method is a promising field for investigation. The bactericide and tissue stimulating action of light is proven, as is also the fact that it can be made to penetrate the body and reach the internal organs. Whether its properties can be successfully employed for the treatment of diseases of the internal organs must be left for future investigation to decide.

22. Berliner klin. Woch.: Monatshefte f. prakt. Derm., April 15, 1902.

23. Ibid., June 15, 1902.

24. Presse Médicale: Ibid., Sept. 15, 1902.

25. New York Medical Journal, July 12, 1902.

26. Dietetic and Hygienic Gazette, October, 1902.

20. Münchener Med. Woch.: Monatshefte f. prakt. Derm., Oct. 1, 1902.

21. THE JOURNAL A. M. A., Jan. 3, 1902.