mouth-breathing or nose-breathing appliances, to consider any but the respiration chamber method. While one may say that the placing of a subject in an enclosed chamber for the purpose of studying the respiratory exchange is a distinctly abnormal procedure, repeated experiments with the large calorimeters have shown that, except in very rare cases, the subject immediately adjusts himself to the conditions and frequently sleeps through the entire experimental period. On the other hand, the insertion of the nosepieces or the mouthpiece is almost invariably followed by a distinct alteration in the type of respiration, even though often there are no external evidences of discomfort. This is no more sharply shown than when pneumograph records are secured of the respiration with the subject breathing normally and comparisons made with records obtained with the subject breathing through a mouthpiece or nosepieces. Very noticeable alterations of the respiratory type are almost invariably found, especially with untrained subjects. Furthermore, personal conversation with many physiologists, both in America and in Europe, has led us to believe that it is the almost universal opinion that breathing through nose or mouth appliances, save in the case of exceptionally well-trained subjects, produces an abnormal type of respiration.

Consequently, in deciding upon a method for studying the respiratory quotient to be developed in the Nutrition Laboratory, the original plan was to construct a chamber which would meet all of the conditions requisite for successful experimentation and use it attached to the universal respiration apparatus. This method seemed especially feasible, as smaller chambers had been successfully used with this apparatus in studies with dogs and infants. In devising this respiration chamber we profited by the previous experiences of the Laboratory with the large chambers used in the respiration calorimeters and attempted in so far as possible to eliminate all particularly troublesome factors.

Recognizing, also, that any form of apparatus so complicated and costly as the respiration calorimeters in the Nutrition Laboratory and the Russell Sage Institute of Pathology was practicable for only a very few hospitals, the attempt was made to develop a respiration chamber for use with the universal respiration apparatus which would provide for the measurement of the carbon dioxide production and the oxygen consumption of patients in relatively short periods and still not be prohibitive in cost. It was not possible in such an apparatus, therefore, to introduce costly appliances for the direct measurement of the heat output. Fortunately, however, almost all of the evidence that has accumulated at the present time points to the fact that the larger proportion of experiments may well be made by determining the heat production indirectly from the measurements of the carbon

dioxide production and the oxygen consumption, provided that these are accurate.

A detailed description of the apparatus as finally developed and a number of results obtained with it on pathological subjects are given in Part II of this paper.

REFERENCES.

¹ Carpenter: Carnegie Institution of Washington Publication No. 216, 1915.

(To be continued.)

STUDIES OF THE BASAL METABOLISM IN DISEASE AND THEIR IMPORTANCE IN CLINICAL MEDICINE.

BY J. H. MEANS, M.D., BOSTON.

Henry P. Walcott Fellow in Clinical Medicine, Harvard University.

[From the Medical Service of the Massachusetts General Hospital.]

IN a recent paper, Professor Lusk has referred to the calorimeter as the interpreter of life processes.¹ This might be broadened to calorimetry, and calorimetry, as I shall presently explain, can be carried out without a calorimeter.

What is calorimetry?

As generally used at present, the term includes (a) the determination of the heat given off by an organism, by actual measurement; that is, the heat elimination; and (b) the calculation of the heat production from the gas exchange. The former is called direct, the latter indirect calorimetry. Direct calorimetry can be done only by means of the calorimeter, an instrument costing several thousand dollars; indirect calorimetry, on the other hand, can be carried out with any of a number of different types of respiration apparatus, the cheapest of which need not cost over \$75.*

Studies of the gas exchange in man and animals have been made by scientists since the days of Lavoisier, but it is only comparatively recently that Magnus-Levy,² and more particularly the workers at the Russell Sage Institute at Bellevue Hospital, New York,³ have shown the importance of calorimetry in internal medicine.

The elimination of heat and the production of heat, as determined by direct and indirect calorimetry, respectively, usually agree within a few per cent. so that studies by the latter method alone will serve most purposes. When one obtains the elimination of heat by direct calorimetry he expresses the result in heat units—calories. If, now, we wish to calculate the heat production in calories, we may do so by determining the oxygen absorption and the respiratory quotient. The latter is the volume of carbon dioxide eliminated divided by that of the oxygen absorbed in any given time unit.

* I refer to the Douglas respiration apparatus, which may be had from Siebe, Gorman & Co., London.

The respiratory quotient is of importance because it is an index of the kind of material being burned in the body. In the case of any given food substance, whether burned in the body or in a retort, the consumption of a definite amount of oxygen will result in the evolution of a definite amount of carbon dioxide. In other words, the ratio $\frac{\text{CO}_2}{\text{O}_2}$ will be a constant, for any given

substance, but will vary with different substances according to the structure of their molecules. The more highly oxidized a substance is at the start, the smaller the amount of oxygen that will be required to complete the oxidation, hence the higher will be the ratio $\frac{\text{CO}_2}{\text{O}_2}$.

Of the three classes of substances that may be burned in the body, carbohydrates, fats and proteins, we know the respiratory quotients. For example, in the case of grape sugar, which has an empirical formula of $\text{C}_6\text{H}_{12}\text{O}_6$, and which is completely oxidized according to the reaction

there will be given off the same number of molecules of CO_2 as were absorbed molecules of oxygen (and in the case of gases according to Avogadro's hypothesis, equal numbers of molecules make equal volumes at the same pressure and temperature); hence the respiratory quotient, $\frac{\text{CO}_2}{\text{O}_2}$, of grape sugar is 1.00. In the case of fats,

$\frac{\text{O}_2}{\text{O}_2}$ however, substances less highly oxidized than sugars, but also completely burned in the animal body, relatively more oxygen will be required. With sugar, oxygen was required to oxidize the carbon only; with fats some is also required to oxidize the hydrogen. Therefore the respiratory quotient of fat will be below unity. As a matter of fact, it is .71. Proteins occupy an intermediate position, having a quotient of .81. Knowing the theoretical quotients, then, of carbohydrates, fats and proteins, and having obtained a subject's respiratory quotient by experiment, we can calculate how much of his total heat production is due to the combustion of carbohydrate, of fat and of protein.†

In addition to knowing the respiratory quotients of the three classes of foods, we also know their heats of combustion, and since they produce the same amount of heat when oxidized in the body as when burned in a retort, it has been possible to calculate the value in calories of a liter of oxygen for any respiratory quotient between .70 and 1.00.‡

Hence if we know the respiratory quotient and also the oxygen absorption, we can calculate the heat production thus:

† In the case of proteins this statement is not strictly true, for proteins are not, like fats and carbohydrates, completely oxidized. To be absolutely accurate we should have to take the urinary nitrogen into account. For routine work, however, the protein metabolism may be disregarded since the R. Q. of the non-protein katabolism differs very little from that of the total.

‡ For these values see Williams, Riche and Lusk: Jour. Biol. Chem., Vol. xii, p. 357.

Oxygen absorption per hour	×	Calorific value of oxygen for the R. Q.	=	Calories per hour
----------------------------------	---	---	---	-------------------------

which, in case the oxygen absorption were 12 liters an hour, and the respiratory quotient were .85, would become

$12 \times 4.863 = 58.4$ calories per hour, 4.863 being the calorific value of oxygen for a quotient of .85.

So much for methods of calculation. The next problem is in regard to what constitutes the normal metabolism and what factors may alter it. It is not my purpose in the present communication to discuss these questions, it is merely to point out their possible clinical importance. To answer the second question first, I may say that the taking of food, the doing of muscular work, and the exposure to cold, are the common factors which cause an increase in heat production in the normal individual. In the field of pathology, fever, acidosis and certain disturbances in internal secretions may also cause a rise.

Now it will be asked what constitutes the normal heat production, and is it a constant or a variable? It is very definitely a constant provided we use for comparison the minimum or, as it is more commonly called, the basal metabolism. By basal metabolism is meant the heat production when all the extraneous elements previously mentioned, food, muscular work, cold, etc., are eliminated. We can determine this basal metabolism in practice by having the subject at complete muscular rest, in the post-absorptive condition (that is, fourteen or more hours after food) and in an ordinarily warm room. If the effect of food, cold and muscular work are ruled out the metabolism will be found in any individual from day to day and from year to year (provided his weight remains the same) to vary very little. Further, it has been found that the metabolism of one organism is comparable with that of another, provided the total heat production of each is divided by the surface area of their bodies. This law was shown by Rubner to hold for all warm blood animals. Per unit of surface area, a mouse, a fowl and a man have the same basal metabolism.

Recently Du Bois⁴ has made it possible to determine the surface area in man with great accuracy by means of calculations based on a number of linear measurements of the body. Using this formula, I have found the average basal metabolism of 16 normal men and women to be 38.8 calories per square meter an hour.⁵

This figure agrees quite closely with that obtained at the Sage Institute, and the average variation from it was only $\pm 4\%$, and the extreme, less than $\pm 10\%$. Following the Sage investigators, therefore, I shall consider any metabolism abnormal that is more than 10% from the average figure.

FIG. 1. The basal metabolism of a series of normal men and women. The black columns represent the metabolism per square meter and hour, the shaded columns, that per kilogram and 24 hours. The solid horizontal line represents the average of the series; the interrupted lines, a 10% variation above or below the average. The subsequent diagrams are all constructed on the same plan as this one. The base line in each case is zero, so that the height of the columns indicates the actual amount of heat production.

That the metabolism is more nearly proportional to the surface area than to the body weight will be seen by a glance at Fig. 1. The black columns representing the metabolism per square meter, are more nearly the same height than the shaded, which represent that per kilogram of body weight. This fact becomes more striking in Fig. 2, where the metabolism of two very fat women is contrasted with that of a woman of normal build, and a rather thin man. The four black columns representing the surface area relation are all within 10% of the average, whereas the shaded columns representing the weight relation are, in the case of the fat subjects, 40-45% below the average.

It may now be asked of what interest is all this to the internist. A little consideration will make it apparent that it is of very distinct interest. No one will deny the importance of the body temperature, of the pulse, or the blood pressure, yet about the only information we get from the last of these is that if constantly elevated the patient probably has chronic nephritis, or arteriosclerosis, or both, facts that we could probably arrive at by other means.

In the basal metabolism we are dealing with a factor nearly as constant as the body temperature, and because of the light it throws upon the calorific requirements of the body, quite as interesting.

For example, the basal metabolism of a 60-kg.

man might be 1,500 calories a day. This would be his heat production if he lay quietly in a warm room all the twenty-four hours. When he leads a life of moderate activity this may become 3,000 to 4,000 calories. Now in severe Graves' disease the basal metabolism may be doubled, so that the same man if he had that malady might have a basal metabolism of 3,000 calories and an actual metabolism very much higher. This at once shows that in order to keep patients with this disease in equilibrium they must be fed very many more calories than a normal individual of similar size and under like circumstances. The same applies in typhoid fever as shown by Coleman and Du Bois⁶ and hence the advantage of their high calory diet.

During the past two years I have had an opportunity to make various studies of the basal metabolism in the medical wards of the Massachusetts General Hospital. The apparatus used for the oxygen and carbon dioxide determinations was that of Benedict.⁷ After a series of normal individuals had been secured the work lay chiefly in a study of thyroid disease and of obesity.

Thyroid cases are particularly interesting since in hyperthyroidism there is a marked rise, and in hypothyroidism a marked fall, in the basal metabolism. In Fig. 3 are shown a series of nine cases of typical exophthalmic goitre. The metabolism in all of them shows a marked rise,

FIG. 2. The basal metabolism in obese and normal subjects, showing the close relationship of the metabolism per unit of body surface and the wide divergence of that per unit of body weight.

FIG. 3. Basal metabolism in Graves' disease. The third column in Case 1 was after one x-ray treatment, and the third in Case 2 was after hemithyroidectomy.

anywhere from 30% to 80% per square meter, and still more per kilogram, this latter being due to the great emaciation that many of them show.

This marked rise in hyperthyroidism is very interesting because, so far as is known, so marked a rise occurs in no other condition in which fever is not present. These patients were all afebrile when studied. A slight rise in the metabolism may occur in acidosis, that of diabetes or of uremia, for example, but it seems likely that a marked rise (50% or more) may be used as factor in the differential diagnosis of hyperthyroidism.* Early Graves' disease is sometimes confused with early phthisis. I have not as yet been able to make any determination on the latter, but one advanced cases of phthisis that I observed showed a normal metabolism.

In addition to the fact that an increased heat production is a characteristic of hyperthyroidism, there seems to be good evidence that the severer the intoxication the higher the metabolism.

Case 1, a middle-aged woman, on whom three observations are reported, was a very severe case with all the classical signs of Graves' disease, marked tachycardia, tremor, nervousness and eye signs. The third observation was about three weeks after the first, and during the interim she had been on the hydrobromide of quinine and in addition had received one x-ray

* A slight rise in the metabolism, so Dr. Du Bois writes me, occurs in such conditions as acromegaly, dyspnoea, anaemia, and cancer, but these conditions can easily be differentiated from hyperthyroidism.

treatment. A moderate fall in the metabolism had occurred.

In Case 2, also a severe one, the first two observations were made before treatment was instituted and the last three weeks after a hemithyroidectomy. Here, again, the metabolism decreased somewhat. In both cases the fall was coincident with subjective and objective improvement.

The three observations on Case 3 extended over a period of five weeks, during which time she was getting x-ray treatment. No fall in the metabolism took place, though clinically she was better. She was not so ill at the start as Cases 1 and 2.

Case 4 was a very advanced and malignant type with great emaciation, who later died following ligation of the thyroid arteries.

Case 8 was an early case with only slight tachycardia and eye signs, but with extreme nervousness and definite thyroid enlargement.

Cases 5, 6, 7 and 9 were all moderately severe cases, none of whom had had any treatment when the determinations were made. Of these Case 5 deserves comment because she had had Graves' disease seven years before but had apparently been cured by hemithyroidectomy followed by x-ray treatments. A few months before my observation was made, symptoms of Graves' disease had returned and at the time of examination there was marked enlargement of the remaining half of the thyroid, with bruit, eye signs, tremor, tachycardia and nervousness.

Reference to Fig. 4 will show that in myx-

FIG. 4. Myxedema and atypical thyroid cases.

edema the metabolism shows the reverse change to that found in hyperthyroidism. Cases 10 and 11, both typical myxedema, who had never taken thyroid gland, showed a marked reduction. The last two columns in Case 10 were after thyroid administration had been begun and show a tendency for the metabolism to increase toward normal.

Case 12 was a patient with a large cystic goitre of long standing, who recently had begun to have symptoms of hyperthyroidism, tachycardia, tremor, nervousness, etc., but no eye signs. The high metabolism found makes it seem certain that she actually did have some hyperthyroidism.

Cases 15 and 16 had both had typical Graves' disease in the past, but were completely recovered when observed; both showed a normal metabolism.

Cases 17 to 20 were obscure cases where hypothyroidism was suspected and the test was done as a means of differential diagnosis.

Case 20 was a tachycardia of unknown origin. Electro-cardiography pointed toward a cause outside the heart, and that, together with the increased metabolism and nervousness, led us to consider it a case of hyperthyroidism, even though typical signs were lacking.[†]

The second group of cases that I have given most attention to is that of obesity. In Fig. 5 the results obtained in ten cases are shown.[‡]

At this point I must mention the fact that with increasing years the metabolism gradually declines[§] so that it is not fair to compare the metabolism of a person of fifty years with my average figure of 38.8 which was obtained from

[†] These thyroid cases will all be published in full in a subsequent paper.

[‡] Of these ten, Cases 21, 24, 25 have been published. (See Means, J. H., Jour. Med. Res., Vol. xxxii, p. 121, 1915.) The others will be published in detail in the near future.

a series of persons between twenty and thirty. Some of the obese subjects were well over thirty (Cases 21, 23 and 25) and for them 35 calories per square meter and hour would be a fairer standard than 38.8. Using 35 calories as a standard, then, it will be found that Cases 21, 23 and 25, all of whom were enormously obese, were well within 10% of the standard. In Case 22 some of the observations fall slightly below the —10% line. So also do Cases 24 and 26, but not nearly to so marked a degree as the two cases of myxedema. Cases 27 to 30, which comprise all the men, are within 10% of the normal. In Cases 24 and 26 there was some clinical evidence that a disturbance of internal secretion was present. Neither of these cases would be considered more than moderately obese in comparison with Nos. 21, 22, 23 and 25.

A few miscellaneous cases were collected. One of Addison's disease, one of myasthenia gravis, one of Raynaud's disease all showed a normal figure. A case of pernicious anemia with a hemoglobin of 50%, also showed a normal metabolism. In pernicious anemia with a lower blood picture, Meyer and Du Bois have found a slight increase due, they think, to an increased blood flow and utilization of blood oxygen.

In regard to the kind of katabolism in these various conditions, as shown by the respiratory quotient, we cannot say a great deal. Benedict[§] has shown that the *nüchtern* quotient in man is influenced to a marked degree by the amount of carbohydrate in the previous diet. As in the majority of these cases the diet was not measured we can draw no definite deductions.

In conclusion I should like to emphasize the following points:

1. Basal metabolism can be readily studied

FIG. 5. Basal metabolism in obesity. Cases 21, 23 and 25 being considerably older than the subjects in the normal series, it is probable that for a standard we should take 35 calories per square meter and hour, which would make them lie within 10% of the normal, as explained in the text.

in a hospital clinic with comparatively inexpensive apparatus.

2. The normal basal metabolism is a fairly constant affair, and hence wide variations from it in disease are of interest to the clinician.

3. A marked rise occurs in hyperthyroidism.

4. A marked fall occurs in hypothyroidism.

5. In regard to hyperthyroidism it seems probable that the basal metabolism furnishes

(a) The best index as to the severity of the disease, and hence is a quantitative means of following the course and of judging of the effectiveness of treatment; and

(b) A valuable aid in differential diagnosis.

6. Enormous grades of obesity are possible in the presence of a normal basal metabolism.

7. When a reduction in the metabolism was found in obese subjects there was also clinical evidence of defective internal secretion.

8. A clearer conception of the food requirements in disease is furnished by the basal metabolism than by any other factor.

REFERENCES.

- ¹ Lusk, G.: *Science N. S.*, Vol. xiii, p. 816, 1915.
- ² Magnus-Levy: *Zeitschr. f. klin. Med.*, Vol. xxxiii, pp. 258-313, 1897.
- ³ *Arch. Int. Med.*, Vol. xv, No. 5, Part II.
- ⁴ Du Bois, E. F. and D.: *Arch. Int. Med.*, Vol. xv, p. 808, 1915.
- ⁵ Means, J. H.: *Jour. Biol. Chem.*, Vol. xxi, p. 263, 1915.
- ⁶ Coleman and Du Bois: *Arch. Int. Med.*, Vol. xiv, p. 168, 1914.
- ⁷ Benedict, F. G.: *Deutsch. Arch. f. klin. Med.*, Vol. cvii, p. 160, 1912.
- ⁸ Magnus-Levy and Falk: *Arch. f. Anat. u. Physiol.*, Suppl. 1899, p. 314; also Du Bois, E. F.: *Harvey Lecture*, Nov. 27, 1915.
- ⁹ Benedict and Higgins: *Influence on the Respiratory Exchange of Varying Amounts of Carbohydrates in the Diet*, *Am. Jour. Physiol.*, Vol. xxx, p. 217, 1912.

THE PHYSICIAN AND THE PREVENTION OF INDUSTRIAL ACCIDENTS.

BY HERBERT J. CRONIN, M.D., CAMBRIDGE, MASS.

WITH horror we read tales of the jungle where the wild beast leaps from his lair; seizes a human victim, and slowly devours him in spite of his futile efforts to escape. Mentally we picture his struggles to avoid the pitiless jaws, hear the crunching of bones and the weakening cries for help. Our imagination is fired and we fancy ourselves shooting the beast and rescuing the unfortunate. Then, it is dismissed from our minds as merely fiction,—such things do not happen nowadays. Yet almost daily in this country a workman will be seized by a huge machine and drawn into its cruel crushing gears. It is more powerful, merciless and destructive than the beast; it quickly kills or horribly mutilates. A campaign is being waged for the protection of our workmen from such disasters. It was first brought to the attention of the public by the Safety First slogans. Accident prevention in industries is now a topic of first importance; yet

the medical profession has not been actively enlisted in this campaign nor has there been an organized effort to interest them.

Until the passage of the Workmen's Compensation Acts, accident prevention was given little consideration. Labor was cheap, and if a man was injured, it caused but an expression of passing regret or sympathy; he was dropped from the payroll and another filled his place. He was allowed to seek his own help; the business that caused his injury had no interest in him. Labor was scrapped in the same way as the machines it worked upon. We had a superabundance of foreign labor, which has been one of America's greatest assets; it has done the hard, laborious work of the country and our mines, steel factories and mills have depended upon it. It is fortunate that we have started this campaign

FIG. 1. Shows how a boy's arm was lost on a mule-spinning frame.

to conserve labor, for it is doubtful if we shall receive many men from Europe after the war. They will be kept home to build up their devastated countries.

For years plants have taken precaution against fire. They have had local fire organizations, apparatus, sprinklers and other safeguards. They were proud of their efforts because they not only protected the plant from destruction by fire but helped to reduce the fire insurance rates. Yet the same firms did nothing to safe-guard the lives of employees from death-dealing machinery. Employees were expected to look after themselves and it was their own loss if they were injured. If the machine was dangerous to operate, they must use all due care; but no attempts were made to make a dangerous machine safe until the advent of the Compensation Laws. Now the costs of the hazards of an industry are charged up to that industry. Many accidents mean increased premiums. It is business policy to try to prevent accidents and place