indications for uterine irrigation and drainage are not difficult to determine, as every text-book and journal treating of the subject recommends irrigation and efficient drainage, so that the question confronting most of us is not *when*, but *how*, to employ this measure; the solution of which is left entirely to our own resources.

Should I now ask you, what are the best means at our disposal for securing uterine drainage, if I mistake not, your answer would unanimously be, iodoform gauze. This substance is the material *par excellence* to carry off primary wound secretion, as well as a valuable tissue stimulant and drain in tubercular or other fistulous tracts, and almost an indispensable mechanical hemostatic and uterine stimulant after curettement; but, as stated by competent observers, instead of acting as drainage in the uterine cavity, it is simply expelled or acts as an obstructing plug to the discharges.

Next Walter S. Wells' spiral wire tube, filled with wicking and supposed to establish osmosis between the uterine cavity and a medicated sponge placed in the vagina, has proved to be a failure.

Then comes Cherron² with 20 to 100 strands of silkworm gut introduced on a special holder, the free ends of the gut being fitted to a celluloid plate. This may in a few cases prove of some value as a capillary drain.

Combined Uterine Irrigator and Drainage Tube.³

We have Bolt's and Wylie's hard rubber drainage tubes, to both of which the same objections are found applicable. In gynecology these tubes answer better than in puerperal sepsis, but in either case they fall short of our expectations; first, on account of the grooves being partially closed by the tissues pressing into them; second, and the most serious objection to their use, is occlusion by the vagina falling against the free outlets.

In a recent case of puerperal sepsis, where one of these tubes had been placed in the uterus for twenty-four hours, its withdrawal and the simple introduction of an English catheter was followed by the discharge of more than an ounce of exceedingly foul sanguino purulent fluid. Of the two, on account of its freer outlet, Wylie's tube is preferable; but who would think of using a contrivance of this kind for drainage in a surgical wound?

I will here show you a Manton tube which is the best of any on the market, as you will observe it has a truncated extremity and a slot nearly its whole length, but it has the same objectionable outlet as the two last named—hence is of but little practical value.

I now present an irrigating drainage tube, constructed of vulcanized rubber or platinum which has, so far as it has been used, proved to answer every indication for which it is intended. It is at present the ideal uterine drainage tube because:

1. Its outlet can not become obstructed and over-distend the flabby uterus.

2. It is the only tube through which irrigation may be accomplished and that will admit of continuous uterine siphonage or irrigation in those very virulent cases where life would be extinguished in a day or two.

3. By no other means is it ever safe to irrigate the uterus without the use of a speculum, as it is never permissible to pass an instrument through the vagina by the sense of touch directly into the uterine cavity, although this folly is sanctioned by high authorities.

4. We should never trust even a trained nurse to irrigate the uterus, but after once introducing this tube, a nurse can be instructed to use an antiseptic vaginal douche and then attach a No 7 English catheter—from which an inch or more of the tip has been removed—to a fountain syringe, turn on the stream, and guide the catheter into the tube. As the cavity of the tube gradually tapers from the last drainage or return fenestra to the irrigation apertures the catheter will, by introducing it until resistance is felt, fit snugly and throw the fluid into the uterine cavity, returning it through the irrigation openings. In this way, by placing a patient on a douche pan, intra-uterine irrigation may be resorted to, without the least inconvenience or exhaustion to the patient, as removal of the tube for sterilization, as a rule, oftener than every twenty-four or forty-eight hours is found unnecessary even in case of puerperal sepsis. In non-puerperal patients, removal of the tube is not so often indicated. Whatever the indication for which the drainage tube is used, the patient must be kept absolutely at rest until its removal.

In conclusion, let it be understood that I do not wish to go on record as advocating drainage in every case of endometritis, dysmenorrhea, etc.; in fact, its indication is rather the exception than the rule. I am aware, too, that even the majority of cases of puerperal infection will, sooner or later, recover without any local treatment whatsoever; again, a few antiseptic vaginal douches are all-sufficient; or every indication may be fulfilled by one thorough uterine cleansing. But that there are rebellious cases which tax the physician's skill to the utmost, can not be doubted, and where alertness to the danger signals, and effectual and radical treatment, is the lying-in patient's only safeguard. It is equally true that physicians who scout at the idea of all intra-uterine medication and who, in a practice of ten or twenty years, never invade this organ, are far from being alive to their patients' welfare.

When any treatment is to be directed to the endometrium two points, however, should under all circumstances, be kept in mind—namely, gentleness in manipulation and conscientious antiseptic precautions.

TUBERCULOSIS.

Read before the Pennsylvania State Medical Society, 1895.

BY J. M. BATTEN, M.D.

PITTSBURG, PA.

I wish to show, by clinical observation, that the tubercular diathesis, predisposition to, or habit, is inherited to the third and fourth generations, and that consumption has been but very slightly, if at all contagious among those who came under my observation

² Gazette des Hôpitaux, October, 1892.

³ Manufactured by the Aloe & Penfold Company, Omaha, Neb.

who have not inherited that predisposition to the disease.

Mrs. G. had five children. She and four of the children died of phthisis pulmonalis. The remaining child, a female, lived till the age of 84 years and bore ten children, five of whom died of phthisis pulmonalis. Three of the latter were married and had children. The eldest a male had four children, three of whom died of phthisis pulmonalis—one is still living. The second one, a female, married and had one child, and within a year after the birth of the only child, the latter died of children's disease and the parents died of phthisis pulmonalis. The third, female, married and had one child, a male, who is still living. The mother died young and the husband died recently of heart disease. Two females, unmarried, died young of phthisis pulmonalis. Of the five living, the eldest had ten children, two of whom died of typhoid fever. The remaining children are living but delicate. The mother is 75 years old and has paresis. The second, a female, married, had five children all living and healthy. The third, a male, in his younger days had hemoptysis, but removed from the city to the country and is still living. He is married and is the father of five children, all delicate. The fourth, a female, married a delicate man who afterward died of phthisis pulmonalis. She is the mother of five children, all delicate. The fifth, a male, is still living at the age of 58 years. He has hemiplegia and is the father of three children, small in stature but healthy.

One case of another family, Mrs. M., who died of tuberculosis, had borne four children, three of whom died young; two of them of children's diseases, the third at the age of 20 years, of phthisis pulmonalis. The mother, Mrs. M., died at the age of 40 years. Mrs. M.'s parents were healthy. Her mother lived till the age of 86 years, and her father till the age of 74 years. The remainder of the latter's children, five in number, are all living and healthy with no history of tuberculosis in the family. Mrs. M. is the only case who came under my observation, in which the disease may be said to have been acquired.

Of the fourteen cases of widows or widowers who came under my observation, whose husbands or wives died of phthisis pulmonalis, all are now living and healthy.

Now the most popular theory as to the cause of phthisis pulmonalis seems to be that the cell must have a deficiency in vital energy and also must have inherited the tubercular diathesis before the tubercle bacillus can attack it with any degree of success. Then it follows that a cell may have a deficiency in vital energy, not having inherited the tubercular diathesis, yet the tubercle bacillus may not attack it, or the cell may have a sufficiency of vital energy and an inherited tubercular diathesis and it may be invulnerable against the tubercle bacillus' attacks. But I have no doubt, in the case of Mrs. M., that the tubercle bacillus successfully attacked the cell with deficient vital energy, without it having had inherited a tubercular diathesis.

On the other hand, there are those who may claim the old theory that the disease is transmitted from parent to offspring, and they might ask the following pertinent questions: If there was no tubercle bacillus, would there be any tuberculosis, and if there was no inherited tubercular diathesis would there be any consumption? It is a well-known fact that "the tree is known by its fruit." You all remember how Jacob of old, transformed the sheep and cattle into "ring-streaked, speckled and spotted," in order to take advantage of his father-in-law, Laban. With our present knowledge of the laws governing procreation we would say that such a change in the color of those animals was caused by the impression made on the parents at the time of copulation. That is, the change in the color of the animals depended on the impression on the ova, either through the female or through the fructifying influence of the male—the semen—at the

time copulation took place. Consequently, there could not have been any other factor to cause the change in the color of the animals. When a person of tubercular diathesis is impregnated, or impregnates, there may be claimed to be a constitutional condition in the offspring similar to that of the affected parent. This constitutional condition may affect the circulatory, nervous or glandular system in such a way as to interfere with a healthy physiologic action of the same and, thereby, producing a weakness in these various systems, and finally diseased cells and consumption are developed.

I believe that the proper plan to prevent this disease, tuberculosis, is to educate young men and young women how they should make their selections, in order that their posterity may not be tainted by disease, and they should also be taught the great importance of making healthy selections.

Fifth Avenue and Ross Street.

SOCIETY PROCEEDINGS.

Chicago Ophthalmological and Otological Society.

Regular meeting, held Oct. 8, 1895, at 155 Dearborn Street. There were seventeen members in attendance, DR. GRADLE in the chair.

Minutes of the last meeting were read and approved.

DR. COLEMAN reported a case of ulcer of the cornea treated by a Saemisch incision and subconjunctival injections of corrosive sublimate. De Wecker says in the *Annales de Oculistique* for June, 1895: "I return to large injections of sublimate, 1 to 2000 with eserine. I have ordinarily injected half a Pravaz syringe full even in very advanced cases of suppuration with hypopion. Only six or eight injections are necessary to obtain recovery, with clearing up of the diseased portion." Abadie, in the *Annales de Oculistique*, 1895, writes: "If the patient comes to us with very extensive suppuration of the cornea, and it is not in our power to restore the diseased parts, we are always able to preserve the portion which still remains intact. To accomplish this object several methods are at our disposal—the method of Saemisch, the actual cautery. What I wish to say is that to-day we have a much better method, that is subconjunctival injections of sublimate, and we can in a large number of cases dispense with the actual cautery. The following cases prove a surprising success: one, a man, aged 48, with an ulcerated point on the cornea and a little pus in the anterior chamber. Injection of two drops of sublimate, 1 to 1000. This was repeated on the third day. Patient was cured without the slightest alteration in the cornea.

Second case, aged 40, with extensive ulcer and large hypopion. Had three injections; quite cured in seventeen days.

The *Annals of Ophthalmology and Otology*, April, 1895, contains an abstract of Dr. Bull's paper on the question of the "Efficacy of Subconjunctival Injections of Mercuric Bichlorid in Ophthalmic Therapeutics." A review of the literature showed great diversity of opinion among the various observers as to the value of this new treatment during the past year. Dr. Bull had employed these injections in forty-eight cases. He summarized as follows: "Pain is always severe in spite of cocaine, and reaction is apt to be very severe. The only class of cases in which the sublimate injections seem to exert any positive effect in allaying symptoms and shortening the duration were those of scleritis and irido-choroiditis of the non-syphilitic type. The treatment