infections. In the isolation wards of the hospital in Providence 37 cases have been admitted with no cross infection among nearly a thousand other patients admitted. At one time 3 cases developed in the diphtheria ward where they had been cared for by a nurse who was coming down with the disease.

TUBERCULOSIS

As tuberculosis has its chief, and apparently primary, seat in the lungs, it was only natural to assume that the route of infection is by the inspired air. Koch's studies on the resistance of the bacillus and its survival in dust greatly encouraged this view, as did the extensive observations of Cornet as to the presence of the germs in dust obtained in the neighborhood of careless consumptives. Then came Flügge, who attempted to show that dust is of little moment, but that infection by floating droplets is the chief mode of infection. The somewhat active controversy between the followers of these two men has, I think, merely served to convince most of us that under the conditions of the experiments both dust and droplets are capable of causing the disease. In tuberculosis, the pathologists take a prominent part in the discussion of causation, but are at variance as to where the primary lesion is to be found and whether the bacillus enters by the respiratory or alimentary tract. Many think that most tuberculous infection dates back to early life, and not a few derive it from the ingestion of milk, though this latter view seems to be losing ground. The weight of evidence derived from these lines of study seems to indicate that the infection is commonly, if not usually, air-borne, and that the primary lesion is in the respiratory tract. There are not a few, however, who hold that contact infection plays a prominent, if not the larger part, in the distribution of the disease and that bacilli-holding sputum carried to the mouth and nose in countless ways reaches the lungs by insufflation or by absorption through the tonsils or through lower sections of the alimentary tract.

Tuberculosis is such a chronic disease that it has thus far been impossible to devise observations on human beings which would throw much light on the route of infection. Recourse must be had to animal experimentation, and most of this has been done under conditions far from natural. Still, a good many experiments have consisted in exposing test animals in cages in apartments occupied by consumptive animals or human beings. A large number of tests made by the Department of Agriculture by exposure to animal sources of infection resulted in a very few cases of tuberculosis. Exposure of test animals in cages in rooms with human beings in Germany, France and this country has almost invariably resulted in the development of the disease in some of the animals, but unfortunately in no instance could contact infection be absolutely excluded, except in an experiment made in a house in Providence occupied by a careless consumptive. Here two sets of guinea-pigs were exposed in cages, one set fed by the patient and the other excluded from any possible form of contact. Most of the animals in both sets developed the disease. The animals in the locked cage covered with wire gauze were, I believe, infected by mouth spray, as the patient often held his face right in front of the box and talked to the animals. There is then a good deal of evidence to show that tuberculosis may be, oftentimes, an air-borne disease. This evidence is certainly far more convincing than for any other disease, except perhaps anthrax. If, as seems probable,

the air may serve as a vehicle for the carriage of tubercle bacilli it is probably because, in this disease, more germs are discharged indoors in places where they are likely to get into the air than is the case in other diseases, and because the bacilli have a fairly high resisting power.

ANTHRAX

On theoretical grounds, anthrax ought to be more easily air-borne than any other disease. The spores resist drying for many years, their virulence is high and they are, at times, found in large numbers in materials handled by workmen in such a way as gives rise to much dust. Workers in wool and hair, coming from anthrax-infected regions, have always been subject to this disease. Formerly, when no precaution was taken to prevent the breathing of dust, the disease was very much more prevalent than it is now. During nine months, in 1882, thirty-two cases occurred in the woolen industry in Bradford, England, of which twenty-three were of the pneumonic type and presumably due to the inhalation of dust. Last year there were seventeen cases, and of these only three were of the internal type, so that it appears that with increased freedom from dust the latter type of the disease has shown marked decrease while the external type has not. Careful observers in England and Switzerland have noted the occurrence of anthrax in animals grazing on pastures on which dust from wool-sorting or hair-cleansing rooms was blown.

After this review, a summary scarcely seems necessary, or even a formal conclusion. We have seen that a number of important diseases, formerly considered exclusively air-borne, have been shown never to be such. There is little evidence that, among the diseases which commonly occupy our attention in this part of the world, aerial transmission is a factor of importance. In most it is under ordinary conditions of home and hospital, a negligible factor. For tuberculosis alone is there evidence that air-borne infection is a factor of moment, but the last word has not been said as to the etiology of this disease. We may be sure that the sewer-gas bogey is laid, the notion that dust is a dangerous vehicle of every-day infection is unsupported and that mouth spray is usually effective only at short distances.

HYPERACIDITY OF THE STOMACH

ADOLF SCHMIDT, M.D., GEH. MED.-RAT.

Professor of Internal Medicine, Vereinigte Friedrichs-Universität,
Halle-Wittenberg
HALLE, GERMANY

There is hardly a single condition concerning which opinion and treatment have so much changed during recent years as so-called hyperacidity of the stomach. Practically this term is taken to mean those cases of increasing gastric disturbances which appear at various intervals after the taking of ordinary food or special kinds of nourishment. The pain is sometimes accompanied by eructations of gas and small quantities of soured fluid (heart-burn). On examination of the gastric contents obtained with the stomach-tube, there is found either hyperacidity, large quantities of normal stomach-juice, or acid gastric juice. Thus we have either hyperacidity or hypersecretion. The latter may be separated into a digestive and a continuous type.

The continuous type of hypersecretion, when appearing periodically, has been called by some authors

gastroxynsis. The most serious cases of continued hypersecretion are known as Reichman's disease or gastrosuccorhea.

When further analysis of the cases for various abnormal stomach contents is attempted, many difficulties arise from the fact that the contents do not represent a simple mixture of the test-meal and the gastric secretion. In fact, they first undergo certain uncontrollable changes by their evacuation into the duodenum, which occurs almost immediately after eating. Furthermore, hydrochloric acid may be partly neutralized by the alkaline fluid of the duodenum and the mucus produced by the superficial gastric epithelium. Neither the most exact examination of the stomach contents, performed with repeated aspiration during digestion, nor the valuation of the residue by the Matthew-Raymond method is able to give us a true idea of the different factors here active.

It is important to know if, in the disease spoken of, the amount of secreted hydrochloric acid is the same as in normal conditions.

In consequence of investigations made on the lower animals by Pawlow and on men by Rubow, Hornborg, Bickel and others, the opinion has been almost generally accepted that pure gastric juice has always the same rather high percentage of hydrochloric acid, even under pathologic conditions. According to this opinion we are led to believe that under normal circumstances the pure glandular product is diluted or partly neutralized by alkaline or gastric mucus, since we do not meet such high degrees even in the empty stomach. According to Forschbach and Schutz, we ought further to believe that the different kinds of hyperacidity or hypersecretion arise either from the secretion of stomach juice being too copious or lasting too long compared with the digestive stimulus represented by the test-meal, or by irregularity of the intervals occurring between the gastric secretion and the emptying into the duodenum. Recently Gregerson, by careful investigations, has proved that at least so far as the human stomach is concerned, Pawlow's opinion must be abandoned, unless we presume that the superficial epithelium produces concentrated alkaline fluid, together with pure gastric juice, a suggestion which is rather paradoxical. Comparing the stomach contents after the usual test breakfast (tea and toast) with those stated after the so-called dry testmeal, Gregerson found, shortly afterward, the same amount of fluid and nearly the same percentage of acidity in both cases. Hence we must conclude that the stomach secretion varies considerably in the amount of hydrochloric acid which it contains.

Under pathologic conditions, these variations occur with greater intensity in view of the fact that the gastric secretion, as well as the motility of the stomach, is dependent on the nervous influences which arise not only from the stomach contents, but also from the brain. It is not remarkable that the gastric contents, as well as the other symptoms of the disease, should vary with the state of the central nervous system. Patients themselves will talk of the whims of their stomachs. One suffers regularly from eating sweets, another from taking too much fat and a third stands the same food either badly or well, according to his mood. Nearly all of them agree that sorrow, excitement or other psychic influences increase their complaints.

One of the most interesting questions connected with this disease is whether or not there is an etiologic relationship between the hyperacidity of the stomach con-

tents and the so-called acid pains, gnawing pressure in the stomach some hours after meals and the heart-burn. The common experience of instant cessation of these symptoms after a dose of sodium bicarbonate or other alkaline salts would seem to confirm it. But it is a well-known fact that hyperacidity and hypersecretion will continue even after the symptoms are completely relieved. Moreover, patients with diminished gastric secretion, achylia or hyperacidity complain sometimes of quite similar pains, so that no experienced observer will venture to take the hyperacidity for granted on the basis of anamnesis without confirming the condition by the use of the stomach-tube. As a matter of fact, the question has been decided in the negative by the results of a simple experiment. It has been found that large quantities of highly concentrated hydrochloric acid introduced into a normal stomach (Schmidt took solutions of 5 per cent.) do not produce such pains. The so-called acid pains, therefore, must be due to some other cause, either a spasm of the stomach sphincters or a hypersensibility of the mucous membrane, or hydrochloric acid, or something hitherto unknown. Cinematographic roentgenoscopy of the stomach during digestion, which has reached its highest perfection in the United States, has shown that very often the beginning of the so-called acid pains takes place at the same time as the spastic contraction of the pylorus.

Of more practical importance, but associated with the theoretical points just discussed, is the question whether the state of hypersecretion or hyperacidity occurs as a disease, *per se*, dependent only on nervous cases, or is always based on certain organic lesions such as gastritis, ptosis of the stomach, ulcers, adhesions, scars, gallstones, etc. A short time ago we did not hesitate to take the majority of the cases as pure stomach neuroses, provided that the presence of a clear sign, as hemorrhage or periodic stenosis of the pylorus, was not against it. To-day our opinion has changed chiefly in consequence of the rediscovery of duodenal ulcer and the description of its clinical signs by American and English authors. I shall neither enter into the difficult question of the exact diagnosis of duodenal ulcer, nor discuss the causes of its high frequency in America compared with the small number of cases occurring in Europe. Yet I shall attach weight to the merit of the Mayo brothers and other surgeons who have shown that an ulcer of the duodenal or pyloric region is often hidden by the symptoms of hyperacidity. We always have reason to suspect the presence of such an ulcer when roentgenoscopy reveals that peristalsis, initially increased, stops in the later periods of digestion; that the well known six hours' residue remains, and that these two periods are separated by a spasmodic contraction of the pylorus, which corresponds to the setting in of hunger-pains. Of course, roentgenoscopy gives more convincing signs, among which may be mentioned the deviation of the pylorus to the right, the worm-eaten cap and the constant filling of the duodenal bulb, but they are seldom to be found in cases of mere hyperacidity running along without vomiting, loss of blood and other decisive symptoms of ulcer. Here I wish to emphasize the importance of a most careful examination by all methods available in each case of hyperacidity or hypersecretion in which there could possibly exist a hidden ulcer of the stomach or the duodenum, or scars, adhesions, etc., caused by them.

If the results are negative we must go on searching for other causes of hyperacidity. Certain rare cases,

like beginning cancer of the pylorus, the occurrence of gall-stones and simple catarrhs of the stomach must first be considered. We can take it as a fact that in many liver diseases, especially those accompanied by pains, the secretion of the gastric acid is increased as the result of a reflex influence. There can also be no doubt that some acute, as well as chronic, inflammations of the stomach result in increased acidity, whether they are connected with atonic ptosis or not. These acid catarrhs may cause some difficulty for diagnosis, since the mucous product of the epithelium, the characteristic sign of catarrhal inflammation, is dissolved by the gastric juice, and therefore does not reappear in the washed-out specimen. The patients are afflicted often with an abnormal irritability of the stomach, so that even a little too much free acid leads to a spasm of one or both sphincters. When the stomach is washed, these patients choke continually and vomit violently. Here we encounter the hyperirritability of the mucous membrane already mentioned, which should be considered as a local manifestation of the general nervousness with which most of these patients are suffering. The irritability of the autonomous nervous system in the so-called vagotonic patients sometimes leads to spastic contraction of the stomach vessels, and this phenomenon is regarded by some authors (Bergmann, Roessle), as the very cause of the ulcer.

This shows that the present state of science does not allow a large space for the purely nervous cases of hyperacidity. It is due to the increasing refinement of our diagnostic knowledge that here, as well as in other branches of medicine, the number of those doubtful cases is more restricted, and the opinion is daily spreading that the nervous moment is only an accompanying symptom of an organic lesion, though the nervous feature may completely overshadow the real disease. Yet we should be premature in entirely denying to-day the clinical unit of hyperacidity. We confess, with resignation, that in practical medicine we often must be satisfied in stating the changed chemism of the stomach.

In treatment, the principal point is not to restrict ourselves to the stomach alone, but at once to aim at calming and strengthening the nervous system, the state of which is of so great importance for the comfort of the patient. Some patients suffering with hyperacidity can get rid of their disease in no other way than by giving up their work for some time, leaving their families and undergoing a real confining treatment in a hospital or sanatorium. I demand this in every severe case, and believe that I owe many good results to this procedure.

Naturally we try first to reduce the increased secretion of the stomach by application of medicines. Atropin is known as a substance which diminishes the secretion of the gastric glands as well as those of the salivary and sweat glands. Therefore the extract of belladonna, which at once levels the tonus of the vagus and removes the accompanying spasm of the sphincters, is a popular part of many stomach powders. Von Tabora recommended regular injections of 0.5 or 1 mg. of atropin before meals, but it is surely not advisable to continue this for a long time, if there is a chance of removing the real cause of the illness in another way. The alkalies which have already been taken for a time by most patients not only neutralize the superfluous acid, but, according to Bickel, also stop their production to a certain degree. On the other hand, they irritate the stomach glands, particularly if carbonic acid is set

free by their reaction with the stomach acid, for the carbonic acid is not indifferent to the mucous membrane. Therefore, we prefer to give the magnesium oxid instead of sodium bicarbonate, and we combine it with small quantities of belladonna and a purgative salt, such as sodium sulphate. With the latter I intend to produce a transudation of fluid into the intestine and so to diminish the secretion of the stomach glands. The effect of the Carlsbad water, which often proves beneficial, seems, indeed, to result from the favorable combination of alkaline and purgative salts, but it should always be given in small quantities. Some other acid-binding drugs, as powdered charcoal, kaolin and aluminum silicate, act in another way, by mechanical absorption, each of them being worth trying occasionally. More heroic is the cure of Henriques and Hambart, who for a time withdraw all chlorids from the food. Winternitz and Petri of Halle followed a new principle by introducing magnesium peroxid into the stomach. These authors observed that small quantities of hydrogen peroxid diminished even in a pronounced degree the acidity of the stomach contents, and that this fact is based on an irritation of the surface epithelium, the alkaline product of which neutralizes the present acid. Here we have to deal with the opposing effect of one irritation to another. Lastly, silver nitrate belongs to this group. Its excellent effect on hyperacidity is caused partly by its astringent qualities on the secretory cells, and partly by blunting the hyperirritability of the mucous membrane. For the latter purpose it is to be given as a medicine to be taken half an hour before meals; for the former it is used as a solution to rinse the empty stomach in the morning by means of the tube.

As regards the diet, which is perhaps even more important than the treatment by medicine, opinions differ, though some controversial points are already cleared by the results of experimental medicine. There is no doubt that all kinds of foods and drinks which belong to the class of secretory stimulants must be avoided, as, for example, too much salt, sparkling drinks, high-percentage alcohols, coffee, meat-juices or extracts of beef, and spices. More difficult is the question of how to arrange the meals. We distinguish three groups of diet specialists corresponding to the single groups of food; those who prefer albuminates, those who prefer carbohydrates and those who prefer fat. The advocates of the first group emphasize the fact that only the albuminates can bind the hydrochloric acid, and therefore have the same effect as the alkalines. They sometimes go so far as to prescribe that when the pains appear the patients should take regularly some food prepared from meat, eggs, etc., lest the stomach should become empty.

Contrary to this, the advocates of the carbohydrate diet assert that they are more correct, since the ingestion of carbohydrates stimulates the production of gastric acid little or not at all, while albuminates first stimulate and then neutralize the acid. They declare their diet as a rational one superior to the symptomatic albuminate diet. They not only avoid stimulating the production of the acid, but also try to produce a reflux of alkaline juice from the duodenum into the stomach, which is most easily done by taking one or two tablespoonfuls of pure olive-oil before meals.

With regard to these three groups, it must be stated, from a practical point of view, that it is most difficult to give one special kind of food for any length of time. Moreover, it is really unnecessary because experience

shows that some other dietary prescriptions are more important. I long ago returned to a mixed food and am quite satisfied with it, but strictly observe the following rules, which I believe to be decisive:

1. All food must be well cooked and most carefully minced. This order is based on my own theory that the chief business of the stomach is to reduce the food into small pieces, not by its muscular force, but by chemical influences. These influences consist in the dissolution of the intermediate substances, the connective tissue of meat and fat, the gluten of the bread and the so-called "middle lamina" of the vegetables. As to the connective tissue, I have shown that only the gastric juice is able to digest raw or insufficiently cooked connective tissue, and that during the digestion of meat in the stomach it is always connective tissue which is first dissolved, so that the muscular fibers are set free. This phenomenon is the base of the connective-tissue proof which is a part of my test diet.

As to the disintegration of bread in an efficacious gastric juice and its remaining compact in achylia or hyperacidity, it is sufficient to mention the phenomenon which is generally known as amylorrhexia. The middle lamina of the vegetables is not really dissolved by the stomach juice, but is only prepared by it for dissolution in the alkaline intestinal juice. In this way the chemical stomach work leads to the reduction of all sorts of food into small pieces, and this is a much more significant act of the organ than solution of albuminous matter, as the latter is more surely and with more power effected by pancreatic juice. As the food pieces are broken up by the chemical influence of the gastric juice, they are pushed into the duodenum. Pieces not quite broken up remain for a long time in the stomach, while fluids are rapidly pushed forward. This process is also of importance for the evacuation of the stomach.

2. The stomach must come to rest at least once during the twenty-four hours. This can best be arranged by changing the usual time of meals. The patients should eat large quantities of food in the morning, but only a small quantity in the evening. This means that during the forenoon, when, according to my experience, the patients suffer the least from their pains, they should take two or three meals with intervals of two hours so that there is little appetite left for lunch. After lunch they do not take anything until 7 p. m., when only porridge is given. They will now fast until the next morning, and this change also has the advantage that the production of gastric acid, which has been adapted before to lunch and dinner, is regulated in a new manner; the chief secretory reflexes, easily provoked during the former style of living, are now put out of circuit. I can say certainly that I have obtained very good results with this method.

3. Drinking should be generally diminished and restricted to those hours when the stomach is not filled with food. I therefore give the meals rather dry, except the first breakfast and the evening soup, and let the patients drink a short time before lunch or in the afternoon. This prevents loading the stomach, which measure is of special importance to patients suffering from atonic ptosis.

4. If the condition is a severer one or if it is combined with ptosis of the stomach, the patients have to stay in bed for two or three weeks. I think this advice most important. It results from the experience that many patients have their acid pains only when being about or walking, never when lying down. It is further based on

the frequency of latent ulcers which lurk behind the symptoms of hyperacidity.

It may be remarked that my dietetic treatment of hyperacidity comes into close touch with the old Leube cure in gastric ulcers. I confess that there is only a certain difference of degree.

I sometimes also order hot compresses in simple hyperacidity, if they give relief to the patient, but I do not apply them continually as Leube does—only twice daily for two hours. For the night I substitute cold hydropathic compresses for them.

Washing the stomach is indicated only if the hyperacidity rises on the basis of a catarrh. Then I rinse the empty stomach in the morning with 1:1,000 solution of salicylic acid or 1:5,000 solution of silver nitrate. Moving the bowels must be carefully regulated.

What is to be done if all these prescriptions do not lead to recovery? This question is equivalent to the other—"Under what circumstances are we allowed to operate in hyperacidity or hypersecretion?"—and cannot be answered generally. Indeed, the nature of the process alone, that is, its being a functional one, cannot always be decisive; it may sometimes be of less importance than the violence of the pains, or the general state of health or the social position of the patient. There is a difference between an apparently neurasthenic patient whose condition remains unchanged after a well-performed gastro-enterostomy which only reduced his stomach pains—for example, an old man who has given up business and can stand his pains if he takes care with his food—and a young man overwhelmed with his business who wishes at all costs to get rid of his sufferings. In the latter case I should advise an operation. There is no doubt, in my opinion, that a case of hyperacidity or hypersecretion, even without complication, which cannot be relieved by internal treatment in order to make the patient fit for work, gives an indication for operation. The manner and procedure of operation cannot be discussed here in detail. This depends on the state of the patient's condition found during operation. If an ulcer of the pylorus or duodenum or a scar is the cause of the hyperacidity, gastro-enterostomy is the right procedure, and if possible, the pylorus should be closed. Perigastric adhesions must be loosened, and gall-stones removed by cholecystotomy or cholecystectomy. But what is to be done if there is no anatomic lesion at all? In these cases I should advise gastro-enterostomy, but without the closing of the pylorus. The latter can be recommended only if violent pylorospasm is present. The patient should never be allowed to get up and eat anything he wants immediately after recovering from the operation; he must go for some time with the diet and drugs in the same manner as before operation, as the operation does not restore health alone, but only establishes conditions favorable for internal treatment.

Fee-Splitting.—The secret division of fees between two practitioners is indefensible in theory and reprehensible in practice. Viewed from any and every angle, there is no argument that will validate the physician's attitude of auctioneer—selling out his patient to the highest bidder. Nor can the purchaser, by any specious argument, justify his part in so foul a transaction. The practice is rampant not only among those of smaller incomes, but with those enjoying large means as well. Poverty does not justify crime any more than low wages immorality; and certainly, cupidity lays a poor foundation for fair dealing, but builds well for moral strabismus.—Arthur J. Patek, M.D., *Wisconsin Med. Jour.*