right hemisphere again yielded the same organism isolated at the two previous spinal punctures.

BACTERIOLOGY OF THE CASE.

As has been already stated, the same organism was isolated from the spinal fluid at the two spinal punctures and again at the postmortem, as also from the throat, and a description of one will serve for all four strains.

Morphology.—The organism is a short bacillus measuring 2.5x0.5 microns. It stains readily by all anilin dyes, also retaining Gram's. Preparations from 48-hour cultures show a beaded appearance of the organisms when stained with Loeffler's methylene blue. There were also marked involution forms, such as club-shaped and dumb-bell-shaped organisms of variable size.

Culture Characteristics.—The organisms grow well on all the ordinary culture media, but best on blood serum, where at the end of 10 hours a creamy slightly elevated layer is formed. On blood-agar plates we find at the end of 24 hours pinhead colonies without hemolysis, though when glucose blood-agar was used often a zone of greenish tinge developed about the colonies.

Animal Experiments.—Guinea-pigs were inoculated subcutaneously with 1/20 of a 24-hour blood serum culture suspended in normal salt solution. The animals appeared sick at the end of 12 hours, with beginning evidence of local infiltration. Death occurred in from 48 to 56 hours. At autopsy there was found locally marked edema and bloody exudate into the subcutaneous tissues, extending some distance beyond the point of inoculation. The internal organs, as a rule, showed some pallor, while the adrenals were usually large and deep red in color. Cultures from the heart's blood and internal organs remained sterile, while the organism was always recovered in pure culture from the local process.

Unfortunately we were forced to discontinue the work for two months, and on resuming it the organisms had lost their virulence so that it was impossible to test the protective influence of diphtheria antitoxin.

To differentiate the bacillus isolated from the *B. xyro*s and *B. pseudodiphtheria*, the method of Knopf¹ was adopted. Serum sugar media were prepared with inulin, dextrose, saccharose and dextrin and inoculated with several strains of *B. diphtheria* and also the bacillus C, described above, with the results as shown in the table.

FERMENTATION OF SERUM SUGAR MEDIA BY *B. DIPHThERIAE*.

Organism.	Serum Sugar Media.			
	Inu- lin.	Dextrose.	Sac- charose.	Dextrin.
P ₁	—	Fermented.	—	Fermented.
P ₂	—	Fermented.	—	Fermented.
R	—	Fermented.	—	Fermented.
Bacillus C isolated from case	—	Fermented.	—	Fermented.

We see from the table that Bacillus C corresponds in its ability to ferment sugar to the other known *B. diphtheria*.

CONCLUSIONS.

Bacillus C isolated from the spinal fluid in the case of meningitis thus produces lesions in guinea-pigs typical for the *B. diphtheria*. It agrees, moreover, with the same morphologically and culturally, as also in its ability to ferment certain sugars, and we are, therefore, justified in classing it with this group. The case just

reported also shows that these organisms occur more frequently in the central nervous system than suspected, and on careful search could undoubtedly be recovered in a large number of cases.

ETIOLOGY OF CEREBRAL PALSIES IN CHILDREN.*

WILLIAM J. BUTLER, M.D.
CHICAGO.

A number of classifications have been suggested for the cerebral palsies of children; some based on their pathology, others on the clinical manifestations and one on the time of their occurrence. The latter, suggested by Sachs, is very convenient. He divides them into antenatal, natal and postnatal cases.

It is not always easy to determine the etiology of cerebral palsies in individual cases. This is due chiefly to their coming late under observation or because the autopsy findings, obtained sometimes years after the onset of the disease, may not reveal the nature of the initial lesion and its cause. This applies more particularly to prenatal palsies and those occurring after birth.

The cerebral lesions having their origin *in utero* seem to depend partly on conditions affecting the mother and secondarily the fetus and partly on the development of the fetus. The brain of the fetus may suffer lack of or defective development as may other fetal organs. The pathologic change commonly found in prenatal cases is porencephalus, a localized brain atrophy, which is an end result of a primary vascular lesion, as embolism or thrombosis, which, in turn, has arisen from infections in the mother and fetus, as syphilis, etc. Injury to the abdomen of the pregnant woman has resulted in isolated instances in cerebral hemorrhage in the fetus and spastic contractures have been found at birth (Cotard, Gibbs).

Birth palsies are dependent on intracranial hemorrhage, the result either of trauma to the infant's head during precipitate labor, prolonged labor, forceps delivery or the result of prolonged asphyxia at birth. Occasionally a history of breech presentation is noted in a birth palsy. The hemorrhage is meningeal in location and arises from laceration of the pia vessels usually close to the longitudinal sinus, from which point it may spread and envelop a considerable part of one or both sides of the brain. While this is the invariable finding in recently posted cases, if they survive a number of years, as is common, secondary changes follow which may render the primary lesion unrecognizable and, instead, we may have a lobar sclerosis, etc. This applies, however, not only to natal cases, but to all spastic cerebral paralyses in children who survive the first insult, and it is this point which has caused so much discussion as to the etiology of postnatal cases. As the prognosis of those cases happening in childhood is usually favorable as to life, the autopsy findings of acute cases are correspondingly infrequent. There is little doubt of the prominence of the rôle of vascular lesions as a cause of many of these cases of hemiplegia, in fact a predominating factor in their production, as in early posted cases, hemorrhage, embolism and thrombosis are frequently found.

The paucity of pathologic information, however, has

* Read in the Section on Diseases of Children of the American Medical Association at the Fifty-seventh Annual Session, June, 1906.

1. Jour. Med. Res., 1904, xii, pp. 475 to 479.

developed considerable conjecture, and as a result various views have been advanced in their explanation. Gowers believes many of them due to thrombosis of the veins of the pia and the longitudinal sinus. Sachs is convinced that the majority of them are caused by cerebral hemorrhage. Strümpell, on the other hand, thinks that the acute spastic hemiplegias of children are the result of an encephalitis, dependent on an acute infection. This latter view was based entirely on clinical study originally and only recently received postmortem confirmation in a few reported cases.

While the only absolute proof of the exact nature of an organic disease usually lies in the pathologic findings, however typical the clinical picture may be, we are, nevertheless, often forced to depend on a study of the latter, and if this is permissible clinical observation of acute hemiplegias and occasionally a chronic one would seem to admit of a reasonably accurate pathologic diagnosis in many instances.

The cause of spastic hemiplegia in children may be briefly stated as follows: First, vascular lesions, to-wit: (a) hemorrhage from venous or arterial rupture, the result of trauma or occurring spontaneously in intense congestions, as in convulsions or the paroxysm of whooping cough; (b) embolism from cardiac or arterial thrombi or endocarditis; (c) endarteritis and thrombosis. Second, inflammatory changes, to-wit: acute encephalitis and meningoencephalitis. Third, rarely, a tumor.

CASE SUMMARIES.

The following cases would seem to illustrate some of the above causes:

CASE 1.—*Diagnosis.*—Intracranial hemorrhage, either delayed traumatic or vascular rupture occurring in the course of a convulsion which in turn was the result of fright.

History.—James B., aged 4 years, fell from a baby carriage to the ground. No injury was evident at the time, but he was greatly frightened. On awakening the following morning, he sat up and called for a drink of water. A moment later he was found in a convulsion, followed by coma. On awakening he had right hemiplegia and aphasia; no fever. Examination a week later gave the condition as above, no trace of injury, no heart lesion, urine negative, electrical reaction unchanged.

CASE 2.—*Diagnosis.*—Left hemiplegia, caused by an embolism from a heart thrombus, forming in the course of acute myocarditis in diphtheria.

History.—Martha O., aged 10 years, had diphtheria; antitoxin was given on the third day. In the second week there was palate paralysis, marked dyspnea, palpitation, edema of the extremities and scanty urine. A few days later while in this condition she was given a tub bath, and during the ordeal she became very cyanotic and seemed unconscious. Ten days later left hemiplegia was noticed. In the course of several weeks all symptoms subsided, but paralysis persisted. Examination at the third month revealed spastic paralysis of the left side, heart area enlarged, second pulmonic loud, no murmurs, urine normal.

CASE 3.—*Diagnosis.*—Juvenile progressive dementia, resulting from syphilitic disease of the brain in the form of an endarteritis and thrombosis with consequent cerebral changes.

History.—James O'B., aged 10 years, at the age of six weeks had a generalized eruption, also suppuration around the nails. All cleared up under treatment in a few weeks. He developed normally to the sixth year; thereafter he showed failing intelligence, spastic gait and stammering speech. Examination revealed rigid pupils, right spastic hemiparesis with exaggerated deep reflexes. No ataxia. Eye grounds normal; no change in electrical reactions, no disturbance of rectal or vesical function. *Petit mal* attacks.

CASE 4.—*Diagnosis.*—Acute encephalitis.

History.—Bennie H., 14 months old, always well up to the present illness, was seized with vomiting, fever and drowsiness.

Inside of 48 hours left-sided paralysis was noticed. Examination a week later showed left spastic hemiplegia, deep reflexes exaggerated, electric reactions normal, no evident mental impairment. Subsequent observation showed continuous improvement. He started to walk a couple of months later and shows but slight traces of hemiplegia. The child is playful and has no apparent discomfort.

The similarity of the clinical history of this case to that of a poliomyelitis is evident. Attention was called to this similarity by Strümpell in his original article on the acute encephalitis of children. In this he also suggested that they were caused by the same infection, the difference in the clinical picture depending only on the location of the process; in the anterior horn in poliomyelitis and in the cerebrum in the polioencephalitis. This view was reinforced by the report of the occurrence of a case of poliomyelitis in one child and encephalitis in another in the same family at about the same time.

An encephalitis may also occur during the course of or following an acute infectious disease, in which event it is assumed it is caused by the infectious agent or its toxin. This has been reported in many of the acute infectious diseases, as measles, scarlet fever, etc., but notably in influenza, in which Nauwerck and others have demonstrated the cerebral lesion to be due to a local influenza infection by finding Pfeiffer's bacilli in the diseased foci.

Osler and Dercum have reported cases of hemiplegia dependent on cerebral tumors. Seeligmüller laid some stress on solitary tubercles as a cause of the spastic paralyses, but apparently without any convincing effect.

The following case is presented as one in which a spastic hemiplegia was a prominent symptom. I am indebted to Dr. J. A. Harvey for the privilege of reporting this case.

CASE 5.—*Diagnosis.*—Anatomic: solitary caseous tubercles of the brain—(a) right lobe of the cerebellum, (b) left crus and internal capsule, (c) left occipital cortex. Tubercular meningitis. Chronic internal hydrocephalus.

History.—Marie P., aged 4 years 9 months, was in fair health to the third year. She had a series of infections, including measles, pertussis and pneumonia. Thereafter she was less active mentally and physically. Occasionally she had convulsive seizures, dizziness, headache, cerebellar ataxia, etc. At the age of 4 she had an attack characterized by fever, vomiting and opisthotonos. This subsided in five days, leaving left ptosis and facial paresis. Several days later right spastic hemiplegia developed. Double optic neuritis. On spinal puncture a clear serous fluid under pressure was obtained. Bacteriologically negative. Temperature, 99 to 101 degrees *per rectum*. Ultimately she succumbed in coma.

On reviewing the etiology of infantile spastic hemiplegia of postnatal origin, the conclusion seems unavoidable that the causes are essentially the same as produce hemiplegia at later periods of life.

While we do not find in children the miliary aneurisms or arteriosclerosis that are found in later life, and which with high arterial tension is the source of spontaneous hemorrhage at this period, we may have a hemorrhage resulting from spontaneous rupture during sudden, intense, venous congestions, as in convulsions or the paroxysms of whooping cough, or we may have a hemorrhage from head injuries. Likewise an infantile spastic hemiplegia may depend on a specific cerebral endarteritis and thrombosis, or an embolus from a nearby arterial or cardiac thrombus or an endocarditis, resulting in areas of cerebral softening, all of which are common causes of hemiplegia at all periods of life.

Although the frequency of inflammatory processes

as a cause of infantile spastic hemiplegias may remain a disputed question, the fact of their occurrence seems to have been placed beyond a doubt, in view of the pathologic reports of Ganghofner, Nauwerk, Raymond, Batten, etc. Encephalitis, however, is a lesion that may occur at any time of life, and is probably an important factor in the causes of cerebral paralyses in children. Close clinical observation of some cases of infantile spastic palsies seems to allow of no other assumption than that of an acute encephalitis as its cause.

Cerebral tumor is hardly to be considered as a factor in these cases, because hemiplegia, if it exists, will merely form one of a series of clinical manifestations of the tumor, the diagnosis of which will take precedence over a symptom.

BACTERIOLOGIC AND BLOOD FINDINGS IN THIRTY-SIX CASES OF BOWEL INFECTION OF INFANTS.*

J. C. COOK, M.D.
CHICAGO.

The 36 cases which I am about to present have been selected from 350 cases of diseases of children, which came under my supervision in the Jackson Park Sanitarium during the summer of 1905.

Those cases, the clinical aspect of which showed more or less severe type of anemia, are the ones selected for the special work to which I wish to call your attention. The work was taken up with the hope of obtaining findings pointing to the relation of the condition of the blood, the enlargement of the glands and spleen and the bacterial and metabolic intoxications.

The most important data obtained in each case are incorporated in the table. The cases have been arranged with regard to the red blood count, the highest to the lowest, that being the most logical order on which to make comparisons.

In every case of examination of the blood, including a count of the red and white cells, a differential count of the white cells and a hemoglobin test was made. In cases in which a general infection was suspected, cultures from the blood were made. The results of microscopic examinations from the stools and throat were recorded, as they seemed to have direct bearing on the clinical aspect of the case.

Of the physical findings, only the enlarged glands and spleen and lesions which might serve as a clue of infection were recorded. Such other information as is of interest in the consideration of the subject is also placed in the table.

The diagnosis which is recorded in each case was that made at the time the case was sent into the hospital, and, in view of the careful examination made afterward, would seem to have been, in many cases, symptomatic.

CLASSIFICATION OF CASES.

Accordingly, I will classify the cases as they are numbered in the table.

1. SYMPTOMATIC CLASS.¹

12 cases of enteritis (Cases 1, 6, 7, 15, 18, 21, 25, 26, 28, 30, 31, 33).

1 case of anemia (Case 17).

1 case of constipation (Case 13).

2 cases of malnutrition (Cases 16 and 30).

3 cases of indigestion (Cases 12, 32, 34).

1. *Wien. med. Wochschr.*, 1894, xlv, pp. 401, 464, 516, 550, 613. Medical Association, at the Fifty-seventh Annual Session, June, 1906.

1. *Wien. med. Wochschr.*, 1894, xlv, pp. 401, 464, 516, 650, 613,

2. ACUTE INFECTIONS.

2 cases of vaginitis (Cases 23 and 24).

2 cases of pneumonia (Cases 4 and 29).

1 case of bronchitis (Case 5).

3. CHRONIC DISEASES DUE TO DISTURBED NUTRITION.

4 cases of rickets (Cases 8, 9, 10, 20).

1 case of scurvy (Case 3).

1 case of marasmus (Case 13).

4. CONGENITAL DISEASES.

1 case of syphilis (Case 11).

1 case of cerebral palsy (Case 23).

1 case of cretinism (Case 36).

5. CASES IN WHICH THE BLOOD FINDINGS ARE OF INDIVIDUAL INTEREST.

1 case of hemiplegia (Case 27).

1 case of icterus neonatorum (Case 2).

1 case of goiter (Case 19).

1 case of cretinism (Case 36).

BLOOD EXAMINATION.

The normal number of red blood cells in children² is fairly constant after the eighth day, at about 5,000,000. In the cases of anemia, however, I wish to note that the red blood counts varied from 8,850,000 to 3,256,000. The white count,³ which normally is about 10,000, varies from 6,600 to 53,000.

The differential count of white cells in infants⁴ is: Small lymphocytes, 50 to 70; large lymphocytes, 6 to 14; neutrophils, 28 to 40; eosinophiles, 5 to 10. In these cases we have differential counts as follows: Small lymphocytes, 4 to 77; large lymphocytes, 6 to 56; neutrophils, 8 to 87; eosinophiles, 0 to 10.

Normal hemoglobin in children is 85 per cent.⁵ In these cases the hemoglobin ranges between 45 and 100. The color index, which is normally 8.5, ranges between 1.5 and .4.

Even a hasty glance at the table will convince one that those cases having a normal red count are normal in respect to the other findings. Hence the logical conclusion to make in regard to the variation of the red count, namely, that there is a concentration of the blood in those cases showing clinical aspects of anemia and high red counts.⁶

If the anemia were due to similar etiologic factors, taking into account the ages of the patients and digestive leucocytosis, we would expect a similarity in the white and differential counts; but a comparison of Cases 1 and 6, 16 and 30, and of 12 and 34 will readily convince one that there are not similar causes exciting the leucocytosis.⁷ The same condition will hold true when a comparison of the percentage of hemoglobin is made if we take into account the number of red cells, which condition is represented by the color index.⁸

Again, if we attempt to account for the enlarged spleens on the same basis, and a comparison is made of Cases 3 to 18 and Case 30, we are compelled to look for a different cause to account for the variations in the number and differential count of the white cells. The same

2. Sorenson. Cited by von Lulbeck: "Grundriss einer klinischen Path. des Blutes," Jena., 1896.

3. Whitney and Wentworth. Cited by Roach: "Pediatrics," 1896, page 348.

4. Gondobin: *Jahrb. f. Kinderh.*, 18893, v, No. 35, p. 187.

5. Da Costa: "Hematology," 1902, p. 280.

6. Oppenheim: *Deutsch. med. Wochschr.*, 1889, No. 42; also Hay: *Jour. Anat. and Physiol.*, 1882, p. 430.

7. Counts were made a sufficient time after feeding to eliminate digestive leucocytosis, which, according to Cabot, might vary from 1 to 30,000.

8. Cabot: "Clinical Examination of Blood," p. 303, 1901.