NOTES.

The Editor desires to point out that the pages of the Journal are open for the inclusion of short notes dealing with analytical practice and kindred matters. Such notes are submitted to the Publication Committee in the usual manner.

SOUR MILK SAMPLES AND GERBER'S FAT METHOD.

I HAVE recently had occasion to re-test, after a lapse of several weeks, a number of milk samples for fat by Gerber's method. These had been "preserved" by the addition of about 0.05 per cent. of formalin, which of course is not effectual for more than a few days, and the samples were very sour and decomposed. About 2 per cent. of strong ammonia was added to make pipetting possible, and, after correction for this, the results were up to 0.2 per cent. higher than those of the original tests. As would be expected, the ammonia is not responsible for this, as eight tests on five different milks averaged—

With ammonia	...	...	3.67
With ammonia, corrected	...	...	3.74
Without ammonia	...	...	3.76

The possibility of the formation of amyl esters suggested itself, and a sample of milk was divided into several parts, to which acetic, lactic, and butyric acids in known amounts were added. To make all comparable, 0.05 per cent. of formalin was also added, and 2 per cent. strong ammonia to neutralise the acidity. The following table shows the apparent increase in fat over the control, also the apparent increase in the case of several sour samples tested at the same time.

	Total Acid (as Lactic Acid).	Approximate Increase in Fat per Cent.
Milk + 0.5 per cent. acetic acid	0.57	0.10
" + 0.5 " butyric "	0.42	0.32
" + 1.0 " " "	0.63	0.57
" + 0.5 " lactic "	0.70	0.05
" + 1.0 " " "	1.20	Nil
" + { 0.17 " acetic " }	0.57	0.12
" + { 0.17 " butyric " }		
" + { 0.17 " lactic " }		
" + { 0.17 " acetic " }		
" + { 0.33 " butyric " }	0.80	0.19
" + { 0.33 " lactic " }		
Milk 3 days old, no smell of butyric acid	0.72	-0.06
" 1 week old, " " "	1.02	0.18
" 3 weeks old, smell of butyric acid	1.12	0.12
" 5 " " " "	2.10	0.19

The results in the case of the artificially soured samples were corrected for the volume of acid added, and those of the naturally sour samples for the ammonia. The fat separated from the most acid samples had a distinct odour of amyl esters.

It is evident that the use of Gerber's method for sour milk is apt to give very

misleading results, owing to the formation of amyl esters of the lower aliphatic acids. Lactic acid appears to have no effect, acetic acid little, but butyric acid a large effect. I was unable to try the effect of propionic acid, but one would expect it to be intermediate between that of acetic and butyric acids, in which case it would be an important source of error.

F. E. DAY.

LANSDOWNE, LIMERICK.

THE IDENTIFICATION OF ACIDS BY PHENACYL BROMIDE.

A useful method for the identification of organic acids by means of the m.pt. of their phenacyl esters has been described by Rather and Reid (*J. Amer. Chem. Soc.*, 1919, **41**, 75) and abstracted in the *ANALYST* (1919, **44**, 103).

The method depends upon the use of ω -bromoacetophenone for the esterification of the acids; and the writer thinks it worth while to call attention to the fact—learned by unpleasant experience—that special care should be taken in preparing and using this substance and its solution. Even small quantities produce very painful blisters on the skin if allowed to come in contact therewith. The substance also has a very irritating effect upon the eyes.

H. E. Cox.

NEWPORT, MON.

MASKING THE PRESENCE OF FREE SALICYLIC ACID IN ACETYL SALICYLIC ACID.

The writer, (*Pharm. J.*, 1920, **51**, 90), in a paper describing a new method for the estimation of free acetic acid in acetyl salicylic acid, also drew attention to the masking of the free salicylic acid present in aspirin tablets when tested for with ferric chloride test solution. The substance causing this was tartaric acid, some colourless non-ionised salt of iron being formed, there being no free ferric ions available for the formation of the violet-coloured ferric salicylate. Certain other strong acids will produce the same phenomenon, particularly citric acid (1.0 per cent. of the latter being sufficient to mask the presence of about 0.2 per cent. of free salicylic acid).

Tartaric acid, citric acid, etc., when present, can readily be detected after removal of the salicylic acid, acetyl salicylic acid, etc., from a cold aqueous extract of the tablet, and, conversely, the free salicylic acid can be estimated colorimetrically, after separation by means of some suitable solvent, such as ether, chloroform, etc., which does not extract the masking substance; when either tartaric or citric acid is present, a mixture of equal parts of ether and petroleum ether is a better solvent than ether alone, since both acids are slightly soluble in ether. All samples of aspirin tablets, therefore, which, after being powdered, give no violet coloration with ferric chloride test solution, should be viewed with suspicion.

A. NUTTER SMITH.

MESSRS. BOOTS' PURE DRUG CO., LTD.
NOTTINGHAM.

THE TREATMENT OF RED LITMUS PAPER.

Owing to the sizing of the original paper with rosin (colophony), it is frequently found that red bibulous litmus papers are not readily wetted by aqueous solutions, and that the colour indication is indefinite and slow in appearance.

This trouble may be obviated by the following procedure: A number of litmus books are rolled into a double layer of filter paper and inserted, backs downwards, in a Soxhlet extractor. They are then covered with 85 per cent. (or stronger) alcohol,

containing 1 per cent. of concentrated hydrochloric acid, and allowed to soak overnight, when a further quantity of acid alcohol is added, and extraction carried through in the usual manner, until no more red colouring matter is removed. The books are then squeezed dry in a cloth, and finally dried in an oven at 100° C. The alcohol removes, in addition to the rosin, some of the colour from the litmus and some from the red dye of the bookbinding.

T. J. WARD.

THE LABORATORY,
STAG BREWERY, S.W. 1.