THERAPEUTICS.

UNDER THE CHARGE OF

REYNOLD WEBB WILCOX, M.D., I.L.D.,

PROFESSOR OF MEDICINE AND THERAPEUTICS IN THE NEW YORK POST-GRADUATE MEDICAL SCHOOL AND HOSPITAL; VISITING PHYSICIAN TO ST. MARK'S HOSPITAL,

ASSISTED BY

HENRY HUBBARD PELTON, A.M., M.D.,

INSTRUCTOR IN MEDICINE IN THE NEW YORK POST-GRADUATE MEDICAL SCHOOL AND HOSPITAL;
CHIEF OF THE MEDICAL CLINIC, PRESBYTERIAN HOSPITAL DISPENSARY, NEW YORK.

The Treatment of Infantile Diarrhoea.—PERRET (*La quinzaine therap.*, 1907, vi, 450) advises careful feeding and intestinal irrigations with boiled water, 0.02 per cent. boric acid or naphthol solution, or normal saline solution. The irrigations facilitate the elimination of toxic substances from the intestine and supply the loss of the body fluids caused by the diarrhoea. During irrigation the child should be upon its right side; a soft, rubber catheter should be used and the receptacle of the fountain syringe, through which the lavage is given, should be elevated not more than five or six inches above the patient. The quantity employed should be from one pint to one quart and the fluid should be cold if the patient's temperature is elevated, hot if the latter is subnormal. As an intestinal antiseptic, three powders consisting each of 5 grains of bismuth salicylate and $2\frac{1}{2}$ grains of benzonaphthol may be given daily. Lactic acid should not be prescribed if the patient is taking milk. If the child is less than one year old he may take a teaspoonful of the following mixture every half hour: lactic acid, 3 parts; quince syrup, 25 parts; distilled water, 100 parts. If older than one year dessertspoonful doses may be given. Bismuth in a mixture containing soluble bismuth phosphate, 2 parts; simple syrup, 10 parts; distilled water 90 parts may act well; the dose is one teaspoonful every hour. If vomiting is present, gastric lavage may be indicated. Injections of artificial serum are often very useful.

Prevention of the Formation of Gallstones.—LANGHELD (*The Therapist*, 1907, xvii, 4) states that the reformation of gallstones can be prevented in a very simple manner by exercising pressure upon the gall-bladder, through displacing the liver by about two inches, the activity of the viscus being thus considerably enhanced. The procedure is as follows: The patient, in loose clothing, sitting upright, leaning against the back of an arm-chair, places his arms on the arm-rests and takes a deep breath, holding it while counting 15, breathes out deeply and again counts 15 before inspiring again. Thus the liver is moved downward by about two inches. Undertaking the exercise for about fifteen minutes, morning, and evening is sufficient for the purpose. For those who have suffered from cholelithiasis it is wise to continue the measure permanently.