THE LARYNGOSCOPE.

VOL. XVII. ST. LOUIS, MO., OCTOBER, 1907. No. 10.

ORIGINAL COMMUNICATIONS.

(Original Communications are received with the understanding
that they are contributed exclusively to THE LARYNGOSCOPE.)

SURGERY OF THE LABYRINTH.*

BY JOHN D. RICHARDS, M. D., NEW YORK.

The following paper is based upon 11 cases of labyrinthine sup-
puration and necrosis.† Of these, four occurred in the course of
acute or subacute mastoiditis; seven, in the chronic suppuration.
In *four*, neither symptoms were present, nor was there anything in
the previous histories suggesting inner ear involvement; the laby-
rinthine invasion was discovered at the time of the mastoid opera-
tion.

In *three*, no symptoms were present at the time of operation,
though the previous history in each pointed to labyrinthine dis-
turbance.

In *four*, decided symptoms were present at the time of operation.

In *three* of the eleven cases infective sinus thrombosis compli-
cated the labyrinthine disease. In two of these the jugulars were
thrombosed.

This relative frequency of sinus involvement is of surgical inter-
est when we consider the vascular connection between the veins of
the vestibule, the semicircular canals and the lateral sinus, and those
of the cochlear and the inferior petrosal.

Three of the eleven cases died. The first death was from menin-
gitis resulting from operative interference. The second death
occurred in one of the sinus thrombosis cases and was due
to meningitis resulting from operative interference. Though
the sinus was occupied by an infected clot, the thrombosis was not
an important factor in the patient's death, as the post-mortem re-

* Read before the Thirteenth Annual Meeting of the American Laryngo-
logical, Rhinological and Otological Society, New York City, May 30, 31, and
June 1, 1907.

† The writer has since operated upon a twelfth case in which a cholesteatomatous
mass invaded the internal ear through the oval window, forming a cast of a portion of
the labyrinth.

vealed the pathway of infection to be from the labyrinth along the course of the auditory nerve. The third death occurred in one of the sinus thrombosis cases and was due solely to the infective thrombosis. In this instance the jugular was occluded by an infected purulent clot which extended below the clavicle into the deep thoracic portion of the vein, it being impossible to get below the cardiac limit of the clot. In this instance the labyrinth was in a perfectly healthy condition and had been so for a period of over ten days before the development of the sinus symptoms. The first two deaths resulted from improper technique and the specific errors—for the sake of emphasis—will be referred to in their respective places. I regard these deaths as sacrifices which could now be avoided.

The labyrinthine capsule showed the following lesions:

In five, the only visible lesion was a loss of the prominence of the horizontal semicircular canal.

In one, the oval window was the seat of fistula through which granulations protruded. The vestibule, the semicircular canals, and the lower half of the first cochlear whorl were involved.

In one the prominence of the horizontal canal had disappeared. The oval window was perforated and filled with granulations; a second perforation was present through the centre of the promontory through which polypoid granulations sprouted and pus oozed. The canal system was filled with granulations, the vestibule and cochlea with granulations and pus.

In three, the horizontal semicircular canal had disappeared and the oval window was the seat of fistula and filled with granulations. The canal system, the vestibule and the cochlea were involved in each.

In one, there was a perforation just above the horizontal semicircular canal in the region of the solid angle. The inner vestibular wall was also the seat of a fistula which communicated with a large epidural cerebellar abscess in the vicinity of the internal auditory meatus. The entire labyrinth was involved in the purulent process, which had also destroyed the greater part of the petrous pyramid, necessitating its removal.

Of the 11 cases, therefore, the horizontal semicircular canal, either alone or in combination with other lesions of the capsule, was involved in nine.

The oval window was perforated in five; the promontory, or first cochlear whorl, in one.

The labyrinthine capsule in the solid angle of the semicircular canals in one.

The inner vestibular wall was perforated once.

In all, there were seventeen perforations in the eleven cases,—sixteen through the outer labyrinthine wall, but one through the inner. This is of surgical importance, and one of two conclusions is evident—either in the great majority of cases the labyrinth is invaded by direct extension through the capsule from without, or else the outer wall of the labyrinthine capsule is exceedingly more vulnerable than the inner. Both are true.

The cases show conclusively that the fistulous openings in the external capsule do not always represent avenues of invasion from without inward, but avenues of escape through which pus is successfully discharged from the labyrinth into the middle ear.

In every instance in which the vestibule has been involved the canal system and the cochlea—either in whole or in part—have been involved also.

The first question which from a surgical standpoint we are called upon to decide—and it is one with which we are frequently confronted—is the following: If after the diagnosis of labyrinthine suppuration is made we operate and find in the outer capsule of the labyrinth no fistula, shall we explore it upon the symptoms, fork tests, etc?

To take a correct surgical attitude with reference to this important point, we will consider first, briefly, the diagnostic value of certain chief symptoms and tests; second, the dangers which attend the exploration of the labyrinth; and third, the dangers which attend the non-exploration of the labyrinth. I trust that a brief reference to the surgical value of the first will not be misinterpreted, in the present paper which is concerned chiefly with the treatment of labyrinthine suppuration.

1st. In the vertigo, vomiting, nystagmus and disturbances of equilibrium of labyrinthine origin, there is nothing characteristic.

2d. Even if we can exclude cerebral and cerebellar disease and can fix these symptoms definitely upon the labyrinth, they are not indicative of actual invasion of the labyrinth; they merely signify disturbance of labyrinthine function; but this occurs with comparative frequency from causes entirely extra-labyrinthine: from polypi, cholesteatoma, pus in the middle ear under pressure, etc.

3d. The mere fact that these symptoms are associated with a suppurative process is of little diagnostic value, as the extra-laby-

rinthine causes which produce these symptoms by transmitted pressure are, as a rule, associated with a suppurative process also.

4th. Cases occur, presenting all the above symptoms and in addition fork tests, which, according to orthodox interpretation, point to labyrinthine involvement which when operated upon show no evidence of labyrinthine involvement in the outer capsule, and which completely recover without having the labyrinth explored; these cases are not uncommon.

5th. In my own cases the forks have been of doubtful value in differentiating middle ear from labyrinthine lesions.

Briefly: seven of the eleven cases lateralized to the side of labyrinthine lesion. In six bone conduction was not shortened in duration. Of seven in which the greater part of the entire labyrinth was removed, four not only lateralized to the side of the labyrinthine lesion and bone conduction was not shortened in duration, but by air conduction forks were heard throughout a considerable range of the scale.

6th. In every case in which I have seen the labyrinth encroached upon by either operation or disease, the organ has been rendered valueless for practical purposes of hearing. I refer also to several cases of acute mastoiditis in which I have seen the horizontal semicircular canal injured by accident, and which cases I have had the opportunity to examine.

7th. To expose the interior of a normal labyrinth to an infected cavity is highly dangerous to life. What the mortality is under these conditions, I do not know, as I am unacquainted with the literature of this subject; but from my own experience, I realize that the opening of a diseased labyrinth is a serious thing, and I am confident that it is far more dangerous to open a normal labyrinth in the presence of an infected cavity than a diseased one; for in the latter the pathways along which infection may travel to the intracranial cavity and sinuses have had the opportunity to become sealed through inflammatory processes.

8th. If we do not explore the labyrinth under the conditions mentioned, we may overlook certain cases of labyrinthine suppuration—for my own cases show conclusively that the labyrinth is not always invaded directly from without by encroachment upon the capsule, but that it may be infected by septic particles being transmitted to its interior along the avenue of the vessels, the outer capsule showing no perforation whatever, i. e., by a process of local metastasis. The pathological process is analogous to that in which

infection of the sigmoid sinus occurs from the antrum along the avenue of the communicating vessels, the intervening mastoid structure showing no apparent involvement.

Should we overlook such a case, we court two dangers:

1. Intracranial infection taking place along the communicating avenues with the labyrinth.
2. That the inner wall of the labyrinth will perforate before its outer wall.

With reference to the first: of the eleven cases mentioned, three had infective sinus thrombosis. This relative frequency throws some suspicion upon the labyrinth as a causative factor, for this proportion of sinus infections is entirely too high for the ordinary run of suppurative cases either acute or chronic.

That the infected labyrinth was responsible for one of the infected sinuses is probable, inasmuch as the symptoms of sinus involvement appeared six weeks subsequent to the cessation of aural discharge. This is suggestive of the role played by the communicating avenues from the labyrinth as infection carriers to the intracranial cavity, and the danger of overlooking such a labyrinthine supuration.

The frequency with which the cerebellar fossa is found post mortem to be the site of greatest pathological activity in cases of purulent lepto-meningitis, originating from suppurative disease of the petrous pyramid is also suggestive of the important part played by the labyrinth as a distributing center of infection to the intracranial cavity.

With reference to the second—That the inner labyrinthine wall will perforate before the outer, is not borne out by clinical fact, nor, would we suppose it on anatomical grounds.

Of the seventeen perforations in the labyrinthine capsule in the eleven cases, sixteen were through the outer wall, but one through the inner wall. This ratio of sixteen to one is, of course, misleading, inasmuch as the majority of labyrinthine infections occur by direct extension through the capsule from without, under which circumstances the inner wall of the labyrinth is not subjected to a comparative test of vulnerability.

Considering, however, the doubtful value of symptoms as indicative of actual invasion of the labyrinth, the difficulty of eliminating cerebellar disease, the unreliability of tuning forks in differentiating in this class of cases middle ear from labyrinthine lesions, the practical certainty that we will destroy the organ for the purpose of

useful hearing, the actual danger to life should we commit the error of opening a normal labyrinth to an infected cavity, the certainty that the outer labyrinthine wall will in the great majority of cases perforate before the inner wall—the correct surgical attitude is, not to enter the labyrinth upon symptoms, etc., alone at the primary operation unless there is direct evidence that the labyrinth is involved. While I can imagine a combination of symptoms and circumstances which would cause me to open the labyrinth at the primary operation, even in the absence of recognizable labyrinthine invasion, I have as yet met no case which presented such features, and it is to be sincerely hoped that the labyrinth will not suffer the indignity of reckless exploration and that it will not be entered without a due consideration of the possible results to life.

What route shall we adopt?

The following facts have bearing upon the selection of the route by which we enter the labyrinth. From a surgical standpoint the cases may be divided into two classes.

Of the first class, the first five cases are illustrative, in which a loss of the horizontal semicircular canal represented the only lesion in the capsule. These cases represent caries of the labyrinthine capsule in which as the extension proceeds from without inward by a process of erosion, the limit to which the labyrinth is surgically involved is, as a rule, represented by the carious area. In such cases we merely remove the carious area and follow it to its limit, and the route which we select depends upon the site of the lesion, which, as a rule, is in the horizontal semicircular canal. The reasons for the frequency with which the horizontal canal are involved are evident.

1st. The external semicircular canal, unlike the posterior and superior canals which lie deeply embedded, juts out upon the floor of the aditus as a small ivory-like hump and is exposed to the supuration process.

2d. Its exposure occurs at the point of greatest constriction between the tympanic and antral cavities, and at a point of great pathological activity. The canal is not only subjected to the action of pus but to the deleterious combination of pus under pressure.

3d. The aditus is an unyielding bony ring in the circumference of which the external semicircular canal represents one of its weakest points and it succumbs accordingly.

4th. The vulnerable point in the external semicircular canal is not at the extreme apex of its arch—the point of greatest exposure

—but at a point several millimeters below the summit of its ivory-like cap. This is due to structural differences at the points mentioned and to the fact that the base of this ivory-like hump is encroached upon or surrounded by a mass of cancellated bone which is easily disintegrated and in these cases always diseased.

I have seen no instance in which the extreme apex of the arch has been eroded: in every instance in which the canal has been involved the entire cap has disappeared.

In chronic suppurative cases we frequently have the opportunity to observe the effect of this undermining process as it has attained to different degrees of encroachment upon the lumen of the canal.

We find that the prominence of the canal presents at its base a thin, sharp edge so undermined that it would appear that by placing the finger-nail beneath it we could separate it as a little thin shell.

The manner of invasion in these cases is evident—the labyrinth is opened by a process of gradual erosion from without.

Of the second class, the remaining six cases are illustrative. In these the labyrinth is involved in a true suppurative process, and its various compartments are filled with granulations or pus, or both. These cases are in every way more formidable, more destructive and widespread, and more highly infective than the first and require bolder intervention. The vestibule is the site of greatest pathological activity, and in every instance in which I have found the vestibule involved, the canal system and the cochlea, either in whole or in part, have been involved also. This suggests the oval window as the most common avenue of entrance in the second class of cases, and this is in accordance with the pathological findings in the capsule as shown in the previous cases. In the true suppurative cases the oval window was perforated in every instance with one exception. We may also have in an individual case the two pathological processes mentioned at work at the same time.

Inasmuch, therefore, as the exploration of the canal system, involving to greater or less extent its sacrifice, is necessary in this class of cases, the route which we adopt resolves itself into the selection of that which from a manipulative standpoint is the most convenient, which least endangers the important structures in surgical relation, and which allows us to fully expose the limits of the disease. The only route which fully meets these requirements is posterior to the facial nerve, and the point through which it is safest to enter the labyrinth for reasons to be mentioned later, is the solid angle of the semicircular canals.

We prefer to enter the vestibule posterior to the facial nerve instead of anterior to it, for the following reasons:

1st. When the vestibule is involved the semicircular canal system, in the great majority of cases, is involved also. This necessitates its exploration, which entails its sacrifice.

2nd. The sacrifice of the canal system, even though it is not extensively involved, creates only an ephemeral disturbance, and adds no danger to life, as the canal system is not in important communication with the intracranial cavity except by way of the vestibule.

3rd. We secure the maximum amount of working room, the advantage of which can be appreciated only when we have attempted the actual operation; and we are better enabled to work in the axis of the petrous pyramid, which is the direction of greatest safety.

4th. We get a field unsoiled by blood and gain a clear conception of the condition within the vestibule, of the ampullary areas of the canals, and of the inner vestibular wall.

5th. The route anterior to the facial nerve does not permit an exploration of the canal system which we know as a rule to be involved, nor of the ampullary areas, nor of the major part of the inner vestibular wall, but only of its lower anterior portion—consequently fistulae through the inner wall of the vestibule may be overlooked; I have had one such case in which had I adopted the anterior route, I would have committed this error and sacrificed the patient.

6th. It is based upon the erroneous idea that drainage is the chief desideratum and will suffice. Mere drainage in this class of cases will not suffice, for not only is the interior of the labyrinth involved, but its capsule is, as a rule, profoundly diseased and such portions of it must be removed. In addition the complicated system of channels represented by the labyrinth cannot be satisfactorily drained from any one point, as from an opening in the outer vestibular wall.

That drainage through a given point will not suffice is shown by the fact that in many of these cases the outer capsule contained multiple fistulae—yet with multiple fistulae present, drainage was not accomplished. The mere existence of multiple fistulae, properly interpreted, is a plea for more efficient drainage than can be accomplished by this means. The attempt to drain the labyrinth through an opening in the outer vestibular wall, while it may in certain instances succeed, is comparable to the attempt to relieve cases of suppurative mastoiditis by merely opening the antrum. We

should make the effort to place the present operative procedure upon a thoroughly scientific basis from the first.

7th. From a manipulative standpoint the anterior route is far more dangerous; it is the more cramped; the anterior canal wall interferes seriously with manipulation in the upper portion of the cavity; we are working in a blind pit and are forced to do considerable work from without inward, the direction of greatest danger; first, as the dome of the jugular bulb may rise high, undermining the outer vestibular wall; and second, as the inner wall of the vestibule bulges outward at this point, making the anterior portion of the vestibule comparatively narrow from within outward, an injury to the internal auditory meatus is a much more probable accident.

What are the steps of the technique?

As a description of the complete exenteration of the labyrinth will embrace the various steps, that alone will be mentioned.

1st. A Schwartze-Stacke operation, by which we secure the maximum amount of working room, and in which we exenterate the bony angle included between the groove of the sinus knee and the under surface of the middle fossa, which better enables us to work in the axis of the petrous pyramid. We lower the facial ridge to its absolute limit, remove the fringe of bone on the anterior aspect of the facial ridge back to the descending limb of the facial nerve, remove the outer wall of the hypotympanum, and lower the level of the canal floor, securing by these steps the maximum exposure of the outer wall of the vestibule, and the dome of the jugular bulb should it rise high.

As exploration of the cochlea may be necessary, it is important that we see the exact position of the carotid artery by the exposure of its canal. In consequence we shave down the convexity of the anterior wall of the auditory canal, remove the lip of bone overhanging the mouth of the tube and evulse the tensor tympani.

This exposes the tube to curettement by which we relieve the field of blood, and it gives the maximum width to the apex of the cavity. Should the consistency of the bone permit, the arches of the semicircular canals should be delineated. This enables us to work with accuracy. (Fig. II.) The cavity should be cleansed; the tube packed with adrenalin gauze; the field rendered bloodless; the instruments and hands resterilized.

We next remove the prominence of the horizontal semicircular canal; it is a treacherous structure. The cutting edge of the chisel is placed at a point below the summit, but well above the level of

the fallopian canal, for this prominence separates along definite planes of cleavage, and as the outer lip of this semicircular canal is intimate with the fallopian, a fissure in the former may extend out into the latter. Should the plane of cleavage be on a level with or below the fallopian canal, the facial nerve may either be exposed by having its roof removed, as shown in Fig. X, or else completely undermined (Fig. V), under which latter circumstances we may definitely expect paralysis for it will ensue. The stroke should be made in a direction corresponding to the plane of the canal. The remaining canals are next uncapped, and the condition of the interior is noted. It will be found that the interior of the external semicircular canal most frequently of all shows pathological change. In opening the superior canal a curved gouge should be used. It permits the stroke to be made in the direction of safety. The danger arises from the non-support of this canal by solid bony structure.

Owing to the contrast between the dark interior of the canals and the brilliantly illuminated white bone surrounding them, the labyrinthine fluid may appear dark and be mistaken for blood or granulations; we should not be deceived by this illusion.

We next enter the vestibule through the solid angle of the semicircular canals by creating at this point a conical pit with its apex directed inward and gradually lowered until it enters the vestibule. As it is necessary to remove the inner lip of the horizontal semicircular canal, during this step the chisel should be held perpendicular to the plane of cleavage, and under no circumstances should it impinge upon the outer lip, which is intimately associated with the fallopian canal, together with which it is left as a bridge carrying the nerve which spans the vestibule. (Fig. IV.)

The opening in the vestibule is now enlarged until a full exposure is obtained of this portion of the cavity. Its inner wall should be searched for fistulae. In enlarging this cavity no pressure should be made upon the bridge. For this reason a curette is a dangerous instrument. A small sharp gouge is safer.

Owing to widespread necrosis, it is necessary in some instances to sacrifice the bridge of bone carrying the facial nerve. To accomplish this without injury to the facial, we select a curette with its cutting edge turned backward, utilize the superior rim of the bone cavity as a fulcrum and shave off from above downward in a direction parallel to the course of the nerve, the roof of the fallopian canal, thus leaving the nerve exposed in its gutter of bone, from

which it may be separated and lifted without injury. (Fig. VI.) Any filaments given off from the nerve should be cut and not torn from this structure, as unnecessary traumatism is committed. We next remove such portions of the bridge as are necrotic, but no more, for the nerve here represents a curve, and should it lose the entire support of its bony gutter, it apparently elongates and consequently sags or kinks and becoming enmeshed in this vicious position by the granulations, has its functions interfered with later.

In this condition the nerve stretches as an exposed structure from its point of emergence low down on the facial ridge, across the cavity of the vestibule to its entry into the inner wall of the tympanus, corresponding to a point above and anterior to the original site of the oval window. (Fig. V.)

In knocking out the inner edge of the bony ridge, which corresponds to the upper inner wall of the tympanum, the stroke of the chisel should not be made from behind forward in a direction parallel to the course of the nerve, but from above downward, or, from before backward; i. e., in a direction perpendicular to the course of the nerve and to the fallopian canal.

The reason for this is that the bone at this point tends to separate when struck from behind forward, along a plane of cleavage which crosses the fallopian canal, causing its fracture; as the fallopian canal at this point represents the greater portion of an exposed cylinder (Fig. I), its fracture results in the facial nerve being encompassed by a little annulus or cylinder of bone, which may be slid up and down upon the nerve, just as a ring upon a finger. This complication is—so far as the integrity of the nerve is concerned—a formidable one, and it requires the greatest patience to remove the little annulus without causing injury to the facial. To accomplish it we steady the annulus with mouse-tooth thumb forceps and with a pair of ronguers—the jaws of which are accurately apposed—crush it in a direction parallel to the course of the nerve.

To avoid this complication, it is only necessary before attempting the removal of the upper portion of the bony ridge that the fallopian canal should have been thoroughly converted into a gutter, by shaving off its roof, as mentioned, and by making the stroke in the direction indicated. Should the accident now occur, the nerve can be easily freed from the little mass of bone by slipping it through the open side of the annulus, or cylinder.

Should we have to expose the nerve as mentioned and shown in Fig. V, no attempt should be made to retract it or draw it to

one side for the sake of gaining room for manipulation, as this is unnecessary; for the fallopian canal at the points at which the nerve emerges from and enters the bone is represented by sharp, serrated margins, against which if the nerve is drawn it may be lacerated at its fixed points. The sharp spicules should be removed so that the margin of the canal at these respective points is represented by an even rim.

The next step is to expose the antero-inferior cavity of the vestibule by removing the posterior aspect of the promontory and the outer vestibular wall. (Fig. VII.) We select a gouge, the width of which corresponds to the distance between the round and oval windows. The cutting edge straddles the little bridge of bone separating these two openings, and the stroke—which is slight, for the posterior aspect of the promontory is brittle and fractures easily—is directed from above, downward and forward in the direction of the first cochlear turn. We must at this step think of the possible position of the dome of the jugular bulb.

The greatest care should be taken that the cutting edge of the chisel does not cross the cavity of the vestibule and impinge upon its inner wall, for the inner wall at this point bulges outward and is separated from the internal auditory meatus by a brittle partition of bone not more than $\frac{1}{32}$ of an inch in thickness. (Fig. XVI.) Should the inner wall be fractured, we lose cerebro-spinal fluid and aside from the inconvenience caused by this accident, the infected operative cavity is placed in direct communication with the intra-cranial cavity and the patient's life is jeopardized; meningitis will probably result.

As the cerebro-spinal fluid is under pressure and is of low specific gravity, gauze does not act as an efficient plug to the rent; sterile wax is preferable.

By enlarging this opening, we secure, with the previous steps, the full exposure of the vestibular cavity and the beginning of the first cochlear whorl. Should we now find that the disease has invaded the cochlea, and this is the rule in the second class of cases, we must continue the exploration of the cochlea until we have followed the diseased process to its legitimate end.

We next remove the roof of the first cochlear whorl from behind forward, exposing its interior to a point just short of the carotid eminence. (Fig. V.) The instrument of preference is a thin, sharp gouge with no shoulder, the width of which is slightly greater than that of the cochlear whorl. Four structures are to be

avoided; the dome of the jugular bulb below, the eminence of the carotid canal in front, the base of the modiolus and the internal auditory meatus internally. The danger to the first two is slight; to the second two it is imminent, and injury to these structures is to be avoided only with the exercise of care.

By confining the removal of the cochlear shell entirely to the roof of its first whorl and not allowing the gouge to impinge upon the inner wall of the cochlear turn these last dangers may be averted. From the carotid artery in front, the cavity of the first cochlear turn is separated by a hard cuff of bone which serves the purpose of an efficient bumper: though thin this partition is sufficient. (Fig. XII.)

If we now find that the limit of the disease has not been reached, we must explore the remaining cochlea. This constitutes by far the most difficult and dangerous step of the procedure; for the cochlea, which represents an extremely small cavity encased in a brittle shell of bone, is hemmed in on all sides by structures which we cannot afford to injure. In front is the carotid artery, below the dome of the jugular bulb, behind the internal auditory meatus, above and behind in immediate proximity to the second half of the first cochlear turn, the knee of the facial nerve. A circle a quarter of an inch in diameter could be so placed as to pass through the majority of these structures.

Were the above factors the only ones to be considered, it would be comparatively easy to select a point on the cochlear shell, which from dead house work we had found to be a safe one, and open its cavity. But the difficulty lies in the fact that within this shell of bone is contained a structure which from its position is exposed to injury, and which from a surgical standpoint is the most treacherous of the internal ear; I refer to the modiolus.

Before approaching the cochlea it will be well to consider certain anatomical features of the modiolus which are of surgical importance. The modiolus represents a small pyramid of bone seated upon the internal auditory meatus, decreasing in size from base to apex. Its apex is its weakest point, but the next weakest point is not immediately below its apex, but at the extreme base; for its base is ex-cavated as shown in Fig. XI (Politzer), by the internal auditory meatus, and the pyramid consequently rests upon a mere rim of thin, brittle bone. If the chisel is applied to the pyramid well above its base and a stroke made, the fracture does not take place at the point of applied violence but at its base; when this oc-

curs the pyramid fractures completely round the circumference of its base and separates as a single piece of bone. (Fig. XIV.) The internal auditory meatus is consequently opened throughout its entire circumference, and as the diameter of the base of the pyramid, or the rim of bone upon which it rests, is about 1-6 of an inch, the loss of cerebro-spinal fluid is rapid. The failure to appreciate the surgical importance of this anatomical feature of the modiolus resulted in the death of the first patient.

If we now examine the modiolus with a strong convex lens we see that the pyramid has an outer casing of brittle bone and a core which is porous, made so by canals running from base toward apex for the passage of various structures. These canals are not completely filled by the structures which they contain, and this permits the cerebrospinal fluid to penetrate out into the modiolus.

During the second operation in which the modiolus had been removed well down toward its base it was noticed that a seepage of cerebrospinal fluid took place through its stump; the intra-cranial cavity had been placed in gross communication with the infected cavity in an altogether unexpected way through the afore-mentioned channels, and the failure to appreciate the significance of this structural character of the modiolus resulted in the death of the second patient. In this case the infection could be traced along the cochlear branch of the auditory nerve. It now became important to determine how far down from the apex toward the base the pyramid could be removed without putting the intracranial and operative cavities in gross communication; for in dealing with the anterior half of the cochlear cavity it is absolutely necessary to get rid of a portion of the modiolus.

It can be shown upon increasing the tension of the intracranial fluid in a cadaver by injection, or in the living subject by pressure over the internal jugular vein, under which circumstances the cerebrospinal fluid will penetrate as far out as possible into the modiolus (i. e., grossly), that the pyramid may be removed from the apex toward its base, down to a point corresponding to the termination of the first cochlear whorl without causing the loss of cerebrospinal fluid; i. e., without placing the operative and intracranial cavities in gross communication. As will be shown later, this suffices for the complete exploration of the anterior half of cochlear cavity.

In approaching, therefore, the cochlea, we must remove its shell in such a way as not to injure the modiolus. We select a point in

the cochlear shell corresponding somewhat to the apex of the cochlear cavity, and with a thin sharp gouge shave it down until the dark interior of a cochlear whorl shows through the thin lamella of bone. The stroke should be made from above downward and forward in a direction corresponding to that of a cochlear whorl.

Not infrequently the shell of the cochlea is scalloped and the position of the cochlear turns roughly indicated. In this way a window is created in the cochlear shell (Fig. X), and we now enlarge this window, completely exposing the upper portion of the cavity. (Fig. IX.)

In enlarging this window the small gouge is the instrument of preference; an attempt to insinuate a very fine curette beneath the opening in the cochlear shell causes its back to impinge or press upon the modiolus, which sticks up as a little tent pole in the cavity, and this may result in the fracture of the pyramid at its base. The gouge merely removes the shell without endangering the pyramid.

To expose the second half of the first cochlear whorl it is necessary to remove the apex of the pyramid down to a point corresponding to the termination of the first cochlear whorl, as shown in Fig. XII. This done, we can look down over the stump of the pyramid, upon the roof of the second half of the first whorl, which is indicated by the probe in Fig. XII, and with a small gouge carefully break through its roof from above as shown in Fig. XIII, exposing in this way the entire interior of the cavity of the cochlea. In breaking through the roof over the last portion of the first whorl we are in direct relation to the facial nerve and the internal auditory meatus, both of which must be avoided.

The probe in Fig. XII is wound round the base of the modiolus merely for the sake of illustrating the relations of the roof of the second half of the first whorl. An attempt to withdraw it may cause the modiolus to fracture; such a step, of course, would not be taken during actual operation.

In one instance I have had to exenterate the major part of the petrous portion of the temporal bone and the following points have impressed me:

- 1st. That the petrous pyramid in this particular instance was pneumatic; we would expect to find such extensive involvement only in bones of pneumatic structure for reasons which are evident.

- 2d. Where removal of the petrous pyramid is necessary it should first be exenterated; later the shell removed; otherwise the dura of the posterior and middle fossae bulge into the wound, constrict

the operative field which is extremely deep, and close it to successful view. Retraction of these structures is then necessary, but the best retractor is the shell of the petrous pyramid, as it commits no trauma, occupies no space, and is accurately applied to the surfaces to be supported.

A second reason for first exenterating the petrous is that the dura enclosing the superior petrosal sinus is firmly attached to the upper margin of the posterior aspect of the pyramid and separates with reluctance from this uneven lip. In consequence the vessel may be torn; sharp hemorrhage follows which is difficult to control, as it is not easy to place a compression plug between the dura and the bone, as the two do not readily separate. If the accident is to occur, it is more convenient that it should occur toward the end of the procedures.

We should attempt to save the facial nerve by preserving a bridge of bone extending from the lower portion of the facial ridge to the internal auditory meatus; with patience this may be accomplished and, in the instance mentioned, was accomplished. Even though the nerve be sacrificed, we should still attempt to preserve the bridge, the purpose being to give the dura of the middle fossa a seat upon which it may rest that it does not sag deep into the wound and become sealed in this position to the granulation bed springing from the floor.

When the dura is not supported it sinks and shuts off the apex of the operative cavity, rendering this portion difficult to handle, and it further obstructs drainage. No amount of pressure by packing with gauze suffices to maintain it in a proper position, as the gauze soon becomes soaked and its supportive influence lost. Should, for any reason, reoperation be necessary—and we must remember that dyscrasiae may underlie such extensive destruction, and portions of necrotic bone may remain which circumstances at the time of operation either cause to overlook or to abandon—the dura sealed in its vicious position constitutes an efficient barrier to further operative interference. For this reason the operation should be thorough and all dead bone in the deep, apical portion of the pyramid removed. When the bridge is sacrificed the nerve, which represents a curve, is exposed over a considerable portion of its length and it kinks and becomes enmeshed later in the granulation bed; this interferes with function.

It is dangerous to attempt to remove sclerotic portions of the pyramid with the rongeur. The force requisite for a bite of the

forceps is great, the bone suddenly gives and as the jaws of the rongeur spring together, sharp slivers of bone are thrown with violence against the tightly stretched dura of the cerebellum, which makes an excellent target to be perforated. I have seen the posterior fossa opened in this way.

It is highly important to be familiar not only with the surgical anatomy in the operative position, but to know the consistency of the bone; to know the result that will follow a stroke of the chisel when made in a definite direction; for the bone of the petrous pyramid is structurally treacherous, changing rapidly from pneumatic to absolutely sclerotic structure, which may crack when struck far beyond the point of applied violence and in a direction contrary to that expected. We can obtain a practical familiarity with this important part of the technique only by repeated work upon the cadaver, and an attempt at a description of this part of the technique is useless. Generally speaking, we should, when attempting to remove sclerotic portions of the labyrinth, make the stroke in the long axis of the cavity which the sclerotic bone covers. While this is not invariably true, it is as near as the facts can be reduced to rule.

In operations upon the labyrinth a point of interest is to be noted with reference to the carotid artery, when through erosion of the carotid canal the artery is exposed to view, there is, as a rule, no visible pulsation in the vessel; I have seen the artery exposed in eight cases (some in chronic suppuration), and in only one instance was pulsation to be detected. If pressure is made upon the vessel with an applicator wound with cotton and the lumen of the vessel momentarily constricted, slight pulsation may be elicited. It is a wise provision that the vessel does not forcibly pulsate at this point and impinge upon the sharp serrated margin of the eroded canal; the walls of the artery in the canal are thin owing to a lack of the necessity for their full development.

The carotid canal serves a two-fold purpose: First, As the exposed artery in the canal shows no visible pulsation, or but a faint pulsation, it is evident that the carotid canal acts as a bumper to the pulse wave, diminishing its impact and shielding the brain from excessive shock.

Second, As the canal is rigid and does not permit the artery to dilate except within narrow limits, it has the practical effect of constricting that portion of the artery within its grasp. The effect is the same as when with the fingers we constrict the lumen of a rubber tube through which water is being forced by a rubber bulb, i. e.,

there is a tendency under these condition for the intermittent or remittent current to approximate a continuous stream. The carotid canal, therefore, not only protects the brain from excess of pulse impact, but it is a mechanism by which the inflow of blood to the brain is somewhat regulated and made more even and continuous. It is analogous to the hydraulic arrangement of the jugular bulb and sigmoid sinus on the venous side of the circulation, whereby an even outflow of blood from the brain is guaranteed.

During and following operation upon the labyrinth certain phenomena occur which are of exceeding interest.

If during anaesthesia we produce intermittent pressure upon the capitulum of the stapes, we may in certain instances induce nystagmus, which—should it occur—would rather point to the fact that the labyrinth is functioning and not grossly involved in a destructive process. This experiment may prove of some value from a diagnostic standpoint in determining labyrinthine involvement, when the capsule shows no fistula.

If during operation we irritate with a probe the interior of the semicircular canals we may at time induce nystagmus. In the cases which I have seen the nystagmus has been bilateral and in the horizontal plane, regardless of which canal was irritated. The irritation of the inner vestibular wall gives a similar result.

The ampullary areas of the semicircular canals are more sensitive to irritation than their arches.

The upper posterior portion of the inner vestibular wall appears to be its most sensitive part.

A nystagmus which prior to operation is very marked may not disappear under ordinary surgical anaesthesia until the semicircular canals are removed and the vestibule opened, when it may entirely cease.

If the horizontal semicircular canal is by accident injured as during the acute mastoid operation, vertigo, vomiting, and nystagmus may appear and may be marked; again, these symptoms may be entirely wanting. As a rule, the disturbances which follow the injury are exceedingly slight and ephemeral, and consist of horizontal nystagmus—most marked when the eyes are directed to the opposite side, and slight vertigo with a tendency on the part of the patient to fall away from the labyrinthine lesion. Profound disturbance of hearing, as a rule, follows.

If the operation has encroached to any extent upon a functioning labyrinth, particularly in cases where both the canal sys-

tem and the vestibule are removed, disturbed equilibrium, nystagmus, and vomiting, which is accompanied by nausea, as a rule, follow. The vertigo is more pronounced when an attempt is made to assume the sitting posture, and closure of the eyes augments the disturbed equilibrium. The patient, as a rule, tends to fall to the side opposite the labyrinthine lesion.

At first the nystagmus may be of marked rotary character—the globes rotate beneath the closed lids, and nystagmus is independent of an attempt to fix the eyes upon an object. Nystagmus may be present in whatever direction the eyes assume. In a few days a marked decrease occurs, and as a rule, though not always, the greatest excursion of the globes is in the horizontal plane. At first the oscillatory movements may be fine; later they are coarse, sluggish and jerky. The apparent fineness or coarseness of the movement appears to be dependent upon the rapidity of oscillation. The nystagmus, as a rule, presents the following features:

1st. It is most marked when the patient looks to either the extreme right or left lateral position.

2d. When the eyes are directed to the upper vertical meridian, slight nystagmus may be observed, which, at first may appear vertical in direction, but upon careful inspection a slight rotary movement will be seen. This latter is possibly due to the action of the extrinsic muscle of the eye.

3d. When the eyes are directed to the lower vertical meridian, the nystagmus is slightest, and, as a rule, absent, even though present in other meridians.

4th. The nystagmus is most marked when the eyes are directed to the extreme lateral position of the normal side. Even after the nystagmus has totally disappeared in all other meridians, it may still be made manifest when the eyes are turned from the labyrinthine lesion. Occasionally the patient tends to fall to the side of the labyrinthine lesion and not to the opposite side. When this occurs, the nystagmus may be most marked when the eyes are directed to the extreme lateral position of the operated side, or equally as marked as when turned to the normal side. There appears to be some harmony, as to direction, in which disturbed equilibrium and nystagmus are most manifest.

The vertigo and vomiting which follow operation upon the labyrinth, cease, as a rule, in a few days, and disturbed equilibrium and nystagmus soon subside. The patient, however, when attempting to walk a straight line, may show for some weeks or months sub-

sequent to operation a tendency to deviate from the involved side; he may so tend to fall immediately after operation. In other cases all disturbances of equilibrium disappear in a few days. This tendency to fall or to deviate from the side of the labyrinthine lesion occurs in cases in which the horizontal canal alone is removed, in which the canal system and vestibule are both removed, or in which the entire labyrinth is removed. In some cases a slight nystagmus may be observed months after the operation; it may persist for years.

In some instances where the nystagmus is not of sufficient degree to be manifest when the eyes are directed to infinity, it may be made so by having the patient suddenly converge and accommodate, though this is not the rule.

The primary direction of the oscillatory movements is away from the labyrinthine lesion, i. e., if the left labyrinth is involved the direction of oscillation is from left to right, or vice versa, no matter in what direction the eyes are turned; I have seen no exception to this.

The nystagmus and vertigo as above described are in no sense characteristic of the labyrinth; I have seen the same phenomena occur in cerebellar abscess, the location of which was in immediate proximity to its central lobe.

Upon the removal of the semicircular canal system, the vestibule, and the lower half turn of the first cochlear whorl, I have seen nystagmus, vomiting, vertigo, disturbed equilibrium, and intense high-pitched hissing noise, which were distressing before operation, immediately cease.

As to why the destruction of the semicircular canal system and vestibule may in the one instance be productive of vertigo, vomiting, nystagmus and disturbed equilibrium, and in the other cause these same symptoms to quickly disappear, I do not know, but even where these symptoms follow operation upon the labyrinth and are marked, they, as a rule, so quickly subside that it is evident first, that the semicircular canal system is not an indispensable factor to the preservation of equilibrium; and second, that we can scarcely attach to it the importance of being the central organ of the statical sense, as its loss can be so quickly compensated for and in some instances seem to produce no disturbance whatever; it evidently is a part of more or less importance in the great circuit governing equilibrium, but it is by no means an indispensable part, nor is the entire labyrinth indispensable.

Following the removal of the major portion of the labyrinth, conjugate deviation of the eyes may occur, the direction assumed being

upward and outward toward the operated side, as if the eyes were attempting to see the labyrinthine lesion. It should be mentioned, however, that in the two cases in which the phenomenon was observed, cerebro-spinal fluid was lost in both. This suggests that the conjugate deviation may have arisen from other causes than the destruction of the labyrinth. The conjugate deviation was noticed immediately upon the patient's emergence from anaesthesia.

Following the removal of the deep portion of the petrous pyramid, we may notice upon the patient's emergence from anaesthesia that both pupils are dilated ad maximum. This dilation probably results from disturbance of the sympathetic, at the deep, apical portion of the pyramid.

The pupils, though dilated ad maximum, may react to both accommodation and to light. The dilatation gradually disappears after several weeks.

Following the exenteration of the petrous portion of the temporal bone, probing or pressure in the immediate vicinity of the stump of the auditory nerve may produce vertigo, nystagmus, and the subjective phenomenon of intense noise, which noise is referred to in the involved ear. The trunk of the auditory nerve is capable of interpreting mechanical stimuli, as sound, and from the case which I had the opportunity to observe, I am inclined to believe that the trunk of the auditory nerve is also sensitive to waves of sound.

Following the exenteration of the labyrinth, the patient may be exceedingly sensitive to noise and to loud musical sounds. The sense of harmony may be lost and after several months regained. The patient, if subjected shortly after operation to loud musical sounds, as the playing of the piano, may become intensely dizzy, the vertigo attaining to such a degree that it necessitates his sitting that he does not fall.

It would appear that we receive through the auditory apparatus certain physical impressions which, when transmitted to higher centers, act as supporting influences to the preservation of station; that these, when not balanced by similar impressions acting upon the other ear cause, so to speak, an auditory imbalance; that the disturbance is due to the lateralization chiefly to one side of those impressions, which give rise to the subjective phenomenon of sound is questionable, inasmuch as the patient who was totally deaf before operation experienced then no such disturbance under similar conditions.

The phenomenon is suggestive of the great variety of factors which may act through the medium of the auditory apparatus and

quickly disintegrate and completely disappear. The nerve should contribute to the preservation of that condition which we speak of as the state of equilibrium—it gives us an insight into the intricacy of the function of the auditory apparatus.

Following the removal of the entire labyrinth, tinnitus may still persist in the involved ear; its character, however, may completely change, and in all the cases the tendency has been—even where the tinnitus before operation was intense and of high pitch—for it to assume a low, buzzing, inoffensive character after operation. Later the tinnitus may entirely cease.

When the facial nerve is exposed throughout its entire circumference, we may with confidence look for paralysis upon the patient's emergence from anaesthesia. The paralysis deepens and in forty-eight hours becomes complete. It persists from four to five months, even though no injury is done the nerve other than its exposure to the wound and its fluids, and though the reactions of degeneration may appear, the face under massage, and with time, regains its nearly normal expression; I have seen, however, no case in which the function has been so completely restored that the variation in the two sides could not be detected upon the act of smiling. When the nerve is exposed in the manner mentioned, it is difficult at the time of operation to make it respond to ordinary mechanical stimulation, and moderate heat does not affect it.

The facial nerve is structurally very delicate. I have seen a nerve after several days' exposure to an infected cavity which had been irrigated but twice with a 1-10,000 solution of bichlorid of mercury, be protected by sterile vaseline and subjected to no pressure by the packing and the dressing should be changed sufficiently often to keep the cavity clear of pus.

Subsequent to operation, meningitis may occur. It manifests itself in from forty-eight to seventy-two hours. It is characterized by the usual features, with the exception that the sensorium is remarkably clear till near the end; the termination is rapid. Though the postmortem may show a diffuse leptomeningitis, the cerebellar fossa is the site of greatest pathological activity. The route of invasion may be apparent, and in one of my own cases was along the course of the auditory nerve.

When the internal auditory meatus is opened, the rapidity of escape of cerebro-spinal fluid suggests the proximity to the canal of large quantities of cerebro-spinal fluid, as if the cerebellar fossa contained or was in direct communication with a relatively great

amount. In the period of approximately a minute, the cavity represented by the Staëke-Schwartz excavation may fill as many as three times.

Following the too rapid drainage of the cerebrospinal fluid from the cerebellar fossa the following symptoms may appear: a small, rapid, irregular pulse; embarrassed respiration, which may assume a Cheyne-Stokes character; an anxious, concerned expression, with cold extremities and pinched, livid features; marked restlessness.

These symptoms may be of such alarming character that death seems imminent, but with a lessening of the escape of cerebro-spinal fluid the whole picture may in the course of a few hours so completely change that the patient appears quite normal. These symptoms are undoubtedly due to a too rapid drainage of the water-bed of the brain, whereby the important nerve centres at the base, having lost their elastic support, sink and are brought into undue contact with the floor of the skull.

The quick disappearance of these symptoms upon the cessation of the escape of cerebro-spinal fluid forbids any other than a mechanical interpretation.

Following operations upon the labyrinth, large quantities of sugar and acetone may appear in the urine, and the clinical picture of diabetes may rapidly develop. These phenomena suggest that operation upon the labyrinth may be productive of a profound systemic shock which does not manifest itself in the usual disturbances of temperature and pulse. The causative relation, however, between shock and diabetes is highly suggestive of the major character of the labyrinthine operation.

In three cases where the major portion of the entire labyrinth was removed, including the semicircular canal system, there appeared to be after a lapse of over two years in the first, one year in the second, and nine months in the third, no sign whatever of muscular atrophy of either side of the body.

Where the destructive process in the labyrinth has existed for a long time and the irritative stage is passed, the organ, so to speak, is functionally silenced and shows no sign of life; its exenteration is followed by no disturbance whatever of equilibrium; no vomiting, vertigo, or nystagmus occur, as the compensating factors, whatever these may be, have made good the deficiency.

That the disturbance of equilibrium, nystagmus, etc., which follow the removal of portions of the labyrinth are due, to some extent, to the loss of these parts is highly probable, inasmuch as these symp-

toms continue, even though the interior of the labyrinth is thoroughly cocainized. That these symptoms, however, are not altogether due to the loss of these parts is evident, inasmuch as under anaesthesia, mechanical irritation of the interior of the vestibule may produce nystagmus, as may also pressure upon the ampullary areas of the semicircular canals. Pressure upon the stapes may give a similar result. Further, if we irritate with a probe certain portions of the labyrinth subsequent to operation, we may in some instances cause the patient to fall, by producing an intense vertigo. Nystagmus may also be induced or augmented by pressure over certain areas. It is highly probable, therefore, that the disturbances of equilibrium which follow the removal of portions of the labyrinth are due not only to the loss of the parts destroyed but also to peripheral irritation, to which the interior of the labyrinth is subjected. It is only upon this double basis that we can explain certain phenomena in connection with these cases.

For instance, in Case X of the writer's series, disturbed equilibrium, vertigo, vomiting and nystagmus which were intense and distressing before the operation, immediately ceased upon the removal of the semicircular canal system, the vestibule and the lower half of the first cochlear whorl. If, therefore, these disturbances are due to the non-activity of the canal system, and not to irritation, why should the patient when in possession of these parts have exhibited intense disturbances of equilibrium, and after their removal, which insured their non-activity, have shown no disturbances of equilibrium whatever?

Following operation upon the labyrinth, the pulse may be both irregular and rapid and this disturbance may be independent of the loss of cerebrospinal or labyrinthine fluid; the condition soon rights itself.

From the following facts certain conclusions may be drawn, which are of some practical interest in that they show the difficulties attending labyrinthine localization. If we remove the arch of the horizontal semicircular canal, there may follow, vertigo, vomiting, nystagmus and disturbed hearing.

The removal of the entire canal system and the vestibule may be followed by the same phenomena.

The accidental removal of the stapes may produce vertigo, vomiting, nystagmus, disturbed equilibrium and disturbed hearing. The destruction of portions of the vestibule and cochlea without destruction of the canal system produces the same phenomena. It is now

evident that if we deny to these various phenomena, representing several different functions disturbed, the entire labyrinth as the anatomical seat of each (for these phenomena follow no matter what portion of the labyrinth is destroyed) we are led to the following conclusions, which seem to be supported by clinical evidence:

1st. That if the integrity of the labyrinth is encroached upon at any point, the function of hearing, as well as the non-acoustic functions, is seriously interfered with.

(If the encroachment has been gradual and there is no apparent disturbance of equilibrium, etc., it is due to compensation.)

2d. That the disturbances which follow the destruction of a given portion of the labyrinth are not necessarily to be attributed to the loss of the parts destroyed.

3d. That the destruction of the labyrinth at a given point affects the function of distant parts as well. If, for instance, we destroy the entire labyrinth with the exception of the cochlea, or, if we merely remove the arch of the horizontal semicircular canal, we are at a loss to know whether to attribute many of the disturbances which follow to the loss of the parts destroyed or to the irritation of the parts remaining.

It therefore follows that an attempt at labyrinthine localization based upon symptoms will always be subject to the grossest error, and that the attempt to determine the functions of the various departments of the labyrinth by the experimental destruction of these will be subject to the same error.

The difficulty of experimentally determining with accuracy the functions of the various parts of the labyrinth with a view to labyrinthine localization, and this is here alluded to merely on account of its surgical interest, is due in large measure to the anatomical arrangement of the labyrinth, the essential feature of which is, that it is suspended in a fluid medium which is *continuous*. A disturbance of the medium at any point is transmitted throughout the entire apparatus and results in the disturbance of distant parts.

While this particular anatomical arrangement complicates the study of the physiology of the labyrinth, it is an essential attribute of an organ which embraces several distinct and separate functions, which are correlated to the extent that each is influenced through the same common medium and to some extent by the same primary cause—waves of sound.

The labyrinth is an example which illustrates the conservation of energy in the evolution of an organ in that the anatomical bases of its

several functions, which in reality represent different organs, should have been so grouped together and suspended in a common medium as to make common use of that medium. This arrangement is evidence of the influence which waves of sound exert upon the non-acoustic as well as upon the acoustic labyrinth and a practical example of this is illustrated by the case previously mentioned where the patient when subjected to loud musical sounds, after the extirpation of the entire labyrinth, was made so dizzy as to necessitate his sitting, his equilibrium having been profoundly disturbed.

While it is the purpose of the present paper to treat of only those conditions which have a surgical bearing, there are many facts in connection with the cases which are of interest in that they are in contradiction to certain theories which have gained a more or less widespread acceptance.

44 N. 49th St.

A Case of Nasal Hydrorrhoea Cured by Electrolysis. DR. VALENTIN. *Le Nord med.*, Jan. 15, 1906, No. 271.

Valentin reports a case of nasal hydrorrhoea in a patient at the menopause. Spraying with cocaine and stovine was without effect as well as constitutional methods such as strychnine and atropine. The application of electrolysis, however, was followed by a complete cure.

SCHEPPEGRELL.

Massage of the Stapes. TREITEL. *Arch. f. Ohrenh.*, Leipzig, Oct., 1904.

In two cases, massage of the stapes with the Lucae's pressure probe resulted in remarkable improvement of the hearing, lasting several years.

YANKAUER.

FIG. II

FIG. IV

FIG. V

FIG. VI.

FIG. VII

FIG. XII