paired in all muscles; when one of his arms is passively raised (his eyes being closed) he cannot find it with his other hand; thinks his mouth closed, after it was continuously opened, while he had his eyes shut. There is also optic, auditory, olfactory and gustatory hypæsthesia. Looking into the light is painful and causes increased lachrymal secretion. Faradic excitability of muscles and nerves normal, galvanic qualitatively normal, quantitatively somewhat increased.

Bechterew considers the hyperæsthesia of the muscles, bones and articulations to irritation and insults of all kinds to be the most characteristic symptom of the disease. The latter is not a psychosis. In atremia, which in some respects resembles akinesia algæra, tenderness of the muscles, bones and articulations to mechanical insults is not present.

ONUF.

Amusia: Musical Aphasia.—Dr. T. G. Edren, Stockholm (*Deutsche Zeitschrift. für Nervenheilkunde*, 6th Vol., I. and II. Parts).

The author reports a case of transient paraphasia and word-deafness and permanent deafness for musical notes (Tontaubheit). At the post-mortem examination lesions were found in the anterior two-thirds of the first temporal, and anterior half of the second temporal lobe of the left hemisphere. Fifty-one other cases, bearing on that subject, scattered mostly in French and German literature, furnish the basis for the conclusions of the author.

The material is arranged in three categories as follows:

1. Aphasia, without amusia, twenty-four cases; four post-mortem examinations.
2. Aphasia, with amusia, twenty-two cases; five post-mortem examinations.
3. Amusia without aphasia, six cases; two post-mortem examinations.

Altogether fifty-two cases, eleven post mortem examinations.

The author believes that the data contained in the mentioned clinical and anatomical material justify the following conclusions:

1. Any pathological process within the skull destroys the musical faculties of the patient in the same way as the speech faculties, thus leading to different clinical forms of amusia.

2. The different forms of amusia have some clinical independence in relation to each other, as well as to the different corresponding forms of aphasia.

3. The clinical forms of amusia show a great analogy with the clinical forms of aphasia. The former are often, not always, accompanied by the corresponding forms of the latter.

4. Amusia may appear in the clinical picture without aphasia, and *vice versa*.

5. Some of the forms of amusia have very probably an anatomical independence, and are always localized in the neighborhood of the areas of the corresponding forms of aphasia.

6. Musical deafness (Tontaubheit), *i. e.*, the perception of notes, accords and melodies, without comprehending their musical meaning, is probably located in the first or first and second left temporal lobe, anteriorly to the area of verbal deafness.

FRANKEL.

Astasia-Abasia and Its Treatment.—By F. Friedlander, M.D. (*Neurolog. Centralbl.*, 1894, p. 354).

Friedlander finds that in astasia-abasia there is not a loss of the conception of movements, but an inhibition of association in the transmission of the conception of movements into voluntary motion. This inhibition is produced by auto-suggestion on the foundation of abnormally exaggerated suggestibility, which latter Friedlander finds to be the most marked characteristic of every hysterical disease. Friedlander thinks, accordingly, that the treatment ought to be suggestive, but is against