great frequency the latter policy may be questioned, for we may be leaving behind a portion of a malignant growth. I judge, however, from his paper that this transformation is not frequent, and therefore, especially in young women, the fear of its recurrence should not materially influence us in discarding the conservative policy. While some distinguished surgeons are inclined to deny the possibility of auto-implantation, I am convinced that it is possible to carry cancer cells from the field of operation into healthy tissue where cancer will recur. I wish to ask Dr. Cullen how often these malignant transformations of fibroids occur and how far his observations justify the physician in favoring the early removal of these clinically benign tumors with a view to preventing the complication.

DR. L. H. DUNNING, Indianapolis—I would like to have Dr. Cullen consider the influence of these findings on our technic in myomectomy. I also wish to ask whether or not he has collected statistics on the recurrence of these growths after complete and after partial removal of the uterus.

DR. C. O. THIENHAUS, Milwaukee—Does Dr. Cullen consider the presence of giant cells and myoclasts in fibroids a sufficient proof of malignancy? Saenger stated that all fibroids containing giant cells and myoclasts had to be classified as showing sarcomatous degeneration. Concerning Dr. Dunning's question, Martin found sarcomatous degeneration 6 times in 205 cases of fibroids, but this percentage is an exception and extremely high, in my opinion. Landau collected from the literature 10 cases in which following retroperitoneal hysteromyomectomy later on a malignant growth developed in the cervix.

DR. C. L. BONIFIELD, Cincinnati—About eighteen months ago a patient came under my care who had had an ectopic gestation that went to full term and who had had a missed labor fourteen years before I saw her. She developed an immense fibroid, larger than my head, which was undergoing sarcomatous degeneration at the time of the operation. I removed the uterus and the tumor and the ectopic gestation, but unfortunately she died a few weeks afterward from a recurrence of the sarcoma.

DR. T. S. CULLEN—It is impossible for me at this moment to tell the exact percentage of myoma cases in which sarcomatous degeneration takes place, but from 1.5 to 2 per cent. show changes definitely sarcomatous in character or strongly suggestive of it. Carcinoma of the body of the uterus is relatively common where uterine myomata exist. Taking all myoma cases we have at least 2 per cent. in which a malignant growth of some character is present. In hysteromyomectomy in favorable cases where there are not many adhesions the mortality will not be higher than this. I fully agree with those who advocate early operation in nearly all cases. Operate before adhesions have formed. I do not wish to deprecate bisection of the uterus. It is a most admirable operation where the uterus is wedged in the pelvis, thus preventing us from reaching the uterine vessels. If we bisect the organ the myoma can be rolled out, more space is thus obtained and the vessels are more easily controlled. But in these cases we must remember the possibility of sarcomatous degeneration of the myoma. We must, however, run the risk of cutting into the sarcomatous myoma as in some of the cases the myoma is so wedged in the pelvis that its removal would be impossible by any other procedure. Recurrences in cases where sarcomatous degeneration has taken place are certainly appalling. In some of the cases where bisection of the uterus was done the patients died of infection in a few days. I have a record of one case where within six weeks after hysterectomy the pelvis was filled with a sarcomatous growth. In one of our cases where temporary recovery took place Dr. Osler saw her six months afterward and there were metastases in one lung and in the mediastinal glands. She soon died. From the histologic standpoint it is most interesting to follow the sarcomatous changes in myomata, but we sometimes have specimens where it is impossible to say absolutely whether a given area is really sarcomatous or merely a suspicious but benign growth. One can not interpret the changes in the muscle cells with the same degree of precision as is possible when considering cell

changes in the endometrium. From a systematic examination of the uterine mucosa we have found in nearly all cases where the tubes and ovaries are normal that it is unaltered. The changes that do occur in the endometrium are dependent almost entirely on the situation of the growth. If it be partially submucous the mucous membrane over its most prominent part will be thinned out more and more until eventually it is represented by nothing but a layer of epithelium. This, after a time, is flattened and disappears. In the depressions between the myomata there is often slight gland hypertrophy or polypi develop. Knowing that the endometrial changes are almost entirely mechanical in nature where normal tubes and ovaries exist, we can open up the uterine cavity in myomectomy cases when necessary, there being little danger of infection. On the other hand, it is rarely, if ever, advisable to do a myomectomy where the appendages are adherent. Several years ago I operated on a patient suffering from adherent tubes and a myoma in the posterior uterine wall. There was no evidence whatever of any recent infection. I made artificial extremities for both tubes and then did a myomectomy. Shortly afterward the patient's temperature rose and she developed definite signs of sepsis. Although we reopened the abdomen and drained thoroughly she died of sepsis a month afterward. In this case some old infection evidently lurked in the tubes and readily extended to the uterine walls at the site of the myoma enucleation. I would strongly advise against myomectomy in any case where the appendages are adherent.

A CASE OF ACUTE DISSEMINATED ENCEPHALOMYELITIS,

PRESENTING THE CLINICAL PICTURE OF MULTIPLE SCLEROSIS.

JOSEPH L. MILLER, M.D.
CHICAGO.

Acute disseminated encephalomyelitis manifests itself in one of two clinical types, viz.: As an acute ataxia or acute paraplegia. The ataxia type is often described as acute ataxia, or better, acute central ataxia, the latter term serving to differentiate it from the ataxia resulting from peripheral nerve changes.

Eisenman in 1863 described a case of post-typhoid ataxia as a result of myelitis. Leyden in 1868 reported a case following trauma. Westphal in 1872 reported several cases following variola and typhoid, and was the first to verify his diagnosis by autopsy. Ebstein at the same time reported a case of speech and co-ordination disturbance following typhoid, and eight years later reported the autopsy findings on the same case. Pollard in 1872, under the title, "Case of Locomotor Ataxia, Beginning Suddenly and Disappearing Under Treatment," described what was probably a case of myelitis. During the past thirty years numerous cases have been reported which, with the pathologic investigation, have verified the findings of the earlier writers.

NARRATION OF CASE.

This patient entered Cook County Hospital in July, 1902, in the service of Dr. Dolamore, through whose courtesy I am permitted to report the case.

History.—Patient was 47 years old, laborer. Family history was negative. He had been a moderate drinker during the past ten years; previous to that he drank excessively; he chews and smokes tobacco. He had syphilis twenty years ago and was treated at irregular intervals for several years. Six years ago, as the result of a trauma, he sustained a fracture of the right femur. Three years ago the patient was in Cook County Hospital with croupous pneumonia; during the course of the disease he suddenly developed an aphasia, which was only complete for a few days, rapidly improving. Whether there was an involvement of the legs at the same time the pa-

tient does not know and no record of it was made at the time. A few days later the patient made his first attempt to walk, and found that he staggered like a drunken man. Slight disturbance of speech has persisted since, particularly manifested in a slowness and monotone with inability to make himself understood when excited. The difficulty in walking varies with periods of marked improvement followed by relapses. There has never been any bladder or rectal symptoms. Sexual power is lessened, but eyesight is unimpaired. There is no gastric disturbance. His memory is good. He is more emotional during the past three years.

Examination.—The patient is well nourished and his color is good. The feet are widely separated in walking; toes abducted, heels striking the floor first. Muscular strength in all groups of muscles of both upper and lower extremities is good. There is no marked atrophy. Touch, pain and temperature sense are normal. There is marked swaying when the patient stands with his feet together, which is increased by his closing his eyes. There is no ataxia of the legs when lying down and very slight inco-ordination of the arms. There is slight tremor of the hands when at rest, increased by intended movement; very marked intention tremor in legs; none of head. The pupils are equal, reacting to light and accommodation, with marked horizontal nystagmus, no optic atrophy. Skin reflexes are present; knee jerk is slightly under normal. The patient speaks slowly and in a monotone, no distinct scanning. The hearing in the left ear is slightly impaired.

ETIOLOGY.

The vast majority of the cases of disseminated myelitis develop during or following the acute infectious diseases, especially variola and typhoid, less frequently whooping cough, erysipelas, dysentery, measles, influenza, malaria, tuberculosis and gonorrhea. Several cases are reported following exposure to wet, cold or severe fatigue. Leyden reports a case two weeks after fracture of the arm, Taylor after injury received in alighting from a train. A definite etiologic factor was not apparent in many cases.

Experimentally, Babinski has produced both transverse and disseminated myelitis by injection of *Bacillus pyocyaneus*, Roux and Yersin with *B. diphtheriae*, Rogers with streptococcus. Eisenlohr found staphylococci in the cord in a case following typhoid. Banir found staphylococci following gonorrhea.

PATHOLOGY.

Westphal examined two cases of the paraplegic type, and found multiple areas of necrosis in the cord. Ebstein's case showed gray foci in the cord and medulla. Examinations of recent cases show perivascular inflammation in the cord, medulla, pons and cerebrum—in the latter in both gray and white matter. The nerve fibers are destroyed, the neuroglia increased.

SYMPTOMOLOGY.

Clinically, the disease manifests itself as an acute ataxia or paraplegia. The characteristic symptoms of the ataxia form given by Westphal were: 1, Slow or scanning speech; 2, ataxia with normal or slightly lessened muscular strength; 3, no sensory disturbances; 4, often psychic disturbances.

The ataxia is often more marked in the upper extremities, or it may be present in the form of a hemi-ataxia. Ebstein's patient could move his legs in bed without difficulty, but was unable to walk. Unconsciousness may mark the onset, the ataxia only appearing later as in Lenhartz's case; anesthesia and sphincter paralysis marked the onset, the ataxia appearing two weeks later. Intention tremor of the head and extremities is usually present. Nystagmus is an infrequent symptom. Speech disturbances are almost always present, varying from

complete aphasia to slight slowing or monotone. Slight if any superficial or deep sensory disturbances are present. In Lenhartz's case the anesthesia present at the onset disappeared in two weeks. Sphincters were, as a rule, intact. Ptosis was present in Pollard's case. Optic neuritis has been occasionally noted. Skin reflexes usually unchanged. Tendon reflexes often exaggerated.

The paraplegic type is characterized by a flaccid paralysis, with loss of knee jerk, sphincter involvement and often sensory and trophic disturbances.

The course is: 1, Early complete recovery; 2, partial recovery, with perhaps only one symptom remaining, as the speech disturbance; 3, periods of improvement with relapses, the condition being progressive. Ebstein's case showed slight improvement in the beginning, followed by gradual progression, the ataxia reaching such a grade that the patient was confined to his bed during the last two years. Westphal believed that many cases reported as recovered would later relapse. Early fatal termination is unusual.

RELATION TO MULTIPLE SCLEROSIS.

The clinical picture in many of these cases, including the writer's, is one of multiple sclerosis. Leyden formerly maintained that the absence of progression differentiated it from this condition; this has not, however been supported by clinical facts, as in Ebstein's case quoted above. Instead of an acute process with healing and permanent defects, the acute onset would appear to mark the beginning of a chronic process, which may become latent, but more frequently tends to advance. Pathologists are agreed that the anatomic findings may be identical with those of multiple sclerosis. The clinical symptoms complete the picture, and it would be preferable, as is insisted on by many writers, to consider these cases of multiple sclerosis with acute onset.

TREATMENT.

In the early stages, rest in bed, with special attention to the skin to prevent bed sores, is essential. Cold may be applied along the spine, without, however, much hope of benefit. Later, the use of massage and electricity is of value. The systematic exercises employed in tabes are of value in lessening the ataxia. Freedom from exposure to cold and over-fatigue may prevent acute exacerbation. The long list of drugs which are reported as curative is evidence that spontaneous recovery, or at least improvement, is not unusual.

100 State Street.

DRAINAGE, SEWERAGE AND WATER SUPPLY OF NEW ORLEANS.*

GEORGE G. EARL.

General Superintendent of the Sewage and Water Board.

NEW ORLEANS.

In undertaking to give you a brief account of what has been done, or is contemplated, in the direction of the draining and sewerage of New Orleans, as well as in the matter of a water supply for her 300,000 thirsty citizens, I understand that I am undertaking a task which will require a most direct method of treatment, to be kept within bounds as to time, and will, therefore, without circumlocution or delay plunge at once into my subject.

* Read at the Fifty-fourth Annual Session of the American Medical Association, in the Section on Hygiene and Sanitary Science, and approved for publication by the Executive Committee: Drs. S. Fulton, Seneca Egbert and H. M. Bracken.