hose with great force, and the stream was directed into the mouth, cleansing it most thoroughly and giving a delightful sense of comfort.

For the purpose of injecting a paste into pockets or between the teeth, the writer has also devised an ointment syringe, the piston of which is driven by means of a screw. (Fig. 2.)

And now a word in relation to all that has been written regarding oral hygiene in the public schools. The dental examination of pupils is of undoubted value, but they seem to be regarded merely as material to be turned over to the dentist by the board of overseers.

However destitute parents may be, they have the rights of individuals, as citizens, as fathers and mothers, which should be respected, and their consent, not the pupil's, should be asked before a dentist, not of their choosing, shall insert instruments or fingers into their children's mouths in the course of the examination of a hundred others. The father of small means has the right to ask that the same or equivalent antiseptic precautions be used for his children as for rich private patients, and his ignorance of these matters should be no reason for depriving him of the privileges belonging to any citizen.

Although many of the before-mentioned subjects have been treated briefly and perhaps inadequately, your essayist now comes to a stop because "a wise man is like a spring lock, always more ready to shut than to open."

31 Beacon Street.

DISCUSSION.

DR. VIDA A. LATHAM, Chicago—I think all of us are familiar with Dr. William Hunter's paper on "Oral Sepsis," a paper which has done more, I believe, toward uniting the medical and dental professions than any other paper ever written in the history of medicine. It is quoted all over the world, showing the great importance of noting oral sepsis. I would also like to call attention to a condition in that curious disease known as pernicious anemia. There we get a most peculiar glossitis, stomatitis, a condition that is so characteristic as to be almost one of the diagnostic points in pernicious anemia. We can also get from the lymphatics of the tongue an almost pure culture of the *Streptococcus longus*.

DR. E. S. TALBOT, Chicago—The essayist says that a perfectly healthy mouth will not emit odor and speaks of decomposition of food about the teeth producing foul odors, etc. I do not quite agree with the statements. The amount of food which collects about the roots of the teeth is very small. Again, the mouth may be healthy but there may be odor from autointoxication due to indigestion or to the non-elimination of effete matter. In cases of interstitial gingivitis or alveolar pyorrhea the worst forms of odor are found. In a great many teeth which I have cracked the odor was frightful and yet the pulps were still alive. In cases of extreme absorption of the alveolar process with exposed tubuli the odor is also bad.

In the treatment I must take exception to the use of a soft or wet brush. My experience is, that there is not a brush made that is sufficiently stiff. In order to have a healthy mouth gums must be massaged with a stiff brush. I am glad to hear the protest regarding the examination of the mouths of school children. The habit of examining the mouths of these children without thorough cleansing of the fingers is abominable.

DR. M. L. RHEIN, New York City—Dr. Eames speaks of three minutes being advised for brushing the teeth. I insist that my patient shall give from four to five minutes to the brushing. I think the average patient gives about 30 seconds to this important matter, and I am accustomed to laying down a law that must be rigorously observed. I agree with Dr. Talbot that the bristles of a brush can not be too hard. I instruct my patients to use the flat sides on the gums as near the roots as possible in order to stimulate the capillaries of the gum tissue. Recession of the gums or necrosis of the tissue can be greatly retarded if the tone of the tissue is kept up by

proper massage. This point I especially emphasize in my instructions to assistants, as to the necessary directions to be given patient in oral hygiene.

DR. G. V. I. BROWN, Milwaukee—We all have our own ways of accomplishing the same results. The broader view of the paper is its significance in relation to the prophylactic feature. I want to correct what may appear to be in print, as I am sure it was not intended to be—a protest against the examination of school children. Very little has been developed out of these examinations, yet there is a work in this direction which ought to be looked after. I think many of the ills of young persons' future lives could be avoided by proper attention to this examination. Matters pertaining to nose and mouth breathing and the general disinfection of the mouth are of the greatest importance. Much could be done in checking epidemics of diphtheria and the prevalence of tuberculosis by insisting that a large vessel of a cheap, suitable disinfectant be placed at the door of the school-room and the children made to take a mouthful of the solution and hold it in their mouths for a few minutes. Whatever may be done with the sputum, by legislative ordinances against spitting, the use of spit-cups, etc., it is obviously true that the destruction of bacilli before they get into the air would be, if practicable, a better and the only true prophylaxis.

DR. A. E. BALDWIN, Chicago—I believe practical results are to be obtained by anything that will properly cleanse the system and promote a healthy condition of the mouth. It seems to me hard to determine whether odors come from within or beyond the mouth. I think the fears regarding tuberculosis are carried to the extreme. I thoroughly believe that tuberculosis is a great scourge and that all possible should be done to prevent it, but when it is realized that during the life of the human race it is estimated by careful investigators that fully 70 per cent. of the people who attain adult life have sometime during their life tubercular trouble and get well, the disease does not seem so hopeless as might at first be considered. I think care in the disposition of the sputa will bring the best results in preventing the spread of the disease. I am a thorough believer in cleanliness in any way it can be accomplished.

DR. EAMES, in closing—In regard to the healthy mouth not emitting a foul odor, I would admit that the mouth may be the outlet of an odor that may come from a distant part, but if that part is diseased, it is a question in my mind whether the mouth can still remain in a perfectly healthy condition, the oral cavity being so readily affected by abnormal conditions in other parts of the body.

The brush I leave to the discretion of the dentist. The idea was to make a distinction between the wet and dry brush. I believe we can cleanse the teeth better with the wet brush if the bristles are sufficiently stiff and plenty of powder is used.

ETIOLOGY AND TREATMENT OF MIGRAINE.*

J. M. AIKIN, M.D.

Lecturer and Clinical Instructor in the Medical Department,
Nebraska State University.

OMAHA, NEB.

Of all the common and much-dreaded nervous diseases we recognize, none are less perfectly understood than migraine; nor is there any other nervous disorder which is so disastrous to the physician's ability for effectual treatment. Under the title I wish to be understood as excluding neuralgia, neuritis, headaches due, proximately, to improper alimentation, nervous exhaustion or brain fatigue. I would exclude the head pains as not migraine that are cured by merely correcting ocular or nasal abnormalities.

The causes for headache seen through the speculum of the gynecologist and proctologist are not a few; and the many radical cures effected by their rational treatment are glowing tributes to medical progress; but

* Read before the Nebraska State Medical Society, Omaha, May 6 to 8, 1902.

these headaches are not true migraine. The uric acid diathesis is a step in the right direction, but even Haig, its foremost advocate, has not reported his own case of migraine as cured by using alkaliphia. It is easy to say what it is not, but difficult to define what it is.

HEREDITY AS A FACTOR.

By far the most frequent and constant individual etiologic factor in migraine is heredity. Clinical facts teach that transmitted conditions continue throughout many generations and render easily possible the evolution of migraine under suitable conditions. Just what these degenerative stigmata are we do not know, but there is a preponderance of evidence locating the disorder in the nervous system. I believe it has a pathologic entity susceptible to and having an affinity for certain toxic elements in the circulatory system. The nerve cells, bathed in the life current, hold out against these ptomaines until their cumulative strength, overcoming the normal resisting powers of the individual, culminate in an explosion. These nerve storms are denominated migraine. They appear periodically, their frequency being determined by the vulnerability of the tissues, intensity of the ptomaines and contributory influences known as reflex causes.

The transmissibility of conditions favoring migraine is evidenced by its appearance in early childhood and its unabating continuance till after the middle third of life.

THE RESULTS OF TOXINS.

As a result of tissue metabolism, toxins are a constant element in the system common to all. These become an irritant only under conditions of improper elimination. Toxicity of the ptomaines is one of quantity rather than quality. Contributory causes arise from our indulgence of the palate, and the improper functional use of any part of the physical organism. The sufferer from migraine anticipates the onset and very often designates the particular indulgence or incident that precipitated the attack.

For a short time after the nerve storm has passed, the system is comparatively immune, for the same causes that operated to precipitate the attack are indulged with impunity until the system is again surfeited with the ptomaines.

The interval between paroxysms varies with each person, and the same person may increase the frequency of attacks by the improper exercise of one or many individual privileges. The person who is normally accustomed to a migrainous seizure once in four weeks may, by dissipation, mental strain, worry, or loss of sleep, have an attack more frequently.

THE RELATION TO EPILEPSY.

The etiology of migraine is quite as obvious, and the transmissibility of conditions favoring its development are more constant than the causes and conditions producing epilepsy, which we all recognize as closely related to migraine. It is not uncommon to see migraine and epilepsy transposed in the same individual, and often in members of the same family. The interchangeability of these two disorders, indicated to us by the masters in medicine, does not obliterate certain individual characteristics in which they are wholly different. Possibly the attacking forces are the same in each; but meeting different conditions, there results in one a sensory and in the other a motor psychosis.

OPINIONS OF OTHER WRITERS.

Alexander Wallace¹ says: "Migraine is believed to be

due to defective or insufficient excretion, partly of the liver, but mainly of the kidneys. The severity of the headache is directly proportionate to the acidity of the gastric fluids."

W. Heind² says: "Migraine is attributed to two principal causes, (1) poisoning of gastric origin; (2) cerebral fatigue."

"Migraine is regarded as a toxemic condition, the toxins (probably albumoses) being absorbed from the gastro-intestinal canal."—C. E. Herter.³

"The occurrence in migraine of almost constant vomiting, frequent attacks of diarrhea, and polyuria or increase in other secretions is proof that many cases of this affection depend upon intoxication."—W. Stekel.⁴

These conclusions from physicians in widely separated parts of the globe show a growing tendency toward recognizing the germ theory of disease.

Intelligent thought has isolated and classified many bacteria that originate external to the body, which, when introduced, develop specific diseases. Less has been said, however, about those originating within the human economy becoming disease producers when allowed to accumulate beyond certain proportions.

THE COMMON TREATMENT.

In the treatment of migraine, almost as many drugs have been used as equal the number of physicians who have treated the disorder. The principal drugs used are in the main motor depressants or cardiac stimulants. These have been used separately and in combination, often without any very definite therapeutic purpose, hoping only for relief from immediate conditions.

When all else fails morphin will quiet the irritated sensory nerves. Attacks are sometimes aborted, or their intensity lessened, by vigorous acceleration of the blood current during the prodromes. The good thus accomplished doubtless comes from the more frequent contact of venous blood with radiating tissues.

TREATMENT FAVORED BY THE AUTHOR.

Seeing good results from such treatment led me to believe that diuretics, diaphoretics and hydragogues would materially assist in the eliminative process. We have many well-known drugs that act promptly through these several avenues, but in each case their efficiency is greatly enhanced by water as a diluent.

From the inception to the conclusion of a migrainous attack, digestion is practically suspended. This condition certainly demands the withholding of nutrients with immediate dilution and elimination of the gastric and intestinal contents. Emesis and lavage of the stomach are efficient, but often objectionable to the patient and inconvenient to the physician. Consent is much more readily obtained for emptying the lower bowel with a soapsuds enema, followed immediately by high irrigation with large quantities of hot normal salt solution.

This, with small but oft-repeated draughts of hot water by the mouth, continued from six to twelve hours, has given better results in my treatment of migraine than any purely drug medication.

Between the attacks, daily and copious drinking of water will do more to lessen the severity, if not prevent recurring paroxysms, than any or all drugs with only a minimum amount of water ingested.

It would be quite irrational were we to neglect correcting any existing ocular, aural, nasal, gynecologic or

2. Provincial Medical Journal, Oct. 2, 1894.

3. Journal of Nervous and Mental Diseases, January, 1897.

4. Wiener Med. Woch., Nov. 13, 1897.

1. The Lancet, Jan. 14, 1893.

rectal defect. We must also teach them how to eat and warn them against abuse of their already defective nervous system.

A SUBSEQUENT REPORT ON A CASE OF CARCINOMA, DISCHARGED AS HOPELESS AND REPORTED AS A FAILURE.

WILLIAM ALLEN PUSEY, A.M., M.D.
CHICAGO.

In a report of cases treated with Roentgen rays, published in *THE JOURNAL*, April 12, 1902, I recorded a case as follows:

CASE 24.—Man, aged 70, referred to me by Dr. William H. Wilder, with extensive carcinoma involving all the structures in and around the orbit of the right eye. The growth had progressed within the cranium when the treatment was begun. The treatment in this case continued two months, with decided amelioration in his condition for awhile, both as regards the appearance of the growth and his pain, but later evidences of rapid intracranial growth appeared and the treatments were discontinued.

Looking back at this case, it is Dr. Wilder's opinion and mine that we should have taken out enough of the bone of the orbit to have laid bare the structure beneath, so that the *x*-rays could have been thrown directly on the intracranial growth. Dr. Wilder suggested to me that had this been done, there might have been a chance of controlling the disease, so marked was the effect on the diseased tissue which was directly exposed.

When this case came under observation the eye was destroyed, the orbit filled with carcinomatous tissue, the orbital bones involved and there was every reason to believe that the disease had extended within the cranium. The clinical diagnosis of carcinoma had been confirmed microscopically by Dr. Wilder. Treatment was undertaken with *x*-rays with the feeling that there was only the remotest fighting chance.

During the first month that the patient was under treatment the acute pain from which he had been suffering disappeared and the tumor shrunk considerably. During the second month of treatment, however, in spite of the fact that *x*-ray effects were carried to the point of producing an acute *x*-ray irritation, the tumor increased and excruciating localized headache appeared. From the character of the headache there seemed no doubt that the tumor was developing rapidly in the meninges and, as the disease was progressing even on the surface where the *x*-ray effect was greatest and was apparently being uninfluenced by the treatment, we concluded that the case was certainly beyond control and that further treatment was useless. November 23 he was accordingly discharged, having at that time an acute *x*-ray inflammation over the whole involved area. There seemed every reason to believe that he would die quickly. On April 30, however, five months after treatment was discontinued, I received the following letter from Dr. T. Sprague of Sheffield, Ill., to whose thoughtfulness I am indebted for the ability to make this report:

DEAR DOCTOR: I write you regarding the condition of Mr. —, an old gentleman whom you treated for carcinoma of the right eye, discharged as incurable (if I am correctly informed by Mr. —) about the last of November. He was a charity patient and I think was referred to you by Dr. Wilder. He called on me on his return from Chicago, had a recipe calling for tablets of a quarter of a grain of morphin, to be taken when in pain. I saw that he had about twenty tablets. I noticed the condition of his eye at that time and concluded from my examination that it was only a question of time, and short time at that, with him. I saw him yesterday and he presents an entirely different appearance. The mass that occupied the orbit has shrunken, allowing the lids to close. He

is free from pain, eats and sleeps well, uses no anodyne. There are a few little nodules on the nose near the commissure of the eyelids. With that exception he looks quite well. He has certainly gained since coming home, and I feel assured it is the result of the *x*-ray treatment. I should like you to see him again, but, of course, his means being limited, he would have to go to the Eye and Ear Infirmary as before.

This case is instructive in at least two particulars. It illustrates very strongly, in the first place, the persistence of *x*-ray effects. Here is a case in which the effects produced by the *x*-rays persisted for months after discontinuance of treatment and were sufficient to cause the disappearance of a tumor and at least the symptomatic relief of a case, which upon the basis of all experience could end only in death within a short time. We have been made familiar by unhappy experience with the progressive and long-continued effect of *x*-rays in *x*-ray injuries; it is something of a recompense to find these same qualities serving us in the therapeutic use of the agent. The next fact is the evidence in this case of the deep-seated effect of *x*-rays. That this tumor involved not only the orbit and orbital bones, but the surrounding intracranial tissues as well there is almost no room to doubt. The result, therefore, gives unmistakable evidence of the deep-seated action of the rays. Such a case offers another lesson, and that is the agreeable surprise from the use of *x*-rays that may confront one in even the most desperate case of circumscribed carcinoma.

GENERAL NERVOUS MANIFESTATIONS IN RELATION TO THE JAWS AND TEETH.*

GEORGE V. I. BROWN, A.B., D.D.S., M.D., C.M.
MILWAUKEE.

In a paper read before the Section on Neurology and Medical Jurisprudence of this Association in 1898, and this, the Section of Stomatology in 1897, I outlined the theoretical propositions upon which were based methods of treatment adopted by myself in the care of nervous affections intimately related to disturbance caused by the habit of grinding and clenching the teeth, many general as well as local manifestations being attributed to this etiologic factor. The term "jaw-strain" was used in the same sense as "eye-strain."

Attention was called to the fact that wherever the natural teeth remain in the mouths of patients suffering from nervous disorders, it is noticeable that the occlusal surfaces show abrasion, due to constant grinding and rubbing, the effect of extreme pressure brought to bear during paroxysmal muscular effort, or long-continued, excessive pressure, a condition frequently resulting from hours of painful suffering. This has frequent mention by writers as a marked symptom among those who are victims to the various neuroses, but in no instance do I find where an author has thought fit to reverse the order of things and make the habit of the jaws, which is responsible for this condition, one of the etiologic factors in bringing about the disease, rather than a result, as it is generally held to be. This hyperkinetic condition of the muscles of mastication is undoubtedly due to irritation of the brain centers governing these muscles, whether as a symptom of other neural disturbance, or vice versa.

Since the papers referred to were written an almost continuous observation of such cases, and the practical application of these principles, have convinced me more and more that the importance of the subject is but little

* Read at the Fifty-third Annual Meeting of the American Medical Association, in the Section on Stomatology, and approved for publication by the Executive Committee: Drs. G. V. I. Brown, M. H. Fletcher and R. R. Andrews.