THREE CASES OF TUBERCULAR DISEASE OF THE LOWER END OF THE FEMUR, ILLUS- TRATING SOME POINTS IN PATHOLOGY AND TREATMENT.

By W. I. DE COURCY WHEELER, M.D., F.R.C.S.I. ;
Surgeon, Mercer's Hospital.

[Read in the Section of Surgery, January 10, 1913.]

RECENT publications on the subject of tubercular disease of the bones and joints in children, more particularly in relation to the work of Stiles, have thrown light on a subject which heretofore has been in a mire of confusion. Little knowledge could be derived from the involved descriptions, complicated nomenclature, and inaccurate pathology which marred the descriptions in the majority of text-books dealing with osseous tuberculosis.

In this communication I propose to concentrate attention on three recent cases of tubercular disease of the juxta-epiphyseal (metaphysis) portion of the lower end of the femur. The disease commences and progresses along definite anatomical and pathological lines in a corresponding position in all bones.

Pathology.—It may be stated as an axiom that the great majority of cases of osseous tuberculosis owe their origin to infection from contaminated milk, and until this source of infection is recognised and dealt with by the public there will be no diminution of the stream of children suffering from the disease which seeks admission to the hospitals. In a typical case the tubercle bacilli can be traced from the

138 *Tubercular Disease of Lower End of Femur.*

milk to the metaphysis of the bones. Possibly the child suffers from adenoids or enlarged tonsils, the cervical, bronchial, or mesenteric glands become involved. The bacilli are, in the first case, carried through tributaries of the jugular vein into the pulmonary circulation.

Reaching the systemic circulation, the survivors of phagocytosis and endothelial activity are deposited in the metaphysis of bones, to remain latent or to set up active disease. The reason for the lodgment of tubercular emboli in this situation is an anatomical one.

Three sets of capillaries anastomose in the metaphysis of bones :—

1. The end branches of the nutrient artery (diaphyseal group).

2. Branches from the arteries supplying the joint entering the metaphysis (metaphyseal group).

3. Branches from the same source as 2 supplying the epiphysis (epiphyseal group) perforate the cartilage and anastomose with the diaphyseal and metaphyseal vessels.

The circulation through the cancellous spaces of the bone is slow, and this fact, combined with the free capillary anastomosis, makes it obvious why the site is a favourable one for the deposit of a tuberculous embolus.

A slight injury, breaking down some of the trabeculae of the cancellous spaces, may determine the onset (by setting free the embolus) of active tuberculosis.

Once established, the disease is at first frequently central, and subsequently is carried by the Haversian systems through the compact bone to the periosteum. An osteoplastic periostitis ensues, and a thin layer of new bone, very porous and delicate, is laid down between the thickened periosteum and compact bone. Finally, as caseation and liquefaction proceed, an abscess forms in the

soft tissues, and is followed by a sinus leading down to diseased bone.

The formation of the thin layer of porous bone, coupled with the presence of thickened periosteum, is responsible for the very important clinical sign seen early in these cases—viz., thickening. (Plate I., Figs. 1 and 2). Tubercular disease may arise in the epiphysis or in the diaphysis, but for the reasons given, not nearly so commonly as in the situation just described. Furthermore, it is impossible to have a tubercular epiphysitis unless centres of ossification have already appeared in the cartilage. For example (with the exception of the epicondyle) the inner side of the epiphysis of the lower end of the humerus could not be diseased in a child under eleven years, nor could the outer side be effected until the child had reached the age of two. On the other hand, the upper and lower epiphyses of the femur (but not nearly so frequently as the metaphysis), owing to early appearance of ossific centres, may readily be the site of tubercular disease.

The relation which tubercular metaphysitis bears to disease of the neighbouring joints is very definite.

Thus, in the hip-joint, the metaphysis or juxta-epiphyseal portion of the neck is the first portion of the joint to be involved. But here the metaphysis is entirely enclosed within the capsule, and the joint cavity can scarcely escape rapid infection from direct extension.

On the other hand, the metaphysis of the lower end of the femur is not enclosed within the joint cavity of the knee, and is separated from it by a large epiphysis. In this situation the joint is not, as a rule, involved in the customary manner, the primary focus being either synovial or epiphyseal. What has been said of the knee-joint applies equally to the case of the shoulder-joint.

140 *Tubercular Disease of Lower End of Femur.*

The three cases which I mention below, however, demonstrate that tubercular metaphysitis of the lower end of the femur is no less common than in other situations. The disease progresses along the lines of least resistance, through the thin trigone of the femur, presenting as a popliteal abscess above and behind the knee-joint.

In the elbow-joint the metaphyses of all three bones are within the joint cavity, and, as in the hip-joint, tubercular metaphysitis is followed very rapidly by general joint infection.

Diagnosis.—There is little to be said concerning the diagnosis of central tuberculosis of the metaphysis in children. The only early diagnostic sign is thickening of the bone in the region of the epiphysis. Occasionally a similar condition is present in cases of congenital syphilis. In every case an *x*-ray photograph must be taken to demonstrate the tubercular focus and to estimate the extent of operation necessary for its treatment.

Treatment.—It is only necessary to dwell in this paper on the line of treatment adopted in the three illustrative cases—the method is applicable to practically all cases, but I believe it has not been applied to cases of the femur with the frequency which the results deserve, from fear of shortening and lateral deviations.

In some instances the radical operation may give place to more conservative non-operative methods, the progress of the disease being kept under close observation in order that operation may be recommended in progressive cases before the formation of sinuses and mixed infection. The old operation of exposing the bone, and of eradicating the disease by a process of drilling, scraping and drainage may be indicated occasionally, and gives good results.

The operation of choice is, however, to divide the bone

with a Gigli's saw at a level well above the disease, as shown by the skiagram, and to wrench the metaphysis and diaphysis away from the epiphysis by means of a powerful forceps. In no instance can the "epiphyseal line" be injured, for this is part of the epiphyseal mass, which could not, by the use of any reasonable force, be pulled away from the joint. As a rule, in the presence of disease at or near the juxta-epiphyseal line little force is required to remove the diseased portion from the epiphysis, and if difficulty is experienced a few touches of the chisel will complete the separation.

The periosteum in these cases is thickened, and easily separated from the bone. The latter is removed from the tube of periosteum, as is familiar in rib operations, only with very much more facility. Whatever the *rôle* of the periosteum in the production of new bone, it at least acts as a tubular guide to promote regularity of growth.

Without the aid of *x*-ray photographs it would be difficult to believe with what accuracy the bone is replaced by osteogenetic forces in a comparatively short time. (Plates II., III., and IV.)

In my first case, operated upon in 1909, before any similar cases had been put on record, I removed the lower third of the femur in a boy suffering from tubercular metaphysitis. In this case a mixed infection was present, and an old-standing sinus discharged in the popliteal space. The result of the operation is shown in the accompanying skiagram (Plate II.). The boy has perfect use of his leg; there is three-quarters of an inch shortening, and the movements of the knee-joint are normal. The treatment after operation consisted in the application of an 8 lb. weight and pulley with lateral sand-bags.

The second and third cases were also septic from the ex-

istence of old-standing sinuses, but nothing could have excelled the results of "metaphysectomy." The third case, which has but recently left hospital, shows in the skiagram the formation of the new bone as yet incomplete (Plate IV.). The portions of bone removed from the last two cases are illustrated (in Plate I., Figs. 1 and 2)—the specimen from the first case (Plate II.), which was in every respect similar, but larger, was unfortunately lost.

I must again acknowledge my indebtedness to Mr. Stiles, whose work I had an opportunity of seeing, and who has helped to put this whole subject on a sound and scientific basis.

MR. J. BOYD BARRETT said that the difficulty appeared to him to be in the after-treatment of such cases. He had no doubt that great care was required to keep the knee in position after the operation. He asked Mr. Wheeler to illustrate a little of the after-treatment.

MR. W. PEARSON said that some of the points raised in the communication were applicable not alone to these cases, but also to the general treatment of osteomyelitis. He did not think that the points in pathology were absolutely new. He considered that this method of treatment is the only sound method in dealing with a bony cavity of any large extent. He referred to the experience in cases treated by scraping, plugging, &c., in which sinuses kept on discharging as if nothing had been done. He asked whether in these cases the cavity left after the removal of the bone is filled with wax or whether the periosteum is allowed to collapse, because he thought the latter would be sufficient. He considered that some extension was necessary while the new bone was forming, otherwise there would be a great amount of shortening of the limb, and he had no doubt that even in the cases shown there would be some shortening. He thought that it would be helpful if one were to be satisfied

with a less extensive operation—viz., not to excise the whole circumference. He was much interested in the cases which illustrated how rapidly the bone formed in young subjects.

MR. HAUGHTON said the subject of the communication was of extraordinary interest, and the cases proved the excellence of the operation. He pointed out that it was stated by Mr. Harold Stiles that the treatment in such bone cases was more difficult than some major operations in abdominal surgery. His experience was that the vital point in the treatment was the removal of all disease. X-ray photographs should be relied on to ascertain the extent of the disease, and the operation could then be planned precisely. He referred to his experience with osteomyelitis, and said that complete removal of the diseased bone was followed by complete recovery. In a recent case he had removed about eight inches of the diaphysis of the femur, and when the sequestrum was removed there was a three-inch gap. Extension was put on, and in three months' time the wound was healed, and in six months' time the patient was walking on a new column of bone. He again emphasised the vital points—*i.e.*, complete removal and *x*-rays to help.

MR. MAUNSELL said that, in the diagnosis of these cases, he thought that a great many of them were looked on as tubercular which were really septic osteomyelitis, but he had no doubt that afterwards they had become tubercular, although at the start they were either acute or sub-acute osteomyelitis. He was in agreement with the removal of all diseased tissue, but he had seen cavities which had not been laid thoroughly flat, and they healed up perfectly well. In osteomyelitis cases, either tubercular or non-tubercular, he thought that it was not wise to proceed with great haste, and that one should be guided by *x*-rays. He thought that certain cases would heal up with a much more modified operation than the total removal of the diaphysis. He looked on the cases as a useful lesson in physiology, as they helped the surgeon in not being afraid of removing a large quantity of bone. He would, however, be slow to remove a large piece of the diaphysis, and would be satisfied with more moderate measures.

THE PRESIDENT said that these cases recalled to his mind

a case that came under his notice some years ago when on the staff of the Adelaide Hospital. He was asked to procure admission for a boy with disease of the leg. The patient was in a most pitiful condition—emaciated, &c.—and recovery appeared to be out of the question at the time. Four-fifths of one tibia was in a shocking condition from chronic disease of the bone. Mr. Barton cut down and removed the sequestrum, the boy made an extraordinary recovery, and a couple of years afterwards emigrated, having in the meantime grown to nearly six feet in height. He considered that this case illustrated the extraordinary power of recovery seen in these cases.

MR. WHEELER, in replying to the remarks, said he was indebted to Dr. Maurice Hayes for the photographs in those cases. Referring to the after-treatment, he said that this seemed important at first sight, but he found that 7 or 8 lbs. of sand bags were good enough for the pull. He pointed out that the interesting part of the subject was really new, and that it was first pointed out by Lexer, who showed that all the blood-vessels anastomised in this region, and that contusions and sprains easily set up tubercular abscesses in the joint. With regard to Brodie's abscess, he would like to dissociate himself from everything but tubercular disease. As to the periosteum being adherent in each of the three cases he found it easy to separate the periosteum. He used bismuth paste and iodoform, and did not attempt to fill the cavity up. Referring to Mr. Maunsell's remarks, he said that there was no bone or other trouble until the abscess was noticed in the popliteal space. The cases were very chronic when he got them, and there was no hurry about them. In the first case he had tried scraping without effect.

PLATE II.
MR. W. I. DE COURCY WHEELER on "Tubercular Disease
of the Femur."

Recent X-ray of case operated on in 1909. The lower third of the shaft of the femur was removed (between the arrows) and was accurately reformed as shown. Functional result perfect

PLATE III.

MR. W. I. DE COURCY WHEELER on "Tubercular Disease
of the Femur."

X-ray of second case one year after operation. (The bone removed is shown
in Fig. 1 and the new bone lies between the arrows.)

PLATE IV.

MR. W. I. DE COURCY WHEELER on "Tubercular Disease
of the Femur."