complications nearly always develop within three weeks from the onset of the illness. Mild cases are allowed to be out of bed at the end of two weeks. The presence of murmurs and a slight degree of irregularity are no contraindication if the first sound is strong and the heart is not dilated.

The severe cases must be watched for four or five weeks, but after this time the danger from serious heart complication is usually over. In cases with gallop rhythm, absolute rest in bed, upon a liquid diet, is necessary. At the first appearance of vomiting food should be given by rectum. Strychnine gives the best results. Alcohol and digitalis are not well borne. Morphine may be necessary. The after-treatment of all except the mild cases consists in watching the effect of mild exercise upon the heart for several months, and grading it to meet individual requirements.

OBSTETRICS.

UNDER THE CHARGE OF

EDWARD P. DAVIS, A.M., M.D.,

PROFESSOR OF OBSTETRICS IN THE JEFFERSON MEDICAL COLLEGE; PROFESSOR OF OBSTETRICS AND DISEASES OF INFANCY IN THE PHILADELPHIA POLYCLINIC; CLINICAL PROFESSOR OF DISEASES OF CHILDREN IN THE WOMAN'S MEDICAL COLLEGE; VISITING OBSTETRICIAN TO THE PHILADELPHIA HOSPITAL, ETC.

Spontaneous Rupture of the Uterus after Cæsarean Section.—KERR (*British Medical Journal*, June 18, 1904) reports the case of a patient upon whom he had previously performed Cæsarean section, employing the transverse incision across the fundus. The patient was not sterilized. She was admitted to the hospital in a succeeding pregnancy, estimated at the thirtieth week. She was given an enema, after which she complained of pain in the abdomen, but shortly afterward fell asleep. The next morning the pain extended over the entire abdomen, tenderness gradually developed, the pulse rose to 90, the temperature became subnormal, and the breathing more rapid. The patient lay with her feet slightly drawn up, and the fetal parts could be felt with great distinctness through the abdominal wall. There was some blood-stained vaginal discharge.

On performing abdominal section a large quantity of dark-colored blood escaped, and the foetus and placenta lay in the abdomen with the membranes unbroken. The uterus was retracted behind the foetus, and had a transverse rupture through the scar of the previous operation. Hysterectomy was performed, leaving a stump of cervix, and the patient made a good recovery.

Cæsarean Section for Contracted Pelvis.—At a recent meeting of the Obstetrical Society of London (*British Medical Journal*, October 22, 1904), KERR based his paper upon 30 cases of Cæsarean section. His maternal mortality was 6.6 per cent. His morbidity based upon a temperature of 100.5° on more than one occasion was 26 per cent.