aminations and radioscopy are alike negative in spite of pronounced burning or cramplike, inconstant, and nonperiodic pain. This condition may be likened to a causalgia. The site of the irritation inducing sympathetic symptoms may reside either in the stomach itself; in a local disturbance in any other organ in the sympathetic distribution; or, the cause of the irritation may be general—an infection or intoxication, or a disturbance of the ductless glands, in particular the thyroid, genital glands, and adrenals.

Udaondo, C. B. GASTRIC ULCER AND POLYNEURITIS. [Prensa Med. Argent., Vol. 4, 1918, April 20.]

A man of 45 developed polyneuritis at the same time that he had an extensive acute gastric ulcer. The neuritic pains seemed to grow worse or better following the course of the ulcer. It does not appear whether the ulcer caused the polyneuritis or *vica versa* and least of all that they both might have had a common source.

Mendel, L. B., Osborne, T. B. THE RÔLE OF SOME INORGANIC ELEMENTS IN NUTRITION. [J. Biol. Chem., 33, III, 1918.]

The authors in one of the first of a series of rational attitudes towards the problems of metabolism have begun the analysis of what is imperative; that is what needs have sprung up in man's energy system whereby he must have certain stable conditions in his environment in order that function may proceed. By feeding rats on rations in which individual ions have as far as possible excluded, but which are otherwise adequate for growth and nutrition, it has been demonstrated that Ca and P are needed in considerable amounts. Growth for about a year at a nearly normal rate was accomplished upon a ration containing only insignificant quantities of Mg, Na, K, and Cl. Where both Na and K were present in the above small amounts, no growth occurred.

Babonneix, L., and David, H. A CONTRIBUTION TO THE CLINICAL STUDY OF MIGRAINE. [Journ. de Med. e. d. Chirurg. Pratiques, 1918, LXXXIX, p. 174.]

The writers claim that meningeal reactions play a part in the symptomatology of ordinary migraine and also the ophthalmic variety: there is in migraine a hypertension of the cerebrospinal fluid: "this explains the benefit which lumbar puncture gives in some cases (J. A. Sicard)." The writers draw special attention to the aggravation of the pain of migraine by coughing or sneezing, either spontaneous or volitional. They think that at each shock of a cough or sneeze the cerebrospinal fluid, in a state of momentary hypertension, disturbs the sensory trigeminal root, and so provokes a pain which is propagated to the meningeal filaments of that nerve. They regard the pain of migraine as due, not merely to a neuralgia, but to a true radiculargia of the trigeminal nerve. And this radiculargia is itself dependent on hyper-