actually overlap in their ranges and the Wood-cat is to be found far out on the plains, the Grass-cat is unknown in the wooded or riverine districts.

The following are some measurements taken of two large males, the first having been killed as far back as 1873 and the second recently :—

	(1st)	(2nd)
Length, inclusive of tail	38 in.	37½ in.
Tail	12	12
Height before	15	13
Height behind	..	14½
Girth of abdomen	16	14
Girth of neck	11	9½
Length of head	6	
Width of head	5	

Weight of first (very thin) 13 lbs. I have heard of one that scaled 22 lbs.

November 28, 1899.

Dr. HENRY WOODWARD, F.R.S., V.P., in the Chair.

Mr. Oldfield Thomas exhibited the skull of a Baboon recently obtained at Aden by Messrs. Percival and Dodson. It appeared to represent a new species allied to *Papio hamadryas*, but distinguished by its small size, the row of upper cheek-teeth being only 41·5 mm. in length. This species was proposed to be named *Papio arabicus*.

Mr. W. Saville-Kent, F.L.S., F.Z.S., stated that he had devoted considerable attention since the meeting of the last session to the subject of trichromatic or three-colour photography as applied to the correct colour-registration of Zoological and Botanical subjects. With the aid of the lantern he submitted a series of examples upon which he had successfully experimented. These included various species of tropical butterflies, orchids, fishes, lizards, and birds. Among the slides displayed, that of a peacock's feather, in which the characteristic tints were reproduced with marked fidelity, was particularly referred to as a successful demonstration of the capabilities of the system. Gold and Silver Carp, Cuckoo Wrasses (*Labrus mixtus*), and other marine species, taken by Mr. Saville-Kent at the Plymouth Zoological Station, yielded appropriate illustrations of the process as applied to the colour-registration of the more brilliant but notably evanescent hues of fishes. In the bird-section, especial prominence was given to the correct colour-portrayal of the gaily plumaged Australian finches *Poephila gouldi* and *P. mirabilis*. These were

represented as lantern-transparencies prepared from photographs of artistically preserved specimens, and also from the replica of a water-colour drawing of a group of these birds executed by Mr. J. G. Keulemans from living examples.

Mr. Saville-Kent explained that all these photographs had been taken by him with the Sanger-Shepherd colour-screens, of which he exhibited a set, in conjunction with the Cadett "Lightning Spectrum-plate." These screens represented the three primary spectrum colours, red, green, and blue-violet, as enunciated by the late Prof. Clerk-Maxwell, and a separate negative of the subject had to be taken through each respective screen. The transparent positives prepared from these negatives were stained with tints complementary to those through which they were severally taken. That was—the positive resulting from the red-screen negative was stained blue or minus red; that from the green screen, red or minus green; and that produced from the blue-violet screen, yellow or minus blue. Due care being exercised in obtaining the right tint-gradation, and the three stained positives being then superimposed in precise register, an optically perfect presentment or counterfeit of the original subject was mechanically produced. The special method of developing and staining the positives exhibited was, as in the case of the production of suitable colour-screens, associated with the name of Mr. Sanger Shepherd, with whom Mr. Saville-Kent had been working in collaboration.

It was recommended, for the acquirement of perfect registration, that all three of the respective negatives should be taken on a single plate in conjunction with a specially constructed multiple back, of which a sample was exhibited. This, however, was not absolutely necessary. It was competent, in fact, for anyone possessing an ordinary camera to secure correct colour-replicas of desirable objects, using only in conjunction with his instrument the Cadett spectrum-plates and colour-screens referred to. As an illustration of this fact, Mr. Saville-Kent explained that the Peacock's feather, and several other subjects exhibited that evening, had been taken by him with a large-sized Kodak camera, across the lens of which he had simply slung, with the aid of elastic bands, consecutive sections of his multiple-back screen.

The negatives taken for the production of these lantern-transparencies were also available for three-colour printing or process work. In conjunction more especially with such a perfected machine as the newly introduced Orloff Colour-printing Press, there was evidently a wide field thrown open for the cheaper reproduction, by printing methods, of Zoological and Botanical subjects in their correct natural colours. For lantern demonstration, at any rate, the Sanger-Shepherd process as illustrated by him that evening would, Mr. Saville-Kent anticipated, strongly recommend itself to adoption by the many naturalists who had hitherto employed their cameras for the delineation in monochrome only of the subjects of their studies. In the instances, more especially, of the brilliant but fleeting tints of reptiles, fishes,

and marine subjects, as also in those of the tegumentary tissues and appendages of birds, this method of colour-registration would be of invaluable aid to the working artist, who having possession of a correctly prepared transparency might utilize it at his leisure for the elaboration of a finished painting. The successful application of the same process to the duplication, as lantern-transparencies, of coloured figures of zoological subjects had been demonstrated by the example of the pictures of the Gouldian Finches submitted to the meeting. In a like manner, coloured illustrations from other more rare and costly zoological works could be correctly reproduced.

The following papers were read :—

1. General Account of an Expedition to the Gambia Colony and Protectorate in 1898–99. By J. S. BUDGETT, F.Z.S.

[Received November 28, 1899.]

I propose to give a short general account of an expedition recently made by me, under instructions from the Council of this Society, to the river Gambia. This expedition had for its object the general study of the vertebrate fauna of the Gambia, and especially the investigation of the habits of *Protopterus* and *Polypterus*.

The river Gambia lies between the 13th and 14th parallels of North latitude. It flows due west through country which, lying about 100 miles to the north of the equatorial forest-region, is nowhere densely wooded but mostly covered with a somewhat sparse vegetation consisting largely of leguminous trees interspersed with gigantic baobabs (*Adansonia digitata*), the African mahogany (*Kaya senegalensis*), figs and sycamores.

Extensive open plains, which in the rainy season become flooded, border this river along the greater part of its course, while at a very variable distance from the river-bank low hills of dark red conglomerate rise, often abruptly, and occasionally in steep cliffs, to form level plateaux, which in the upper river may be 200 feet high.

The river-bank itself is clothed throughout the year with a rich luxuriant vegetation extending usually about 100 yards from the water's edge. Though here the trees and creepers remain green the year round, yet away from the river the trees lose their foliage in the dry season as completely almost as our own trees in winter.

From the mouth of this river to the country just below Nianimaru, the river is shut in by an almost impenetrable wall of mangroves, sometimes 30 feet in height. Above this point the river, though

tidal, is perfectly fresh. The tides in the dry season make themselves felt for over 200 miles up the river, in fact to the end of navigable water, where there is about a foot rise.

The dry season extends from November to May. Tornadoes usually begin in June, while during July, August, and September there is a total rainfall of about 50 inches. During these months, though the tides make themselves felt, yet there is no change in the direction of the flow, while in August there runs a steady current of about 3 or 4 miles an hour.

In passing up the river the first place of interest is the old Fort James, which was formerly the port of export of the Gambia for the black-ivory trade. It is now being slowly washed away.

About 20 miles further up, the Vintang creek joins the Gambia, and at the junction of the two streams is the village of Vintang; it is seldom that a purely native village is seen at the water's edge, as they are usually on higher ground a mile or so from the river. If there are any tall trees in these villages, they are sure to be the nesting-places of Pelicans and Marabou-birds, which in the neighbourhood of the villages are strictly preserved. The vast flocks of these birds and also of the Balearic Crane are a great feature in the lower river, where there is little else to be seen but continual walls of mangroves, though now and again the monotony is broken by the passage of a native canoe or some trading cutter; but further up the variety of the vegetation is much greater.

Of particular interest to myself were patches of a *Pandanus* growing in the swampy ground at the river-side. The native name of this was Fang jani, which means "It burns itself." It certainly looked as though it deserved this name, for wherever it was seen a portion of every patch was charred with fire, and it was not easy to imagine how this could have been set alight by an external agency.

The great trading station on the Upper Gambia is M'Carthy's Island. To this place the trading cutters bring their cargoes of ground-nuts, the fruit of the plant *Arachis hypogæa*, to be shipped to Europe by the Ocean steamers which make their way up to this island.

On M'Carthy's Island there are two trading establishments or 'factories' as they are termed, and the remains of an ancient military settlement, consisting of Government House, Officers' quarters, and Barracks, formerly occupied by a detachment of the West India Regiment, which was withdrawn about 1870. The Government House alone of these buildings has been kept in repair; and here I established myself in company with Mr. Wainewright, the Commissioner of the district, who, though usually travelling about the district, yet spends a considerable portion of his time here as Governor of the island. I stayed on M'Carthy's Island about one third of my time. To the Governor of the Colony, Sir Robert Llewellyn, I am indebted for allowing me the free use of the Colonial steamer, 'Mansah Kilah,' and also for much hospitality. To Mr. Wainewright, the Travelling Commissioner in the M'Carthy's

Island district, I am greatly indebted for allowing me the use of a portion of the Government House at M'Carthy's Island, and also for the use of his huts in the main towns of his district. Very soon after my arrival at my headquarters, I made a tour through the district with the Commissioner to get some idea of the kind of country that surrounded me.

We started from Nianimaru, which was subsequently made my second headquarters, and where I spent even more time than at M'Carthy's. The chief interest in this tour lay in the people themselves, the country we travelled through not being of great interest from the point of view of its scenery.

Travelling was not difficult, as porters were plentiful, and were employed from one village to the next at the rate of 3*d.* a man, if the distance was not more than 5 miles. At the important towns a court was held, and a stay was made of two days. The courts were held in the open, the chief, the head-man of the town, and the people all sitting round the Commissioner's chair.

There was plenty of time for shooting and no need to carry much in the way of provisions. The bag usually consisted of Bush-fowl and the Barbary Quail, *Pterocles quadricincta*, Guinea-hen, *Ædicnemus*, various Spur-winged Plovers, especially *Lobivanellus senegalus* and *Hoplopterus spinosus*, also Doves and Pigeons as many as were required. The finest of these, as game, was the Green Pigeon (*Treron calva*), which is never seen to approach the ground, being especially fond of the fruit of the fig-tree.

The commonest birds around us, which were not shot for the pot, were numbers of four species of *Coracias*, a *Centropus* known as the "foolish bird" from its fearless habits and its call, which resembles a soft laugh, several species of *Bucerotidae*, generally seen flying clumsily from tree to tree in small flocks; while overhead hovered large flocks of Bee-eaters (*Merops nubicus*), swallow-like in flight and song.

Other common birds everywhere seen in large flocks were the Metallic Starlings (*Lamprocolius auratus* and *L. caudatus*); Wood-Hoopoes (*Irisor senegalensis*) seen in smaller flocks; while the commonest solitary birds were the Long-tailed Shrike (*Corvinella corvina*) and a species of Drongo (*Dicrurus assimilis*). The bushes of course swarmed with *Ploceidae* and *Nectariniidae*.

It being the beginning of the dry season, the grass was everywhere yet high, and it was out of the question to do any mammal-shooting; the only mammals visible were Climbing Squirrels and Monkeys. Burrows of *Orycteropus* were seen, though the animal does not appear to be very common in this region.

The towns visited during this tour were mostly far from the river and were taken in the order Nianimaru, Sukuta, Kaihai, Demfai, Tabanani, Sami, Koreantab, and back to M'Carthy's Island.

Near Kaihai there were news of a Giraffe having been seen, but they appear to be extremely rare in these parts. I heard indirectly that there were two in captivity at Kaies on the Senegal river.

Having returned to M'Carthy's Island on December 5, I devoted myself again to fishing and catching *Polypterus*. I found that all the specimens of *Polypterus lapradii* had already returned to the river from the swamps, where they come up to spawn in the wet season. However, large numbers of the young of *Polypterus senegalus* could still be caught by damming up the swamp-outlets.

This is a favourite way of fishing with the natives. They make dams across the creeks at short intervals, and then leave them in connection with each other for some days. Then damming up the connections, they bale out the water from the lowest compartment, collect the fishes, and proceed to the next compartment.

Very much more difficult is it to catch the *Polypteri* in the river. Nets which were very successful with other river fishes, failed utterly with *Polypterus*. The seine-net and trammel were given up, and the native cast-net was used with better success. The results of weeks of patient work were not encouraging however, and I gradually realized that the time to catch *Polypterus* was during the rainy season, when it had betaken itself to the flooded lands.

However, during these fishing days at Nianimaru, many interesting fishes were caught, and most of the common small Passerine birds were skinned. Moreover, this fishing was not without its dangers and excitement, as a look-out had ever to be kept for Hippopotami which swarmed in all the creeks. Moreover, frequently in the morning, when the trammel-net was examined, a Crocodile (*Crocodilus cataphractus*) or a Sawfish (*Pristis perotteti*) had to be slain. Several specimens of the latter were thus caught up as far as M'Carthy's Island, some of them measuring 9 feet in length.

Fly-fishing was tried without success. The line and hook were used more by the natives than myself. The trammel was found to be the best kind of net to use for the *Mormyridæ*, which were seldom caught in other ways. The *Mormyrids* apparently keep to the bottom of the river, and were seldom taken in the seine near shore.

It was noticed that a very large proportion of the fishes caught in this river were brilliantly coloured red in the ventral posterior portion of the body. Of fishes I believe 40 species were obtained, including 2 Selachians, *Protopterus annectens*, *Polypterus lapradii* and *P. senegalus*, 8 species of Siluroids and 7 *Mormyridæ*, and 18 others belonging to various groups. Most of the fish were tried as food, but there was only one that was really good eating: this was, I believe, a grey mullet and was taken far up the river.

Often the creeks in which the cast-net was thrown were very narrow, and the canoe slid silently amongst the most luxuriant vegetation abounding with Bee-eaters and Flycatchers. Altogether representatives of 108 species of birds were shot, measured, and described; but skins were made only of the smaller birds, of which examples of 52 species were obtained, belonging to 23 families.

With Dr. Gadow's assistance, most of these have been identified. Of the *Upupidæ*, in addition to the gregarious *Irrisor* already

mentioned, several specimens of *Scoptelus aterrimus* were seen and a skin of a male preserved.

On April 4 I took my two fishermen, my cook, and canoe up to a small village in the Kunchow creek called Alimaka, and there had some huts built. At this place again the trammel, the seine, and the cast-net were worked with hope of obtaining numbers of *Polypterus*. As a rule in the afternoon I went out to shoot, and found it a fair place for game.

During the fortnight thus spent at Alimaka, only six *Polypteri* were caught. There were caught also in this creek several specimens of *Gymnarchus niloticus* and some fine specimens of a freshwater Turtle, *Cyclanorbis senegalensis*. Lions were heard here frequently, and Leopards were seen, but at neither did I get a chance of a shot.

On April 20 two English gentlemen and a Frenchman arrived at M'Carthy's Island, on their way to some supposed gold-mines about 300 miles to the east of M'Carthy's Island. I accompanied them a short distance beyond the eastward British frontier to the town of Netebulu; the river is not navigable beyond that point.

Netebulu is an important native town, where a powerful chief named Sandian had his castle and harem. Here we stayed several days as the guests of the chief, and then I parted from the gold expedition, and made my way back overland to M'Carthy's Island, staying on the way a week at Koina.

About 50 miles above M'Carthy's Island the river-banks become high and precipitous, the country around being composed of high plateaux intercepted by valleys. Frequently, however, the edges of the plateaux retreat from the river-bank a mile or so, surrounding wide plains, where one could be fairly certain of finding game.

Along the steep cliffs of the river-bank, vast numbers of Dogfaced Baboons (*Cynocephalus babuin*) might be seen wending their way. Sometimes the cliffs extended so far along the river-side that the Antelope were forced to come down to drink at certain places, and here the ground would be covered with their spoor.

April and May are the best months for big-game shooting. At Koina, large herds of Tankong (*Damaliscus korrigum*) were seen almost every day. Several were shot and a complete skin was made, which, however, suffered severely from the attacks of dogs and insects before it reached England. These herds were composed of males, females, and young of every age. The largest males seemed to lead the herd, though fine males mingled with the females and young as they daily made their way back in long procession from the river-banks to the higher lands.

Large herds were also seen of *Hippotragus equinus*, the Roan Antelope, or Dakoio as the natives call it, but this species was not so plentiful as the Tankong in these parts. A herd of Elands (*Oreos derbianus*) are believed to have been seen in the distance, and I was presented with a skull taken by Mr. Wainewright from a carcass floating down the river.

In the open plains, where clumps of tall dead grass were shaded by a few trees, one might generally count on starting a Konkotong (*Cobus kob*), some Gazelles, or a Harnessed Antelope. The smaller solitary Antelopes were usually found in pairs. Enquiries were instituted everywhere as to the existence in this region of a Zebra, but I could hear nothing of it.

The horns either collected by me or from the natives included those of 9 species:—*Bubalis major*, *Damaliscus korrigum*, *Cobus unctuosus*, *Cobus kob*, *Cervicapra redunca*, *Hippotragus equinus*, *Tragelaphus scriptus*, a second species of *Tragelaphus* not yet determined, and *Oreus derbianus*.

Buffaloes were said to be common on Deer Island, but they were not seen by me, though horns of two forms were obtained from natives.

On the way back, a cutter was taken from Fatotenda to M'Carthy's Island, and after a few days spent at the Government House attending to my collections, and my living fishes and reptiles, I paid a final visit to Nianimaru. During this time, being the latter part of May, the rainy season began and the swampy places became filled with water. The Frogs began to spawn, and several series of stages in development of the different forms were preserved.

Here I first obtained free swimming *Protopterus* with ripe ovaries: examples of 8 Frogs, 3 Chelonians, 5 Lacertilia, and 9 Ophidia, including a *Typhlops*, were also collected about this time.

Returning to M'Carthy's Island, it was found that a number of *Polypterus lapradii* which had been kept in a pool connected with the river in the hope of getting them to spawn had been set free by the rising river. However, during the latter part of June and July a large number of *Polypterus* of both species were obtained, the females of which were crowded with ripe eggs. Artificial fertilization was tried with these, without success. Many were kept in confinement, and some, of which a pair are now exhibited, were successfully brought alive to England.

About the 10th of July, in the same swamp where these fishes were obtained, several nests of eggs were found. These eggs coincided in measurement exactly with the ovarian eggs of *Polypterus*. The young larvæ possessed cement-organs on the front of the head so characteristic of Ganoid larvæ; and other characters led me to assume that they were the young of *Polypterus*. None were reared beyond the larval state, and their identity could not well be established. However, having stayed on the Gambia three months longer than I had intended, and having a number of healthy *Polypteri* full of spawn, I decided to return home.

Just a day or so before leaving M'Carthy's Island I obtained eggs of *Protopterus*. These were watched through the early stages of segmentation, but the young could not be reared. On July 25 I left M'Carthy's Island and returned to England.

Several *Polypteri* and *Protopteri*, 12 young *Crocodilus cata-*

phractus, a Python, 3 *Cyclanorbis senegalensis*, 2 Hinged Tortoises, some Chameleons, and a Serval Cat were brought home alive.

Since my return to England, I have definitely decided that the eggs and larvæ obtained are not those of *Polypterus*. I have, however, I believe, learned enough about the habits of *Polypterus* to encourage me to make a second attempt next year to obtain the developmental stages.

In conclusion, I wish to thank the Society for lending me influence and support, without which the little that has been done by this expedition could not have been accomplished.

2. A Note on the Hatching-stage of the Pagurine Land-crabs. By L. A. BORRADAILE, M.A., F.Z.S., Lecturer in Natural Sciences of Selwyn College, Cambridge.

[Received October 12, 1899.]

The life-history of the Land-crabs of the family Cœnobitidæ is one to which considerable interest attaches, and of which, at present, nothing appears to be known. The family comprises the genera *Birgus* and *Cœnobita*, the robber- or coconut-crab and the land hermit-crabs, all of which have given up a sea life for one on land. It need hardly be remarked that changes in habitat, particularly from sea to land or fresh water, have frequently necessitated the suppression of larval stages in the life-history. Among Crustaceans the instances of the cray-fishes, the ditch-prawn (*Palæmonetes varians*), the freshwater crabs (*Potamon*), and at least one species of land-crab (*Gecarcinus*) come at once to mind. The possibility was thus suggested that the land-pagurines might also have lost the whole or a part of their larval life, and leave the egg in something like the adult condition. On the other hand, it had to be borne in mind that some species of land-crabs and all the strand-crabs (*Ocypoda*, &c.) retain the habit of setting free zoœa-larvæ in the sea, where they pass through their earlier stages.

It was probably with these considerations in his mind that von Willemoës-Suhm, when, in October 1874, the 'Challenger' arrived at Zamboanga in the Philippine Islands, wished to investigate the development of the robber-crab from the egg. Unfortunately the time of year rendered it impossible for him to do this, but he was told by an "intelligent native" that the young were born resembling the parent. This statement has since been accepted in a tentative manner by certain text-books, in spite of the fact that the small size of the eggs (and, indeed, of the female genital opening) made it improbable that the development was a direct one, depending on food-yolk. As for the statement of the

native, it is only what was to be expected. A "native," being unfamiliar with the idea of a metamorphosis, will always give the same answer to any question on the subject, namely that the young are born exactly like the adult—but very small.

All doubt on this point has, however, now been removed by the discovery by Dr. A. Willey of a female *Birgus* on the rocks at the brink of the sea at Lifu in the Loyalty Islands.

The abdomen of this specimen was covered with hatching zoæas which were being washed off into the water. The time of the year was the month of January¹.

With regard to the genus *Cænobita*, I have myself recently taken specimens of two species (*C. rugosus* and *C. perlatus*) in Ceylon and the island of Minikoi in the months of May and June, bearing zoæas. The animals were taken on the stretch of wet sand just above the waves. The catches of the tow-net in the island of Minikoi not having yet been examined for specimens of the zoæa of *Cænobita*, the possibility is not completely excluded that the larvæ may undergo the whole or a part of their development within the shell of the mother, which is always wet with salt water. The larvæ, however, did not, on a cursory examination, give any indication to justify such an assumption, and when placed in sea-water lived for a short time and showed powers of swimming in a lively manner. An attempt to rear them unfortunately failed, but this was only to be expected in view of the known difficulty of the operation. It is perhaps worth noticing that the above two species of *Cænobita* are those whose habits keep them nearest to the sea. If an abbreviated development is to be found in the genus, it would more probably occur in forms such as *C. spinosus* which live at a considerable distance from, or at least a considerable height above, the sea.

From the observations just recorded, it is clear that the early stages of the development of the *Cænobitidæ* present no very remarkable features. It may be presumed that later stages follow the ordinary course. The only points of interest remaining for investigation are the assumption of the adult form by *Birgus* and the transition from sea to land, which we may hope to have described by some future traveller in the Pacific Ocean.

¹ The present writer is under great obligation to Dr. Willey for handing over to him a number of these zoæas, which will be described and figured in Part V. of Dr. Willey's "Zoological Results" now being published by the Cambridge University Press. It is intended also to publish an account of the larvæ of *Cænobita*.

1. *Engraulis encrasicolus* 2. *Clupea harengus* 3. *Ammodytes lanceolatus*. 4. *Megalops cyprinoides*.

5. *Thalustes aculeatus*, 6. *Scomber scombrus*. 7. *Salmo salar*.

8. *Mugil capito*. 9. *Hippoglossus vulgaris*. 10. *Osmerus eperlanus*.

11. *Alburna conorhynchus*. 12. *Malapterurus electricus*. 13. *Cyprinus carpio*.

W.G.R. del.

Bale & Danielsson Lith. imp.

14. *Cobitis taenia*.

15. *Escx lucius*

16. *Anguilla vulgaris*.

17. *Labrax lupus*.

18. *Liocassis longirostris*

19. *Callichthys littoralis*.

20. *Clarias magur*.

21. *Perca fluviatilis*.

22. *Zeus faber*.

23. *Anarrhichas lupus*.

24. *Labrus maculatus*.

25. *Exocoetus brachysoma*.

26. *Lophilus piscatorius*.

27. *Cottus scorpio*

28. *Trigla cuculus*

29. *Cyclopterus lumpus*.

30. *Fistularia tabaccaria*

31. *Saccobranchius fossilis*.

32. *Silurus glanis*.

33. *Syngnathus acus*.

34. *Gadus aeglefinus*.

35. *Blennius pholis*.

36. *Orthogoriscus truncatus*.

37. *Tetodon palembangensis*.

3. On the Relations of the Efferent Branchial Blood-vessels to the "Circulus Cephalicus" in Teleostean Fishes. By W. G. RIDEWOOD, D.Sc., F.L.S., Lecturer on Biology at St. Mary's Hospital Medical School.

[Received August 1, 1899.]

(Plates LXIII.-LXV.)

INTRODUCTION.

When engaged in dissecting a Herring some years ago, I was struck by the fact that the four efferent branchial vessels all reached a median vessel which was continuous behind with the dorsal aorta, and that the circulus cephalicus was situated anteriorly to the first pair of vessels. This condition was so totally different from what I knew to be the arrangement of the vessels in the Cod, that I examined the same parts in a third form, the Salmon, and here found a condition intermediate between the two preceding. The results appeared to warrant a further inquiry, and the present investigation was undertaken with a view to ascertaining what are the commonest, and what the extreme modifications of the efferent branchial arteries to be met with among Teleostean fishes.

The greater part of the work was done during the summer vacation of 1892 at the Marine Laboratory of St. Andrews, Scotland; but owing to the difficulty of making the series of fishes thoroughly representative, the investigation has been protracted over a period of seven years. Late, however, as it may now appear, I take the opportunity of expressing my warmest thanks to Prof. W. C. McIntosh, M.D., F.R.S., for his kindness in placing at my disposal for six weeks during the summer of 1892 all the facilities that are afforded by the St. Andrews Laboratory for the procuring and injecting of the indigenous fishes. I have also to thank Prof. G. B. Howes, LL.D., F.R.S., of the Royal College of Science, London, for specimens of *Perca*, *Trachinus*, *Lophius*, *Fistularia*, *Motella*, *Ammodytes*, *Silurus*, *Exocoetus*, *Esox*, *Albula*, *Megalops*, *Chirocentrus*, and *Hippocampus*, and Mr. G. A. Boulenger, F.R.S., of the Natural History Museum, for specimens of *Corvina*, *Equula*, *Gobius*, *Sphyræna*, *Hemichromis*, *Clarias*, *Saccobranchus*, *Malapterurus*, *Callichthys*, *Scopelus*, *Cobitis*, *Marcusenius*, *Balistes*, *Tetrodon*, and *Orthogoriscus*.

The series of forms examined includes 61 species belonging to 57 genera.

LIST OF SPECIES EXAMINED.

ACANTHOPTERYGII.

- Percidæ.
 Perca fluviatilis.
 Labrax lupus.
 Mullidæ.
 Mullus barbatus.
 Scænidæ.
 Corvina nigra.
 Carangidæ.
 Equula edentula.
 Cyttidæ.
 Zeus faber.
 Scombridæ.
 Scomber scombrus.
 Trachinidæ.
 Trachinus draco.
 Pediculati.
 Lophius piscatorius.
 Cottidæ.
 Cottus scorpio.
 Trigla gurnardus.
 Trigla cuculus.
 Discoboli.
 Cyclopterus lumpus.
 Gobiidæ.
 Gobius giuris.
 Blenniidæ.
 Anarrhichas lupus.
 Blennius pholis.
 Centronotus gunellus.
 Zoarces viviparus.
 Sphyrænidæ.
 Sphyræna vulgaris.
 Mugilidæ.
 Mugil capito.
 Gastrosteidæ.
 Gastrosteus spinachia.
 Fistulariidæ.
 Fistularia tabaccaria.
 Gobiæsocidæ.
 Lepadogaster gouanii.

PHARYNGOGNATHI.

- Labridæ.
 Labrus maculatus.
 Chromides.
 Hemichromis fasciatus.

ANACANTHINI.

- Gadidæ.
 Gadus æglefinus.
 Molva vulgaris.
 Motella triciirrhata.

Ophidiidæ.

- Ammodytes lanceolatus*.
 Ammodytes tobianus.
 Pleuronectidæ.
 Hippoglossus vulgaris.
 Pleuronectes flesus.

PHYSOSTOMI.

- Siluridæ.
 Clarias magur.
 Saccobranchnus fossilis.
 Silurus glanis.
 Liocassis longirostris.
 Malapterurus electricus.
 Callichthys littoralis.
 Scopelidæ.
 Scopelus boops.
 Cyprinidæ.
 Cyprinus carpio.
 Hypophthalmichthys nobilis.
 Cobitis tænia.
 Scombresocidæ.
 Exocoëtus brachysoma.
 Esocidæ.
 Esox lucius.
 Mormyridæ.
 Marcusenius plagiostoma.
 Salmonidæ.
 Salmo salar.
 Salmo trutta.
 Osmerus eperlanus.
 Coregonus oxyrhynchus.
 Clupeidæ.
 Engraulis encrasicolus.
 Clupea harengus.
 Clupea sprattus.
 Albula conorhynchus.
 Megalops cyprinoides.
 Chirocentridæ.
 Chirocentrus dorab.
 Murænidæ.
 Anguilla vulgaris.

LOPHOBRANCHII.

- Syngnathidæ.
 Syngnathus acus.
 Hippocampus guttulatus

PLECTOGNATHI.

- Sclerodermi.
 Balistes aculeatus.
 Gymnodontes.
 Tetrodon palembangensis.
 Orthogoriscus truncatus.

LITERATURE.

On searching through the literature of the subject, one cannot fail to be struck by the fact that the arterial system of Teleostean fishes has been very greatly neglected, and the disposition of the efferent branchial vessels particularly so. Meckel and Hyrtl appear to have been the only anatomists to undertake anything like a systematic study of these latter. Meckel in 1831 (14. p. 192) pointed out that the mesial ends of the third and fourth efferent branchial vessels are usually close together, and may unite with one another before joining the aorta; and he furnished a few observations on the efferent branchial system of *Gadus*, *Trigla*, *Perca*, *Pleuronectes*, *Lophius*, and *Muraena*. Seven years later Hyrtl (7) gave a table showing the proportions of the circulus cephalicus in fourteen species of Teleosteans, supplemented by a considerable amount of information concerning the vessels associated with the circulus, and good figures of the efferent branchial system of *Perca*, *Gadus*, and *Tinca*.

Stannius in 1849 (23. pl. v.) published some fairly reliable figures of the efferent vessels of *Cyclopterus*, *Gadus*, *Salmo*, and *Scomber*, but the blood-vessels were only introduced into his figures to act as landmarks for the recognition of the sympathetic nerves, and must not be treated too critically. In his 'Handbuch' of 1854, however (24. p. 242), he made reference to the fact that the circulus cephalicus is wide in *Gadus* and *Lota*, where all the efferent branchial vessels open into it, whereas it is narrow in *Scomber* and *Salmo*, in which genera the last two open directly into the aorta. The only other information on the subject is that conveyed by the figures of the Carp by Duverney (6. pl. ix. figs. 17 and 18), the Perch by Laurillard in Cuvier's 'Histoire Nat. des Poissons' (5. pl. vii. fig. 1), the Cod by Müller (16. pl. iii. fig. 13), the Trout by Vogt (1. pl. L. fig. 2), the Pike by Maurer (13. pl. xi. fig. 1), the Cod by T. J. Parker (20. p. 117), and the contributions by Hyrtl on *Heterotis* (9), *Gymnarchus* (11), *Chanos* (12), and other genera.

GENERAL PART.

As may be gathered from the title, the observations recorded in this paper concern the efferent branchial vessels and the vessels formed by their confluence. The cœliaco-mesenteric and sub-clavian arteries usually arise in relation with the hinder part of the circulus cephalicus, or with that part of the aorta which receives the third and fourth efferent branchial vessels. The positions of these arteries are indicated in the figures, and occasional references are made to them in the text; but the investigation does not profess to deal exhaustively with these vessels, nor with the hyoidean, anterior carotid and posterior carotid arteries, which also are associated with the circulus cephalicus.

The dotted lines in the figures signify that owing to the failure of the injection-mass to pass, or owing to the small size of the fish

and the consequent difficulty of dissection, or to the bad state of preservation of the parts, it was not possible to trace out the vessels so delineated, but that, from analogy with other forms, there is reason to believe that the vessels occupy the positions indicated. There appears to be no reason to suppose that the circulus cephalicus is ever incomplete in front. The lumen of the transverse commissure between the roots of the anterior carotid arteries may possibly be closed in some cases, but the failure of the injection-mass to pass into the commissure is not necessarily a proof of the fact, since the pressure during the process of injection is equal on the two sides of the circulus. The transverse vessel is usually of small size, and its traversing the parasphenoid bone makes it difficult to dissect out with any degree of neatness.

In the selection of characters by which to classify the various types of arterial disposition, it has been assumed that the condition found in *Clupea* (Pl. LXIII. fig. 2) and *Engraulis* (fig. 1), where the circulus cephalicus is small, and does not involve the second, third, and fourth efferent branchial vessels, is the most simple and primitive, and that the connection of all four efferent branchial arteries with the circulus, such as occurs in *Gadus* (Pl. LXV. fig. 34), is the most specialized. This assumption is based partly upon the fact that the Gadoids are highly specialized in numerous other respects, whereas the Clupeoids are generally recognized as among the lowest of the Teleostean series; partly upon the fact that in *Amia*, an admittedly primitive Ganoid with Clupeoid affinities, the last three efferent branchial vessels are unconnected with the circulus cephalicus¹; partly also upon the researches of Ayers (4) upon Elasmobranch fishes, which go to prove that the right and left sides of the circulus cephalicus are not the primitive paired aortæ such as occur in *Amphioxus* and in embryos of the true Vertebrata, but that the true dorsal aorta may persist as a median vestigial vessel traversing the circulus cephalicus, in the same manner as, according to Müller (16), it does in the Cyclostomi.

Pursuing this line of argument, we may legitimately conclude that where, as in the Salmon (fig. 7), Mackerel (fig. 6), and Carp (fig. 13), the circulus cephalicus, receiving the first and second efferent branchial vessels, is separated from the point of entry of the third and fourth by a length of the median aorta, the condition is more primitive than that in which the third and fourth vessels open at the posterior extremity of the circulus cephalicus, as in the Bass (Pl. LXIV. fig. 17). And further, the separation of the third and fourth vessels in the Anchovy (fig. 1) by a portion of the aorta indicates a more lowly condition than that seen in the Herring (fig. 2), where the two vessels open close together. There are thus two lines upon which we may regard specialization as proceeding: firstly, by the circulus cephalicus

¹ If such exists. The circulus appears to be suggested in Allis's figure (3. pl. xxxvi.), but its existence is denied by Ramsay Wright (25. p. 495). The arrangement in *Lepidosteus* is somewhat similar to that of *Amia*. See Hyrtl, 8. p. 235, and Müller, 17. pl. v. fig. 6.

involving the second, and later the third and fourth efferent branchial vessels; and secondly, by the progressive suppression in length of the median aorta, bringing about an approximation of the dorsal or proximal ends of the last three efferent branchial vessels on each side.

Owing to the fact that the one variety of specialization may occur quite independently of the other, or in conjunction with it, it becomes very difficult in some cases to compare the ultimate degree of specialization attained, since there is no evidence to show whether the inclusion of the efferent vessels into the circulus or the approximation of the efferent vessels by the suppression of the aorta is the more important. In the Salmon (fig. 7), for instance, the second efferent vessel opens into the circulus cephalicus—an indication of specialization; but a length of aorta persists between the circulus cephalicus and the third efferent vessel—a primitive character. In *Balistes* (fig. 5) the second vessel is free from the circulus cephalicus, and yet there is obvious specialization in the complete suppression of the aorta in the branchial region, resulting in the second, third, and fourth vessels opening close together, immediately behind the circulus. Who shall say whether, in the disposition of the efferent branchial vessels, the Salmon or the File-fish is the more primitive? Having recourse to the other anatomical features of these two forms, one would conclude that, the Salmon being in general structure the more primitive, the abbreviation of the aorta is as a mode of specialization more important than the backward extension of the circulus to include the second efferent branchial vessels. The conclusion is further justified by the fact that *Albula* (fig. 11), which is undoubtedly allied to *Megalops* (fig. 4) and *Chirocentrus*, differs from these genera in this latter respect.

This hypothesis, however, opens up the further question as to how far a backward extension of the circulus cephalicus is due to the longitudinal splitting of a part of the median aorta. Has, for instance, the condition found in *Megalops* and *Chirocentrus*, in which the circulus cephalicus extends back to the second branchial vessels, been brought about by the longitudinal division of a median vessel such as exists in *Clupea* (fig. 2) and *Engraulis* (fig. 1) between the first and second efferent branchial vessels? The suggestion has much to recommend it; more especially as the suppression of the median aorta cannot have operated here, or the two anterior carotids would be arising close together at the bottom of the fork of the first efferent branchials.

Another line of specialization, independent of the two former, can be traced in the confluence of the third and fourth efferent vessels. Having assumed that the separation of two consecutive efferent branchial vessels by a portion of the median aorta is a primitive feature, it follows that the separate entry into the aorta of the third and fourth vessels in *Engraulis* (fig. 1) is an indication of less specialization than the debouching of the two vessels together, as in *Clupea* (fig. 2); and further, that this latter

condition is more primitive than that found in *Megalops* (fig. 4), in which the third efferent vessel unites with the fourth on each side to form a short common trunk which carries the double charge of blood to the aorta. If this line of argumentation be extended to the cases in which the third and fourth vessels open into the circulus cephalicus instead of into the aorta, we are led to the conclusion that *Blennius* (fig. 35), having the Y-shaped system, is more specialized than *Gadus* (fig. 34), where the third and fourth vessels are disposed in the form of a V, and that this latter is more specialized than *Syngnathus* (fig. 33), where the two vessels find separate outlets into the circulus.

Whether much importance, however, can be attached to this last feature is open to considerable question, for the V-shaped system obtains in *Salmo* (fig. 7) and the Y-shaped one in *Osmerus* (fig. 10), and a similar relation exists between the Chinese Carp *Hypophthalmichthys* and our native Carp *Cyprinus* (fig. 13), while among the Siluroids, a presumably natural assemblage of forms, there are gradations from *Callichthys* (fig. 19) and *Liocassis* (fig. 18) with the V-shaped system, through *Clarias* (fig. 20), and *Silurus* (fig. 32), to *Saccobranchus* (fig. 31), with a typical Y-shaped arrangement.

The disposition of the efferent branchial vessels is independent of the shape of the head, except in so far as the slope of the vessels and the shape of the circulus is concerned. There is no connection, that is to say, between the shape of the head and the degree of suppression of the median aorta, or the entry of the efferent branchial vessels into the circulus rather than into the aorta. In long-headed forms like *Ammodytes* (fig. 3), *Sphyræna*, *Fistularia* (fig. 30), *Anguilla* (fig. 16), and *Syngnathus* (fig. 33) the circulus cephalicus is elongated in an antero-posterior direction; while in *Cottus* (fig. 27), *Lophius* (fig. 26), and others, with a broad, flat head, the gills are widely separated, and the circulus cephalicus is proportionately broad, the common trunks formed by the fusion of the third and fourth efferent vessels being also lengthened.

The differences in the arrangement of the efferent branchial vessels relatively to the circulus cephalicus and the aorta are not correlated with any differences in the position and extent of development of the epipharyngeal dentition. At the commencement of the inquiry, the suggestion occurred to my mind that the development of a large and elaborate dental apparatus might, by some process of natural selection, have resulted in the blood-vessels taking up a position of safety, out of the line of direct pressure between the epipharyngeal bones and the vertebral centra or the base of the skull. A minute examination of the individual cases shows, however, that the vessels do not experience any displacement under the circumstances, but obtain sufficient protection by running in grooves or arches in the epipharyngeal bones; in fact, as often as not, the epipharyngeal teeth lie immediately below certain of the efferent branchial blood-vessels. As an instance of two forms

with very similar vascular arrangement, but with widely different epipharyngeal dentition, may be mentioned the Salmon and the Mackerel. In the Salmon there are a pair of small patches of teeth borne by the fourth epibranchial bones, lying immediately ventral to the aortic extremities of the third efferent branchial vessels; but in the Mackerel there are a pair of great dentigerous pads underlying the mesial ends of the second, third, and fourth efferent branchial vessels, as well as a considerable part of the circulus cephalicus. In the Carp the dorsal aorta obtains the necessary protection by actually traversing the great horn-covered bony projection of the basioccipital, against which the lower pharyngeal teeth bite. In the Wrasse the vascular arrangement conforms to a very average type, being apparently quite unaffected by the large and elaborate pharyngeal mill developed in the vicinity of the posterior half of the circulus cephalicus. In *Exocoetus*, again, the epipharyngeal pad is of relatively enormous size, underlying the upper extremities of all four efferent branchial vessels, the anterior ends of the aorta and cœliaco-mesenteric artery, and the whole of the circulus cephalicus except the extreme anterior part; yet there is nothing very remarkable in the disposition of these vessels which might be accounted for by their relation to the epipharyngeal apparatus. Only in *Gobius*, *Cottus*, and *Molva*, of the forms examined, are the epipharyngeal dental pads situated entirely within the circulus cephalicus. The evidence afforded by forms devoid of epipharyngeal teeth is probably inconclusive, on account of the great possibility of the edentulous condition having been arrived at independently in different groups of fishes. In the Sprat and Pipe-fish the types of vascular arrangement are widely divergent. The Sprat closely resembles the Herring (fig. 2); the vessels of *Syngnathus* are shown in figure 33.

In order to discuss intelligibly the different forms of vascular arrangement met with, some form of classification, however artificial, is essential; and after careful consideration I have found it convenient to adopt the following scheme, based on the relations of the efferent branchial vessels to the circulus cephalicus and the dorsal aorta. In consideration of the unsatisfactory nature of the present classification of Teleostean fishes, a purely artificial scheme, founded upon the single character which forms the basis of the present communication, is likely to prove of more permanent utility for purposes of subsequent reference, than one which relies upon a classification which sooner or later may prove to be an unnatural grouping.

The great majority of the forms studied will be seen to come under the headings B and C, while the rarer and more extreme modifications occupy terminal positions in the classificatory scheme. The division of the groups B and C into the subgroups *b* and *c*, differing only in the transverse or oblique position of the confluent third and fourth efferent branchial vessels, appears arbitrary, but in practice there are very few forms which fail to

fall definitely into the one or the other subgroup. Without this subdivision, such remarkably different forms as *Anarrhichas* (fig. 23) and *Tripla* (fig. 28) would come under the same heading.

CLASSIFICATION OF THE GENERA EXAMINED.

Group A. The first efferent branchial vessel opens into the circulus cephalicus, but the second does not.

Subgroup a. The third and fourth vessels open into the median aorta separately.—*Engraulis* (fig. 1).

Subgroup b. The third and fourth vessels open into the median aorta together.—*Clupea* (fig. 2), *Ammodytes* (fig. 3).

Subgroup c. The third and fourth vessels on each side unite to form a common trunk, which reaches the aorta some distance behind the second vessel.—*Chirocentrus*, *Megalops* (fig. 4).

Subgroup d. The third and fourth vessels on each side unite to form a common trunk, which reaches the median aorta immediately behind the second vessel.—*Balistes* (fig. 5).

Group B. The first and second efferent branchial vessels open into the circulus cephalicus, and the third and fourth into the median aorta at some distance behind the circulus.

Subgroup a. The third and fourth vessels open into the aorta together.—*Scomber* (fig. 6), *Hypophthalmichthys*, *Salmo* (fig. 7), *Coregonus*.

Subgroup b. The third and fourth vessels unite before opening into the aorta, the common stem being transverse to the length of the body.—*Mugil* (fig. 8), *Hippoglossus* (fig. 9), *Pleuronectes*, *Osmerus* (fig. 10), *Albula* (fig. 11).

Subgroup c. The third and fourth vessels unite, the common stem sloping backwards towards the aorta.—*Malapterurus* (fig. 12), *Scopelus*, *Cyprinus* (fig. 13), *Cobitis* (fig. 14), *Esox* (fig. 15), *Marcusenius*, *Anguilla* (fig. 16).

Group C. The first and second efferent branchial vessels open into the circulus cephalicus, and the third and fourth into the aorta immediately behind the circulus.

Subgroup a. The third and fourth vessels open into the aorta together.—*Labrax* (fig. 17), *Liocassis* (fig. 18), *Callichthys* (fig. 19), *Clarias* (fig. 20).

Subgroup b. The third and fourth vessels unite before opening into the aorta, the common stem being transverse to the length of the body.—*Perca* (fig. 21),

Mullus, *Corvina*, *Equula*, *Zeus* (fig. 22), *Trachinus*, *Anarrhichas* (fig. 23), *Zoarces*, *Centronotus*, *Sphyræna*, *Labrus* (fig. 24), *Hemichromis*, *Exocoetus* (fig. 25).

Subgroup c. The third and fourth vessels unite, the common stem sloping backwards towards the aorta.—*Lophius* (fig. 26), *Cottus* (fig. 27), *Trigla* (fig. 28), *Cyclopterus* (fig. 29), *Gobius*, *Fistularia* (fig. 30), *Lepadogaster*, *Saccobranchus* (fig. 31), *Silurus* (fig. 32).

Group D. The four efferent branchial vessels open into the circulus cephalicus.

Subgroup a. The third and fourth vessels open separately.—*Syngnathus* (fig. 33), *Motella*.

Subgroup b. The third and fourth vessels open together.—*Gastrosteus*, *Gadus* (fig. 34), *Molva*, *Hippocampus*.

Subgroup c. The third and fourth vessels unite before joining the circulus cephalicus.—*Blennius* (fig. 35), *Orthogoriscus* (fig. 36), *Tetrodon* (fig. 37).

SPECIAL PART.

In the case of species not figured, comparisons will be drawn with those figured forms which, in the arrangement of the efferent branchial vessels, they most nearly resemble, irrespective of the degree of affinity which upon other grounds may be considered to exist between the forms.

FURTHER REMARKS ON GROUP A.

In *Engraulis* (fig. 1), there is a considerable difference in the size of the four efferent branchial vessels; the second is the broadest, and the first the narrowest. The dorsal aorta behind the fourth branchial vessel is wide and very thin-walled. This is also the case in *Clupea*. Although the scheme of classification which it has been found convenient to adopt brings *Ammodytes* (fig. 3) under the same subgroup as *Clupea* (fig. 2), there are several features which go to show that the association is an unnatural one. The circulus cephalicus in the former genus is much longer than broad, and extends back nearly to the point of entry of the second pair of vessels into the aorta; whereas in *Clupea* the posterior part of the circulus is transverse to the axis of the body, and forms with the first part of the aorta a T rather than a Y. The celiac-mesenteric artery arises in *Ammodytes* close behind the fourth efferent vessels, but much further back in *Clupea*. The origin of the subclavian arteries is slightly more posterior in *Clupea* than in *Ammodytes*. There are no differences between *Ammodytes lanceolatus* and *A. tobianus*, nor between *Clupea harengus* and *C. sprattus*.

Chirocentrus closely resembles *Megalops* (fig. 4). In both genera the median aortic stem found in *Engraulis* and *Clupea*

between the first and second efferent branchial vessels is wanting, owing to the backward extension of the circulus to the point of entry of the second branchial vessels into the aorta. The same feature is to be observed in *Balistes* (fig. 5), where the aortic stem between the second vessel and the common trunk of the third and fourth is also suppressed. The coeliaco-mesenteric artery of *Balistes* arises, not from the aorta, but from the third and fourth branchial vessels of the right side, immediately after their anastomosis.

If we disregard the modification of the efferent branchial circulation brought about in *Gymnarchus* by the increased functional importance of the swim-bladder, the efferent branchial system of this genus can be seen, from the description and figure published by Hyrtl (11. p. 11, and pl. 4. fig. 4), to conform with the type which characterizes subgroup A c.

FURTHER REMARKS ON GROUP B.

The efferent branchial system of *Scomber* (fig. 6) bears a close resemblance to that of *Salmo* (fig. 7), but the coeliaco-mesenteric artery arises from the aorta immediately behind the fourth efferent branchial vessels and the subclavian arteries some distance farther back, whereas in *Salmo* the positions of the coeliaco-mesenteric and subclavian arteries are reversed. In the figure of *Scomber* given by Stannius (23. pl. v. fig. 4) the circulus cephalicus appears much too large, and the interval between the circulus and the entry of the third and fourth branchial vessels into the aorta too short.

In *Salmo trutta*, as also in *Osmerus* and *Coregonus*, the dorsal aorta is wide and thin-walled; but this is not the case in *Salmo salar*. The distance between the posterior angle of the circulus and the point of entry of the third efferent branchial vessel is proportionately longer in *Coregonus* than in *Salmo salar*, and proportionately shorter in *Salmo trutta*. A very reliable figure of the efferent branchial system of *S. trutta* has been given by Vogt (1. pl. L. fig. 2). The figure by Stannius of *S. salar* (23. pl. v. fig. 3) is incomplete, but is correct so far as it goes. In *Hypophthalmichthys* both the coeliaco-mesenteric and the subclavian arteries arise some distance behind the fourth efferent vessel. The circulus in this genus is broader than long.

In *Hippoglossus* (fig. 9), *Pleuronectes*, *Mugil* (fig. 8), *Esox* (fig. 15), and *Marcusenius* the distance between the posterior angle of the circulus cephalicus and the point of entry of the common trunk of the third and fourth branchial vessels is so short that these forms approach somewhat closely those included in subgroups Cb and Cc. In *Pleuronectes* the circulus cephalicus is longer than broad, whereas in *Hippoglossus* it is broader than long; the anterior carotids are closer together, and the transverse commissure between them is shorter than in *Hippoglossus*. The common trunk formed by the union of the third and fourth branchial vessels of the right side is much shorter than that

on the left, and the origin of the cœliaco-mesenteric artery is nearer the middle line than in *Hippoglossus*. In *Mugil* (fig. 8) the circulus cephalicus is considerably longer than broad, and the cœliaco-mesenteric artery arises from the median aorta behind the entry of the third and fourth branchial vessels, and not as in the two preceding genera. In *Osmerus* (fig. 10) and *Albula* (fig. 11), also, the cœliaco-mesenteric artery arises from the median aorta, but the subclavian arteries take their origin immediately behind the point of entry of the common trunk of the third and fourth branchial vessels into the aorta, and not behind the cœliaco-mesenteric artery as in *Mugil*.

In *Scopelus* the circulus cephalicus is small and nearly circular in shape, and the portion of median aorta intervening between its posterior angle and the mesial ends of the common trunks of the last two branchial vessels is unusually long. In *Malapterurus* (fig. 12) and *Esox* (fig. 15) the cœliaco-mesenteric artery arises from the aorta immediately ventral to the mesial ends of the common trunks of the third and fourth vessels; in *Cyprinus* (fig. 13), *Cobitis* (fig. 14), and *Marcusenius* it arises more posteriorly, and in *Anguilla* (fig. 16) considerably farther back. Except in this latter respect, the arrangement of the vessels in *Marcusenius* very closely resembles that of *Esox*.

Judging by the excellent figure of *Tinca* published by Hyrtl (7. pl. iv.), this genus exactly resembles *Cyprinus* in the disposition of the vessels of the efferent branchial system. The efferent vessels of *Cyprinus* were figured by Duverney (6. pl. ix. figs. 17 & 18) nearly 140 years ago; and although the figures are incomplete, the essential features are correctly represented. In *Cyprinus*, and according to Hyrtl in *Tinca* also, there are two pairs of subclavian arteries. The anterior pair, arising in front of the common trunks of the last two branchial vessels, supply the upper part of the pectoral arch, the posterior pair the lower part of the arch and the pectoral fins. A somewhat similar arrangement obtains in *Esox*. The mode of origin of the anterior pair has been remarked by Müller, and quoted by Stannius (22. p. 103, footnote 3) and Owen (18. p. 270, and 19. p. 489), and these vessels are shown, although not named, in Maurer's figure (13. pl. xi. fig. 1).

The circulus is large in *Malapterurus* and *Anguilla*; and in the latter genus a median vessel, occupying the position of the anterior continuation of the primitive median aorta described in Selachian fishes by Ayers (4), may be traced forward from the posterior angle of the circulus. It soon forks, and is ultimately lost in the mucous membrane of the roof of the pharynx. In *Anguilla*, also, the origin of the posterior carotid artery is much farther removed from the entry of the first branchial vessel into the circulus than is usual. Meckel has stated (14. p. 193) that in the marine Eel, *Muraenophis helena* (*Muraena helena*), the anterior lateral trunk formed by the union of the first and second branchial vessels is three times as long as the posterior one formed by the union of the third and fourth.

The efferent branchial vessels of *Chanos* have been described and figured by Hyrtl (12. pl. i. fig. 1). The first and second vessels open into the circulus cephalicus, the third some distance farther back into the median aorta, while the vessels from the fourth gills, after uniting with those from the epibranchial organs, unite with one another and open into the aorta at a point as far behind the opening of the third branchial vessel as the latter is behind the circulus cephalicus. The genus thus falls into group B, but, owing to the exceptional disposition of the fourth pair of vessels, it cannot be included in any of the three subgroups recognized.

FURTHER REMARKS ON GROUP C.

The circulus cephalicus is narrow in front in *Labrax* (fig. 17), but it is broad in the Siluroids *Liocassis* (fig. 18), *Callichthys* (fig. 19), and *Clarias* (fig. 20). In *Labrax* the cœliaco-mesenteric artery arises from the aorta immediately behind the fourth efferent branchial. It has the same relations in *Clarias*, but is somewhat more posterior in *Callichthys*, and considerably so in *Liocassis*. The subclavian arteries arise behind the cœliaco-mesenteric artery in *Labrax*, *Clarias*, and *Callichthys*, but close behind the fourth branchial vessel in *Liocassis*. Although *Clarias* is introduced into the subgroup Ca, it really occupies an intermediate position between Ca and Cb, since there is a very short common trunk on each side between the last two branchial vessels and the aorta. The genus *Heterotis* I have not been able to examine, but it is tolerably certain, from the description given by Hyrtl (9. p. 87), that it should be included in the subgroup Ca.

In *Mullus* and *Corvina* the disposition of the cœliaco-mesenteric and subclavian arteries is as in *Perca* (fig. 21), but the circulus cephalicus is broader in front. The circulus has the form of a regular heptagon in *Mullus*, while in *Corvina* it is pear-shaped, the broad end being anterior. In *Equula* it is oval in shape and longer than broad; but otherwise the relations of the parts are as in *Perca*. With regard to *Perca* itself, Hatchett Jackson, in his edition of Rolleston's 'Forms of Animal Life' (21. p. 88), states that the "cœliaco-mesenteric artery * * * springs from the right epibranchial artery before it fuses with its fellow." This does not accord with my own observations. Hyrtl in his figure of *Lucioperca* (7. pl. i. fig. 1) shows the common trunks of the third and fourth branchial vessels entering the aorta at some distance from the posterior angle of the circulus. The transverse commissure, also, between the anterior carotids ("die vorderen oder kleinen Kopffarterien") is situated farther forward than in *Perca*.

In *Zeus* (fig. 22) the circulus is broader, and the cœliaco-mesenteric artery arises, not from the aorta, but from the common trunk of the last two branchial vessels of the right side. *Zoarces* resembles *Anarrhichas* (fig. 23) in the oval shape of the circulus and in the narrow anterior prolongation of the latter, but the

common trunk formed by the union of the third and fourth branchial vessels is shorter. In *Centronotus* and *Trachinus* the anterior prolongation of the circulus is wanting, but otherwise the vessels of the pharyngeal roof are disposed as in *Anarrhichas*. The circulus of *Sphyræna* is twice as long as broad, and the posterior angle is very acute; the subclavian arteries arise farther forward than in *Anarrhichas*, and have more the relations of those of *Labrus* (fig. 24).

Hemichromis differs from *Labrus* in having a much broader circulus cephalicus, and in the more posterior origin of the subclavian arteries. The circulus of *Exocoetus* (fig. 25) is oval and longer than broad; the coeliaco-mesenteric artery arises from the circulus cephalicus immediately to the right of the aorta.

Owing to the absence of the fourth gill and its efferent vessel in *Lophius* (fig. 26), the right of this form to rank under subgroup *Cc* rather than *Ca* is somewhat conjectural. The matter is, however, of no great importance. The circulus is very wide, and the coeliaco-mesenteric artery, which is considerably thicker than the aorta, branches soon after its origin. Concerning *Lophius* Meckel has written (14. p. 192): "*Lophius piscatorius* hat, statt der gewöhnlichen drei bis vier, nur zwei sehr lange Kiemenblutadern. Von der vordersten Kieme entsteht ein einfacher Stamm, der zweite wird durch die Vereinigung der zweiten und dritten Kiemenblutader gebildet, die ungefähr eben so lang als der gemeinschaftliche Stamm getrennt verlaufen." My own observations are thus not in accord with those of Meckel. Most ichthyologists admit, with Müller (17. p. 47), that the three gills present in *Lophius* are the anterior three of the four gills more normally present; and the coupling of the first and second efferent branchial vessels, the third remaining solitary, is by analogy with allied forms a far more intelligible arrangement than that described in the above-quoted passage from Meckel's text-book.

In *Cottus* (fig. 27), although the circulus is so wide in front, the anterior carotids lie very close together, and the transverse commissure, which takes a curious bend forward, is therefore short. The coeliaco-mesenteric artery of *Trigla* (fig. 28) is a double vessel arising from the common trunk of the last two branchial vessels of the right side. There are no differences between the efferent branchial systems of *Trigla cuculus* and *T. gurnardus*. The anterior part of the circulus cephalicus of *Cyclopterus* (fig. 29) is very curiously shaped, and the transverse vessel may possibly be wanting. I have been unable to find it in the three specimens dissected. In the figure of *Cyclopterus* given by Stannius (23. pl. v. fig. 1) the subclavian arteries are drawn too wide; and they are incorrectly described on page 156 as branchial veins. *Lepadogaster* does not differ materially from *Cyclopterus*, except that the anterior part of the circulus cephalicus conforms more to the normal type. *Gobius* differs from *Cottus* (fig. 27) in the wider separation of the anterior carotids, the greater breadth of the circulus cephalicus, the more posterior entry of the second

branchial vessels into the circulus, and the more posterior origin of the subclavian arteries.

In *Fistularia* (fig. 30) the slope of the common trunks of the third and fourth efferent branchial vessels is so slight that the genus might with equal propriety be classed under subgroup C *b*, and the fact that these common trunks do not enter exactly at the posterior angle of the circulus cephalicus makes it difficult to uphold its claim to come into group C at all. The circulus is long and abruptly terminated in front. The aorta, after giving origin to the coeliaco-mesenteric artery, is unsymmetrical, and runs to the left side of the vertebral centra. In *Silurus* (fig. 32) the posterior angle of the circulus cephalicus is not exactly coincident with the mesial ends of the common trunks of the efferent branchial vessels 3 and 4, as it is in *Saccobranchus* (fig. 31); and thus the form really occupies an intermediate position between the subgroups C *c* and B *c*, in which latter subgroup the Siluroid genus *Malapterurus* (fig. 12) has already been placed. With regard to *Saccobranchus*, it has already been pointed out by Hyrtl (10. p. 306) that the first branchial vessel unites with the second (which is another way of stating that they both open into the circulus), the third with the fourth, and that the efferent vessels of the lung-sac open into the fourth branchial vessel.

FURTHER REMARKS ON GROUP D.

In *Syngnathus* (fig. 33) the dorsal aorta is not median but runs slightly to the left side of the vertebral centra. The coeliaco-mesenteric artery arises at the posterior angle of the circulus cephalicus. From the same place arises a single vessel forking posteriorly into the two subclavian arteries. *Hippocampus* closely resembles *Syngnathus*, but since the openings of the third and fourth efferent vessels are closer together, the genus comes under the second subgroup, D *b*. The circulus cephalicus, also, is less elongated than in *Syngnathus*. In both genera, however, the front of the circulus is broad and its posterior angle very acute. The dorsal aorta of *Hippocampus* is median, and the subclavian arteries arise from its sides directly, and not through the intervention of a common root. In *Gastrosteus* the circulus cephalicus is not more than twice as long as broad. The subclavian arteries arise a short distance behind its posterior angle, and the coeliaco-mesenteric a considerable distance behind.

Molva differs from *Gadus* (fig. 34) mainly in the fact that the subclavian arteries arise from the circulus cephalicus farther from the median line, and consequently more remote from the dorsal aorta. The coeliaco-mesenteric forks close to its origin in both genera, as it also does in *Motella*. *Motella* very closely resembles *Molva*, and it is only the slight separation of the mesial extremities of the last two branchial vessels which causes the genera to be placed in separate subgroups. *Gadus callarias* (*G. morrhua*), figured by Müller (16. pl. iii. fig. 13) and by Stannius (23. pl. v. fig. 2), does not appear to differ materially from *Gadus aeglefinus*,

except that the subclavian arteries arise farther from the median line, just as they do in *Molva*. In *Lota*, as described and figured by Hyrtl (7. pl. i. fig. 2), there is a tendency for the last two efferent branchial vessels to unite before opening into the circulus. The fact that the cœliaco-mesenteric artery arises from the circulus is shown in Hyrtl's figure, and is quoted by Stannius (22. p. 103, footnote 3) and by Owen (19. p. 490).

In *Blennius* (fig. 35) the circulus cephalicus is narrow in front, the cœliaco-mesenteric artery arises from the aorta immediately behind the circulus, and the subclavian arteries are just behind this. In *Orthogoriscus* (fig. 36) the aorta is unsymmetrical, running to the right side of the vertebræ; the cœliaco-mesenteric artery is formed by the union of a pair of vessels arising from the common trunks formed on each side by the confluence of the third and fourth efferent branchial vessels. The mesial ends of all the efferent vessels are closely approximated. The gills in *Orthogoriscus* are remarkably prolonged in a backward direction. The efferent branchial vessel of each emerges from near the middle of the full extent of the gill, and not, as is more usual, from the upper end. It is formed by the union of one vessel coming from the lower part or gill proper, with another from the dorso-posterior prolongation, in a manner already made clear by Alessandrini (2) and Milne-Edwards (15. p. 335, footnote 2). Judging from Alessandrini's description, the peculiar mode of formation of the cœliaco-mesenteric artery in *O. truncatus* does not occur in the species examined by him (*O. mola*).

In *Tetrodon* (fig. 37) there are only three gills on each side, and three efferent branchial vessels, the fourth of the normal series being absent. The reason for putting the genus in the subgroup Dc, characterized by the union of the third and fourth efferent vessels into a common trunk, is to be found in the relation of the cœliaco-mesenteric artery to the third efferent vessel of the right side. The association is such that, were the fourth vessel present, it could not reach the circulus between the third vessel and the cœliaco-mesenteric. Both the right and left subclavian arteries arise from the right side of the circulus, and the latter is nearly circular in shape.

CONCLUSIONS.

It will be seen from the foregoing observations that very considerable diversity in the arrangement of the efferent branchial blood-vessels is to be met with in the Teleostean fishes. The type of vascular arrangement is constant for different species of the same genus, and does not vary to any considerable extent in different genera of the same family. If, as in the Siluroïd fishes, some widely divergent types are included in the same family, there are to be found intermediate types which act as connecting-links between these extremes.

With regard, however, to families which, in the at present accepted taxonomy of the group, are brought into close relationship,

the characters of the efferent branchial system fail to afford any convincing evidence as to the correctness or the reverse of such association. Thus, while the Gadoids have one sharply marked type of arterial arrangement, the Clupeoids another, and the Salmonoids a third, in the disposition of the efferent vessels the Lophobranchii resemble the first, the Ophidiidæ the second, and the Scombridæ the last,—a most unnatural coupling of the families. It does not appear, therefore, that the facts revealed by the present inquiry can be applied with any probability of success to the interpretation of the affinities of the family and larger groups of Teleostei.

LIST OF REFERENCES.

1. AGASSIZ, L., et VOGT, C.—“Anat. des Salmones,” Mém. de la Soc. des Sci. Nat. de Neuchâtel, iii. 1845 (1846), pp. 196, fourteen plates.
2. ALESSANDRINI, A.—“De piscium apparatu respir., tum speciatim Orthagorisco,” Nov. Comment. Acad. Sci. Bononiensis, iii. 1839, pp. 259–381, four plates.
3. ALLIS, E. P.—“The Cranial Muscles and Cranial and First Spinal Nerves in *Amia calva*,” Journal of Morphology, xii. Boston, 1897, pp. 487–808, nineteen plates.
4. AYERS, H.—“The Morphology of the Carotids, based on a Study of the Blood-vessels of *Chlamydoselachus anguineus*,” Bull. Mus. Comp. Zool. Harvard Coll., xvii., Cambridge, Mass., 1889, pp. 191–223, one plate.
5. CUVIER et VALENCIENNES.—Histoire Naturelle des Poissons. Paris, 1828.
6. DUVERNEY.—Oeuvres Anatomiques, ii. Paris, 1761.
7. HYRTL, J.—“Beob. aus dem Gebiete der vergl. Gefässlehre,” Med. Jahrb. der Oesterr. Staates, xxiv. (n. F. xv.), Wien, 1838, pp. 69–91 and pp. 232–248, four plates.
8. HYRTL, J.—“Über das Arterien-System des *Lepidosteus*,” Sitzungsber. Akad. Wiss. Wien, viii., 1852, pp. 234–241.
9. HYRTL, J.—“Beitrag zur Anat. von *Heterotis ehrenbergii*,” Denkschr. Akad. Wiss. Wien, viii., 1854, pp. 73–88, three plates.
10. HYRTL, J.—“Zur Anat. von *Saccobranchus singio*,” Sitzungsber. Akad. Wiss. Wien, xi., 1854, pp. 302–307, one plate.
11. HYRTL, J.—“Anat. Mitth. über *Mormyrus* und *Gymnarchus*,” Denkschr. Akad. Wiss. Wien, xii., 1856, pp. 1–22, six plates.
12. HYRTL, J.—Über besondere Eigenthümlichkeiten der Kiemen und des Skeletes von *Lutodeira chamos*,” Denkschr. Akad. Wiss. Wien, xxi., 1863, pp. 1–10, one plate.
13. MAURER, F.—“Ein Beitrag zur Kenntniss der Pseudobranchien der Knochenfische,” Morph. Jahrb., ix., Leipzig, 1884, pp. 229–252, two plates.
14. MECKEL, J. F.—System der vergl. Anat., v. Halle, 1831.

15. MILNE-EDWARDS, H.—Leçons sur la Phys. et l'Anat. comp., iii. Paris, 1858.
16. MÜLLER, JOH.—"Vergl. Anat. der Myxinoiden, Dritte Fortsetzung," Abhandl. Akad. Wiss. Berlin, 1839 (1841).
17. MÜLLER, JOH.—Über den Bau und die Grenzen der Ganoiden, und über das natürliche System der Fische. Berlin, 1846.
18. OWEN, R.—Lectures on Comp Anat., i., Fishes. London, 1846.
19. OWEN, R.—Anatomy of Vertebrates, i. London, 1866.
20. PARKER, T. J.—Zootomy. London, 1884.
21. ROLLESTON, G.—Forms of Animal Life: 2nd Edn. by W. Hatchett Jackson. Oxford, 1888.
22. STANNIUS, H.—Lehrb. der vergl. Anat., ii., Wirbelth. Berlin, 1846.
23. STANNIUS, H.—Das periph. Nervensystem der Fische. Rostock, 1849.
24. STANNIUS, H.—Handbuch der Anat. der Wirbelthiere, i., Fische. Berlin, 1854.
25. WRIGHT, R. R.—"On the Hyomandibular Clefts and Pseudo-branches of *Lepidosteus* and *Amia*," Journ. Anat. and Phys. xix., London, 1885, pp. 476-499, one plate.

EXPLANATION OF THE PLATES.

Efferent Branchial Blood-vessels of the first, second, third, and fourth Gills of Teleostean Fishes, as seen after stripping off the mucous membrane from the roof of the pharynx.

The dorsal aorta, the cœliaco-mesenteric artery, and the subclavian artery are respectively indicated in all the figures by the letters, *a*, *c*, and *s*.

PLATE LXIII.

- Fig. 1. *Engraulis encrasicolus*, p. 947.
 2. *Clupea harengus*, p. 947.
 3. *Anmodytes lanceolatus*, p. 947.
 4. *Megalops cyprinoides*, p. 947.
 5. *Balistes aculeatus*, p. 948.
 6. *Scomber scombrus*, p. 948.
 7. *Salmo salar*, p. 948.
 8. *Mugil capito*, p. 948.
 9. *Hippoglossus vulgaris*, p. 948.
 10. *Osmerus eperlanus*, p. 949.
 11. *Albula conorhynchus*, p. 949.
 12. *Malapterurus electricus*, p. 949.
 13. *Cyprinus carpio*, p. 949.

PLATE LXIV.

- Fig. 14. *Cobitis tania*, p. 949.
 15. *Esox lucius*, p. 949.
 16. *Anguilla vulgaris*, p. 949.
 17. *Labrax lupus*, p. 950.
 18. *Liocassis longirostris*, p. 950.
 19. *Callichthys littoralis*, p. 950.
 20. *Clarias magur*, p. 950.

- Fig. 21. *Perca fluviatilis*, p. 950.
 22. *Zeus faber*, p. 950.
 23. *Anarrhichas lupus*, p. 950.
 24. *Labrus maculatus*, p. 951.
 25. *Eroctatus brachysoma*, p. 951.

PLATE LXV.

- Fig. 26. *Lophius piscatorius*, p. 951.
 27. *Cottus scorpio*, p. 951.
 28. *Trigla cuculus*, p. 951.
 29. *Cyclopterus lumpus*, p. 951.
 30. *Fistularia tabaccaria*, p. 952.
 31. *Saccobranchius fossilis*, p. 952.
 32. *Silurus glanis*, p. 952.
 33. *Syngnathus acus*, p. 952.
 34. *Gadus æglenus*, p. 952.
 35. *Blennius pholis*, p. 953.
 36. *Orthogoriscus truncatus*, p. 953.
 37. *Tetrodon palembangensis*, p. 953.

4. On the Reptiles, Batrachians, and Fishes collected by the late Mr. John Whitehead in the Interior of Hainan. By G. A. BOULENGER, F.R.S.

[Received October 31, 1899.]

(Plates LXVI.-LXIX.)

During his short stay in Hainan, where he died on June 2 of the present year, Mr. John Whitehead had succeeded in collecting a small number of cold-blooded vertebrates in the Five-finger Mountains, in the interior of the island. The fact that so many of the few species represented in the collection are new, tends to show how rich a harvest these unexplored mountains would have yielded but for the fatal climate which has deprived the zoological world of one of its most enthusiastic and successful members.

REPTILES.

1. DRACO WHITEHEADI, sp. n. (Plate LXVI. fig. 1.)

Head small; snout considerably longer than the diameter of the orbit; nostril lateral, directed outwards; tympanum scaly. Upper head-scales unequal, strongly keeled; 8 or 9 upper labials. Male's gular appendage very large, once and a half as long as the head. A rudimentary nuchal crest. Dorsal scales a little larger than ventrals, irregular, obtusely keeled; on each side of the back a series of enlarged, keeled dorsal scales. The fore limb stretched forward reaches the tip of the snout, the hind limb between the elbow and the axilla. Reddish brown above, with dark transverse

1.

2.

P.J.Smit del. et lith.

Mintern Bros. imp.

1. DRACO WHITEHEADI.

2. ACANTHOSAURA HAINANENSIS.

1.

1a.

2a.

3.

2.

Mintern Bros. imp.

COREOPERCA WHITEHEADI

J. Green del. et lith.

2.

J. Green, del. et lith.

1. GYMNOSTOMUS LEPTURUS.

2. BARILIUS HAINANENSIS.

Macleay, Proc. Linn. Soc. New South Wales, vol. 20, p. 10, 1899.

phractus, a Python, 3 *Cyclanorbis senegalensis*, 2 Hinged Tortoises, some Chameleons, and a Serval Cat were brought home alive.

Since my return to England, I have definitely decided that the eggs and larvæ obtained are not those of *Polypterus*. I have, however, I believe, learned enough about the habits of *Polypterus* to encourage me to make a second attempt next year to obtain the developmental stages.

In conclusion, I wish to thank the Society for lending me influence and support, without which the little that has been done by this expedition could not have been accomplished.

2. A Note on the Hatching-stage of the Pagurine Land-crabs. By L. A. BORRADAILE, M.A., F.Z.S., Lecturer in Natural Sciences of Selwyn College, Cambridge.

[Received October 12, 1899.]

The life-history of the Land-crabs of the family Cœnobitidæ is one to which considerable interest attaches, and of which, at present, nothing appears to be known. The family comprises the genera *Birgus* and *Cœnobita*, the robber- or coconut-crab and the land hermit-crabs, all of which have given up a sea life for one on land. It need hardly be remarked that changes in habitat, particularly from sea to land or fresh water, have frequently necessitated the suppression of larval stages in the life-history. Among Crustaceans the instances of the cray-fishes, the ditch-prawn (*Palæmonetes varians*), the freshwater crabs (*Potamon*), and at least one species of land-crab (*Gecarcinus*) come at once to mind. The possibility was thus suggested that the land-pagurines might also have lost the whole or a part of their larval life, and leave the egg in something like the adult condition. On the other hand, it had to be borne in mind that some species of land-crabs and all the strand-crabs (*Ocypoda*, &c.) retain the habit of setting free zoœa-larvæ in the sea, where they pass through their earlier stages.

It was probably with these considerations in his mind that von Willemoës-Suhm, when, in October 1874, the 'Challenger' arrived at Zamboanga in the Philippine Islands, wished to investigate the development of the robber-crab from the egg. Unfortunately the time of year rendered it impossible for him to do this, but he was told by an "intelligent native" that the young were born resembling the parent. This statement has since been accepted in a tentative manner by certain text-books, in spite of the fact that the small size of the eggs (and, indeed, of the female genital opening) made it improbable that the development was a direct one, depending on food-yolk. As for the statement of the

BATRACHIANS.

1. *RANA GRAMINEA*, sp. n. (Plate LXVII. fig. 1.)

Vomerine teeth in two short oblique series between the choanæ, nearer to each other than to the latter. Head depressed, as long as broad; snout rounded, scarcely projecting, as long as the diameter of the orbit; canthus rostralis well-marked; loreal region concave; nostril nearer the end of the snout than the eye: interorbital space as broad as the upper eyelid; tympanum very distinct, three fourths the diameter of the eye. Fingers and toes rather slender, with small but well-developed disks; first finger not extending beyond second; toes nearly entirely webbed; a single, feebly prominent, oval, inner metatarsal tubercle. The tibio-tarsal articulation reaches beyond the tip of the snout; tibia as long as the distance from end of snout to sacrum. Skin smooth; a moderately broad, feebly prominent glandular lateral fold; another fold from below the eye to the shoulder, followed by a strong glandule. Bright green above, brownish on the sides of the head and body, below the canthus rostralis and the dorso-lateral fold, and on the limbs; upper lip white; limbs with regular dark cross-bars; hinder side of thighs marbled dark brown and yellow; lower parts white. Male with two external vocal sacs, in front of the arms; no humeral gland.

From snout to vent 48 millim.

Allied to *R. erythræa* Schleg. Distinguished by the shorter snout, the longer hind limbs, the external vocal sacs, and the coloration. Also allied to *R. jerboa* Gthr. and *R. whiteheadi* Blgr., in which the digital disks are larger and the hind limb longer still.

Two male specimens.

2. *RANA ANDERSONI* Blgr.

This species, first discovered in the Hotha Valley, Yunnan, by Dr. J. Anderson, has since been found in the Kakhien Hills, Upper Burma, by Signor L. Fea, and at Kuatun, N.W. Fokien, by Mr. J. D. La Touche.

3. *STAUROIS HAINANENSIS*, sp. n. (Plate LXVII. fig. 2.)

Head as long as broad or slightly broader than long; snout short, truncate, projecting; canthus rostralis strong; loreal region nearly vertical, concave; nostril midway between the eye and the end of the snout; interorbital space as broad as the upper eyelid; tympanum distinct, one third or two fifths the diameter of the eye. Fingers slender, first longer than second, with very large disks; toes webbed to the disks, which are a little smaller than those of the fingers; subarticular tubercles feebly prominent; a very indistinct inner metatarsal tubercle. The tibio-tarsal articulation reaches the tip of the snout or a little beyond. Skin smooth above in the adult, warty in the young; lower part

smooth. Olive above, spotted with black, or blackish with pale olive markings; limbs with dark cross-bars; hinder side of thighs with a black reticulation.

From snout to vent 58 millim.

Larva with a large pectoral adhesive disk (see P. Z. S. 1893, p. 526). Beak formed of two pieces, an upper and a lower, feebly denticulate, not ribbed; lower lip not fringed; the horny teeth form 3 uninterrupted and 2 paired series on the upper lip, 2 uninterrupted and 1 narrowly interrupted series on the lower lip,

an arrangement that may be expressed by the formula $\frac{2}{1} \frac{2}{1}$.

Closely allied to *Staurois natator* Gthr. Distinguished by the shorter head. The tadpole, on the other hand, stands nearest to that of *Rana latopalmata* Blgr.

Three specimens: a female, a young, and an advanced tadpole.

4. RHACOPHOBUS LEUCOMYSTAX Gravh.

5. RHACOPHORUS OXYCEPHALUS, sp. n. (Plate LXVII. fig. 3.)

Vomerine teeth in two oblique series between the choanæ, the inner front edge of which they nearly touch. Head as long as broad; snout pointed, as long as or a little longer than the diameter of the orbit; canthus rostralis distinct; loreal region concave; nostril a little nearer the tip of the snout than the eye; inter-orbital space a little narrower than the upper eyelid; tympanum distinct, half the diameter of the eye. Fingers with a distinct rudiment of web; toes entirely webbed; disks of fingers nearly as large as the tympanum, of toes a little smaller; a very small inner metatarsal tubercle. The tibio-tarsal articulation reaches beyond the tip of the snout. Skin smooth or with small warts above; belly granular. Greyish or brown above, spotted or marbled with darker; a dark transverse band or triangular marking, base forwards, between the eyes; limbs with dark cross-bars; groin and back of thighs marbled black and yellow; lower parts white. Male with an internal vocal sac.

From snout to vent 57 millim.

Four specimens.

6. BUFO MELANOSTICTUS Schn.

FISHES.

COREOPERCA.

Coreoperca, Herzenstein, Ann. Mus. Zool. St. Pétersb. 1896, p. 11.

Body compressed; scales small, cycloid, concentrically striated, lateral line complete; tubes straight, occupying the greater length of the scale. Mouth large, protractile; maxillary exposed, with

supplemental bone; villiform teeth in jaws and on vomer and palatines; no canines; tongue smooth; head partly naked; præopercle serrated, with a few antrorse spines on the lower border; opercle with two spines. Gill-membranes separate; seven branchiostegals; pseudobranchiæ present. Dorsal fins confluent, XIV-XV 11-14, the spinous portion much longer than the soft; anal short, III 7-11; caudal rounded. Pectoral symmetrical, rounded, rays 16. Ventrals below the pectorals, close together, with a strong spine and five branched rays, the last of which is connected with the belly by a membrane.

The type species of this genus of *Serranidae*, allied to *Siniperca*, is from the interior of North Corea¹. It is highly interesting to add a second species from the interior of Hainan.

1. COREOPERCA WHITEHEADI, sp. n. (Plate LXV.III.)

Depth of body equal to length of head, 3 times in total length. Snout $1\frac{1}{3}$ diameter of eye, which equals interorbital width, $\frac{1}{4}$ length of head; lower jaw projecting beyond the snout; maxillary extending a little beyond vertical of posterior border of eye, the width of its distal extremity a little less than diameter of eye; præorbital entire; cheeks and opercles scaly, rest of head naked; præopercle finely serrated, without enlarged spines at the angle; opercular spines strong. Dorsal XV 14, originating above base of pectoral; spinous portion twice as long as the soft; spines strong, short, increasing in length to the sixth, which equals $\frac{1}{4}$ length of head; longest soft rays nearly $\frac{1}{2}$ length of head. Pectoral $\frac{1}{2}$ length of head. Anal III 11; second spine longest, a little shorter than longest dorsal spines. Caudal rounded, subtruncate. Sq. 80 $\frac{14}{31}$; l.l. 63. Brown, with dark marblings and whitish dots; a dark streak from below the eye to the angle of the præopercle and another from the eye to a large, black, white-edged ocellar spot

¹ Having had the privilege of examining the type specimens of *Coreoperca herzi*, Herzenst. l. c., preserved in the St. Petersburg Museum, I add a description of them for comparison with *C. whiteheadi*:—

Greatest depth at origin of dorsal fin, 3 to $3\frac{1}{4}$ times in total length, length of head $2\frac{1}{2}$ to 3 times. Snout as long as diameter of eye, $\frac{1}{4}$ length of head, and twice width of interorbital region; lower jaw not projecting; maxillary extending to below posterior third of eye, the width of its distal extremity about half diameter of eye; præorbital entire; cheeks and opercles scaly, rest of head naked; præopercle with two strong bifid spines at the angle and two or three antrorse spines on the lower border; opercular spines strong. Dorsal XIV 11-12; originating above base of pectoral, spinous portion twice as long as the soft; spines strong, increasing in length to the fifth, which equals $\frac{2}{3}$ length of head, but is considerably shorter than the longest soft rays. Pectoral $\frac{2}{3}$ length of head. Anal III 7, spines very strong, third a little longer than first, second longest and a little longer than longest dorsal spine. Caudal rounded. Sq. 76-82 $\frac{14-15}{35}$; l.l. 51-56. Brown; a dark streak from below the eye to the angle of the præopercle; a black spot, edged with white anteriorly, between the opercular spines; body with some dark brown spots intermixed with whitish dots; a regular series of dark spots along the base of the dorsal.

Total length 85 millim.

Pung Tung, Corea.

between the opercular spines; some light dots on the soft dorsal and anal and on the membrane between the ventral rays.

Total length 155 millim.

A single specimen.

2. *DISCOGNATHUS IMBERBIS* Vincig.

A species described from the Karen Hills, Burma.

3. *GYMNOSTOMUS LEPTURUS*, sp. n. (Plate LXIX. fig. 1.)

Depth of body 4 times in total length, length of head 5 times. Head $1\frac{2}{3}$ as long as broad; snout broad, rounded; width of mouth nearly half length of head; lower jaw with a sharp, horny edge; diameter of eye equal to length of snout, $3\frac{1}{2}$ times in length of head, $1\frac{1}{2}$ in interorbital width. Dorsal III 8, midway between end of snout and base of caudal; first branched ray longest, a little shorter than the head, last ray longer than those preceding it, $\frac{2}{3}$ length of head. Pectoral as long as head. Ventrals below middle of dorsal. Anal III 6, as deep as dorsal. Caudal deeply bifurcate, $1\frac{1}{2}$ length of head. Caudal peduncle thrice as long as deep. Sq. $49\frac{7}{8}$; 4 scales between the lateral line and the ventral. Olive above, silvery beneath; an ill-defined dark lateral streak.

Total length 165 millim.

A single specimen.

4. *BARILIUS HAINANENSIS*, sp. n. (Plate LXIX. fig. 2.)

Depth of body equal to length of head, $4\frac{2}{3}$ times in total length. Head twice as long as broad; snout pointed, not projecting beyond the mouth, as long as diameter of eye, which is $3\frac{1}{2}$ times in length of head and equals interorbital width; mouth extending hardly to below anterior border of eye; suborbitals entirely covering the cheek. Dorsal II 7, originating just behind ventral and situated at equal distance from the eye and the root of the caudal; first branched ray $\frac{2}{3}$ length of head. Pectoral a little shorter than head, not reaching ventral. Anal II 14. Caudal deeply bifurcate, as long as head. Caudal peduncle nearly thrice as long as deep. Sq. $46\frac{6}{3}$. Silvery, darker on the back; scales above the lateral line black at the base.

Total length 130 millim.

A single specimen.

5. *OPSARIICHTHYS PLATYPUS* Schleg.

A species known from Japan and Formosa.

EXPLANATION OF THE PLATES.

PLATE LXVI.

Fig. 1. *Draco whiteheadi*, p. 956, with side-view of head.

2. *Acanthosaura hainanensis*, p. 957.

PLATE LXVII.

Fig. 1. *Rana graminea*, p. 958.

1 a. " " Side view of head.

2. *Staurois hainanensis*, p. 958.

2 a. " " larva, lower view of body.

3. *Rhacophorus oxycephalus*, p. 959.

PLATE LXVIII.

Coreoperca whiteheadi, p. 960.

PLATE LXIX.

Fig. 1. *Gymnostomus lepturus*, p. 961, $\frac{2}{3}$ nat. size.2. *Barilius hainanensis*, p. 961.

5. On a Collection of Butterflies made by Mr. Richard Crawshay in British East Africa. By ARTHUR G. BUTLER, Ph.D., F.L.S., F.Z.S., &c., Senior Assistant-Keeper, Zoological Department, British Museum.

[Received September 20, 1899.]

(Plate LXX.)

During the past summer I received from Mr. Crawshay a box of Lepidoptera and a letter dated February 8th, 1899, addressed from Neugia, as follows:—

“A few lines to let you know that I have lately returned from a journey into Maraṅga, the S. and S.W. slopes of Mt. Kenya; and that I was able to take some Butterflies, which, I think, will please you.

“From this—I mean the mention of mighty Kenya and its 18,600 feet—you must not infer that these insects have been collected at any great altitude. Maraṅga is not so high as other parts of Kikuyu to the westward,—for instance the neighbourhood of Fort Smith, which is 6400 feet. As a matter of fact it does not average, I suppose, more than 5600 feet; rising to the N. and N.E. gradually into the mighty belt of forest surrounding the mountain for many thousands of feet, and falling away to the W. and S.W. of the Tana River, which, where we crossed it, is 3850–3900 feet.

“In all the thousands of miles I have travelled in Africa, I have never seen a more lovely and more *possible* country than Maraṅga; nor more splendid specimens of its peoples than are the Wakikuyu—though they are at present suspicious of and hostile to everyone from the outer world. The Wakamba are the most veritable *worms* in comparison with them.

“However, you want to know something more of the surprises which I hope are in store for you in the shape of the Butterflies.

“The most promising of these are Skippers and Blues—one a very large and powerful Blue with almost black wings on the inside, which show a Purple-Emperor-like glow, though with a

F.W.Frohawke del. et lith.

West, Newman chromo.

BUTTERFLIES FROM BRIT. E. AFRICA.

tinge of port-wine colour about it¹. This Blue (of which I think I took a pair) and another very active and cunning insect—as you will see from the note appended on the paper envelope—were taken on a stinking crocodile's head, than which nothing smells more foully; otherwise—I mean without this—I should probably never have seen the former, nor taken either of them.

“A hippopotamus skull far gone in putrefaction also proved attractive, and did me several good turns in enabling me to get on equal terms with other insects: on this I took the only *Charaxes* I saw, I think, on the whole journey, though one I know well. Of Moths I also took a nice lot.”

On his return Mr. Crawshaw had a narrow escape from a wounded bull Rhinoceros, but by a fortunate shot whilst lying in the grass he managed to hit it in the heart, and so escaped with his life.

The collection of Moths will be worked out by Sir George Hampson. The Butterflies are represented by 127 specimens referable to 69 species, of which 3 are new to science; the most interesting to me is a new species of *Chloroselas* allied to *C. tamaniba* of Walker from Suakin, but apparently distinct, and clearly proving that Prof. Aurivillius was in error in referring the Somali insect *C. esmeralda* to that species. *C. tamaniba* is considerably larger than *C. esmeralda*, has the posterior half of the primaries, including the base, blue; and the anal orange spot of the secondaries is well-defined as in the present species.

N Y M P H A L I D Æ.

1. MONOTRICTIS SAFITZA Hewits.

Muthambi River, Ndya, 7th January, 1899.

“They swarm on the banks of the stream” (*R. C.*).

2. NEOCENYRA DUPLEX Butl.

Slopes of Nthatha Hill, Kitwi, 4700 feet, 31st December, 1898.

3. NEOCENYRA GREGORII Butl.

Muthambi River, Ndya, 7th January, 1899.

One much worn example.

4. CHARAXES VARANES Cram.

♂, Muthambi River, Ndya, 13th January, 1899.

“Taken on the crocodile's head (hippopotamus' skull?); the only one of this species which I have seen during my week here” (*R. C.*).

5. PRECIS CLOANTHA Cram.

♂, Plains N. of the Tana River, Kikuyu, 4500 feet, 5th January; ♀, Muthambi River, Ndya, 12th January, 1899.

¹ This proves to be a well-known species of *Crenis*, which occurs also in South Africa.—A. G. B.

The female, though in excellent condition, was caught by Bvalamkombi, one of Mr. Crawshaw's native servants.

6. *PRECIS ELGIVA* Hewits.

♂, ♀, Muthambi River, Ndya, 7th January, 1899.

7. *PRECIS NATALICA* Feld.

♀, Muthambi River, Ndya, 13th January, 1899.

"Fairly plentiful: but this is, I think, the only perfect specimen I have seen" (*R. C.*).

8. *HYPOLIMNAS MISIPPUS*, var. *INARIA* Cram.

♀, Muthambi River, 4500 feet, Ndya, 11th January, 1899.

9. *CRENIS BOISDUVALII* Wilgr.

♂ ♂, Muthambi River, 4500 feet, Ndya, 12th January, 1899.

"An insect which I should probably never have seen, and certainly not taken, had it not been for the putrefying crocodile's head, on which it descended as it were from the clouds. It is worthy of note that this insect shows a lovely tinge of purple or blue on the inside of the wings if these are viewed at an angle in a good light." (*R. C.*)

10. *CRENIS HOWENSIS* Staud.

♂ ♂, Muthambi River, 12th January, 1899.

"The most restless, active, and difficult insect to take I have ever come across; for six days have I been watching and following the movements of some half-dozen in the glade of fig-trees in which I am encamped, trying all I could to take one, but without success. They flip about amongst the trees and occasionally perch on the trunk and branches, but always out of reach. By great good luck this morning two have been tempted to come down and feed on a putrefying crocodile's head, and I was thus enabled to take them." (*R. C.*)

One specimen (a male) is said to have contained a whitish-green ovum, but this must have been an error of observation, as the insect has strongly pronounced male claspers.

11. *HAMANUMIDA DÆDALUS* Fabr.

"Plentiful enough on undulating plains N. of Tana River, 4200 feet, 14th January, 1899."

A single dry-season male of this abundant species.

12. *NEPTIS AGATHA* Cram.

♀, ♂, Muthambi River, 4500 feet, Ndya, Kikuyu, 6th & 7th January, 1899.

13. *NEPTIDOPSIS OPHIONE*, var. *VELLEDA* Mab.

Muthambi River, 4500 feet, Ndya, 6th & 13th January, 1899.

Mr. Crawshaw appears to have seen three examples of this

species on the 6th January, of which he caught two then, and the third a week later.

14. *BYBLIA ILIYTHIA* Drury.

Wet phase.—Machakos to Neugia, 16th December, 1898.

Intermediate and dry phases.—Neugia, 31st January, 1899.

“Quite the commonest butterfly hereabouts these days; indeed I have never anywhere else seen this insect in such numbers” (*R. C.*).

15. *ACRÆA CABIRA* Hopff.

Clue to exact locality and date lost.

An example leading to the variety *A. apocida*.

16. *ACRÆA LYCIA* Fabr. (vars. *sganzini* and *daira*).

Muthambi River, Ndya, 7th & 11th January, 1899.

17. *ACRÆA ONERATA* Trim.

♂ ♀ *in copula*, Neugia, Kitwi, 11th February, 1899.

The sexes are remarkably alike, the wing-borders of the female slightly heavier and the body spotted.

18. *ACRÆA ACRITA*, var. *PUDORINA* Staud.

♂ ♂, Neugia, Kitwi, 16th February, 1899.

LYCÆNIDÆ.

19. *LACHNOCNEMA BIBULUS* Fabr.

Ndya, Kikuyu, 4500 feet, 6th January, 1899.

20. *AXIO CERSEES HARPA* Fabr.

♀, Kitwi, 18th January, 1899.

“Grass-green ova” (*R. C.*).

21. *TERIOMIMA PALLIDA* Trim.

E. of Athi River, about 4300 feet, 18th December, 1898.

Mr. Crawshay regarded this as a worn example of *T. hildegarda*, to which it certainly is very closely related, but the small and inconspicuous spots on the under surface of the secondaries give it a somewhat different aspect; it would not surprise me to find that it was only another variety of that variable species. Mr. Crawshay says he took it for a moth until it was in the killing-bottle.

22. *CATOCHRYSOPE PECULIARIS*, var., Rogenh. (Plate LXX. fig. 1.)

♀ ♀, Neugia, Kitwi, 24th & 29th December, 1898.

Of the first example Mr. Crawshay writes—“A hardly won capture. Caught in my Terai hat when out shooting, and kept under this on the ground, while I covered my head from the

blazing sun until my net reached me, fully three quarters of an hour."

The two specimens are of much interest, as they clear up one of the greatest muddles which has been made over any species of *Lycaenidæ*:—In 1891 Rogenhofer described a butterfly which was quite unknown to English entomologists under the name of "*Chrysophanus peculiaris*"; naturally nobody expected a *Catochrysops* to be called a *Chrysophanus*. In 1892 Dr. Holland described the same species in the 'Entomologist' under the name of *Lycaena perpulchra*; in 1893 I described a male from Wasin and a female from the Victoria Nyanza as *Castalius hypoleucus*; and in 1894 Mr. Trimen described *C. peculiaris* again under the name of *Lycaena exclusa*. Dr. Holland subsequently pointed out that *C. hypoleucus* and *L. exclusa* were synonymous with his *L. perpulchra*. In 1898 Mr. Trimen received some large examples from Mashunaland which he rightly stated to be identical with our Nyanza female; but instead of adopting my name for this form, he called it *Lycaena gigantea*, stating that I had confounded the female with that sex of *L. perpulchra* (entirely overlooking the fact that the Nyanza female was described by me as the type of that sex of *Castalius hypoleucus*). At the commencement of the present year, when working out the species of *Chrysophanus*, I recognized *C. peculiaris* Rogenh. as the oldest name for the present species, and entered it in my paper on Mr. Crawshay's last collection. Professor Aurivillius also recognized Rogenhofer's species in his '*Rhopalocera Æthiopica*,' where, however, he retained Mr. Trimen's name for the larger form, ignoring the fact that my female unquestionably takes priority.

The two examples now sent home by Mr. Crawshay are quite intermediate in character between the large and small forms, the colouring of the upper surface agreeing most nearly with *C. peculiaris* ♀, the expanse of wing being nevertheless equal to that of my Victoria Nyanza female. Like the latter and a worn and faded female from Zomba (which Trimen refers to as the Nyasa female), they have from one to two extra black spots in the discal series on the under surface of the primaries; the black discocellular spot is smaller than in the little Zomba female, but varies in size in the two examples. It is absolutely impossible to say that these examples belong to one form rather than to the other, and I do not doubt that they represent an intermediate phase between *C. hypoleucus*=*gigantea* the wet phase, and *C. peculiaris* the dry phase, of one and the same species.

23. *AZANUS NATALENSIS* Trim.

♂, Tana River, 3800 feet, 16th January, 1899.

"The only specimen seen" (*R. C.*).

24. *AZANUS ZENA* Moore (*Lycaena maculenga*, Trim.).

♂, Kitwi, 4000 feet, 30th December, 1898.

25. *EVERES KEDONGA* Grose-Smith. (Plate LXX. fig. 4.)

♂ ♂, Plains N. of the Tana River, Kikuyu, 5th January, 1899.

Mr. Crawshay sent us the female of this extremely pretty species in his last collection (see P. Z. S. 1899, pl. xiv. figs. 3, 3a).

26. *TARUCUS PLINIUS* Fabr.

♀, Athi Valley, 16th December, 1898; ♂, Ndya, 4500 feet, Kikuyu, 16th January, 1899.

Of the female Mr. Crawshay observes—"Common, and all evidently newly emerged."

27. *ZIZERA KNYSNA* Trim.

♀, Slopes of Nthatha Hill, 4700 feet, Kitwi, 31st December, 1898; ♂, ♀, Muthambi River, 4500 feet, Ndya, 8th & 10th January, 1899.

28. *ZIZERA GAIKA* Trim.

♂ ♂, Tana River, 3800 feet, 2nd January; Neugia, Kitwi, 7th February, 1899.

Of the second example Mr. Crawshay observes—"The smallest butterfly I have ever seen." It is a starved specimen.

29. *LYCÆNESTHES AMARAH* Leféb.

♀, Slopes of Nthatha Hill, Kitwi, 4700 feet, 31st December, 1898; ♂ ♂, Plains N. of the Tana River, 5th January, Tana River, 3800 feet, 16th January, 1899.

30. *CACYREUS LINGEUS* Cram.

♂ ♂, Clue to exact locality and date lost.

31. *PHLYARIA VIRGO* Butl.

♀, Muthambi River, 4500 feet, Ndya, 10 January, 1899.

"The only specimen I have seen in these parts, and like, if not identical with, another which I used to take in the mountains of Nyika" (*R. C.*).

According to Prof. Aurivillius this is the female of the West Coast *P. heritsia*, but some of the details of marking make me hesitate to accept this dictum until I have seen East-African males. In my opinion the border of the secondaries is too narrow and the markings on the under surface of these wings too feeble for *P. heritsia*: I may prove to be wrong, but I very strongly object to putting species together by guess.

32. *CHLOROSELAS AZUREA*, sp. n. (Plate LXX. figs. 2, 3.)

Intermediate between *C. tamaniba* from Suakin and *C. esmeralda* from Somali-land; evidently nearest to the former. It is considerably larger than *C. esmeralda*, the shot-colouring on the upperside of the wings being glistening deep sky-blue instead of emerald shaded with blue: in the primaries this colouring is restricted to the outer half of the internal area and the internal margin nearly

to base; the remainder of these wings is smoky brown, slightly cupreous, with a large ill-defined but distinctly darker patch over the end of the cell: secondaries cupreous brown, brilliantly glossed with sky-blue between the second subcostal branch and submedian vein; a conspicuous orange anal spot between two short tails: anal lobe small, silver-spotted and fringed with black; body blackish; frons silvery whitish, collar with pale edges; antennæ annulated with white, the club externally edged with tawny: under surface pale fleshy buff, with the discoidal area of the primaries and the anal patch of secondaries orange-ochreous; the internal area of the primaries blackish grey towards the base, white slightly opalescent beyond, the marking on the wings much as in *C. esmeralda*, silver with black margins; body below white. Expanse of wings 26 millim.

The female is rich copper-brown with white fringes, and an orange spot on the secondaries between the tails, as in the male; the under surface similar to that of the male: size uniform.

♂, Slopes of Nthatha Hill, Kitwi, 4700 feet, 31st December, 1898; ♀, Plains N. of the Tana River, 4500 feet, Kikuyu, 5th January, 1899.

C. tamaniba, according to Walker, expands one inch and one line (or 28 millim.), and is therefore the largest species in the genus; the primaries are described as being shot with blue at the base and on the hind half, and the secondaries as blue, with a narrow brown border; "an orange spot adjoining the tail, into which it extends, bordered on the outer side with glittering chalybeous." These characters do not at all correspond with those of *C. esmeralda* (which Prof. Aurivillius has unaccountably placed as a synonym of it), and differ considerably from those of the present species. Many years ago (1870) I saw the type, but I cannot pretend to remember exactly what it was like. I am, however, quite certain that Walker's measurement is rather under than overstated: the specimen is, of course, incorrectly described as a female. *C. esmeralda* not only differs from *C. pseudozeritis* in having only one tail to the secondaries, but in the absence of the brown clouding on the under surface of these wings. It is mere guesswork to suppose that a small insect like this from Natal is at all likely to be identical with one from Somali-land, or even that the latter is likely to be the same as one from the western shore of the Red Sea. It is quite possible that there are many species of *Chloroselas* scattered over Africa, and that, ten years hence, the species hitherto confounded will be generally regarded as amongst the best marked of all. Structurally, the species may at present be separated by the tails of the secondaries as follows:—

Species with two tails.

Small—*C. pseudozeritis* Trimen.

Large—*C. azurea* Butler.

Species with one tail.

Small—*C. esmeralda* Butler.

Large—*C. tamaniba* Walker.

But in many respects the two small species and the two large species are more nearly related to each other respectively in other characters—the bluer shot and the internal position of the blue colouring, as well as the more conspicuous anal orange spot of the secondaries, being characteristic of the larger species, the greener shot and the basal position of the greenish colouring, as well as the less conspicuous anal orange-tinted spot of the secondaries, of the smaller species.

33. *MYRINA FICEDULA* Trim.

♂, Neugia, Kitwi, 22nd December, 1898.

“Taken sitting on an outstanding branch of a small tree about noon in the hottest possible sun” (*R. C.*).

34. *VIRACHOLA ANTALUS* Trim.

♂, Neugia, Kitwi, 22nd January, 1899.

35. *VIRACHOLA DERONA*? Smith.

E. of Athi River, some 4300 feet, Kitwi, 18th December, 1898.

“I have no recollection of having taken this insect before. It was most accommodating too in awaiting about a quarter of an hour. I was stalking a Hartebeeste and my net was behind.” (*R. C.*)

This appears to correspond with Mr. Grose-Smith’s description, but it is a somewhat faded example.

PAPILIONIDÆ.

36. *MYLOTHRIS AGATHINA* Cram.

♂♂, Muthambi River, 4500 feet, Ndya, 7th & 10th January, 1899.

37. *NYCHITONA MEDUSA*, var. *ALCESTA* Cram.

♂, Tana River, 3850 feet, 4th January, 1899.

38. *COLIAS ELECTRA*, var. *EDUSA* Fabr.

Ndya, 8th January, 1899.

“The first specimen of *C. edusa* I have seen since leaving Massai” (*R. C.*).

39. *TERIAS SENEGALENSIS*, var. *BISINUATA* Butl.

♂, Tana River, 4th January, 1899.

40. *TERACOLUS CALAIS* Cram.

♂, Kitwi, 19th January, 1899.

“An average specimen of this species as regards size” (*R. C.*).

41. *TERACOLUS ERIS* Klug.

♂♂, Athi Valley, 4000 feet, 16th December, and Msokani, Kitwi, 20th December, 1898.

Of the first specimen Mr. Crawshay writes—“By no means

common: an active and exceedingly restless insect of strong flight—very difficult to take if once missed.” Of the second specimen, however, he writes—“ Newly emerged, evidently an easy prey in the wet and cold of early morning.” This, then, is clearly the time to secure it.

As might have been expected, the males are referable to the typical Northern species, not to the more southerly *T. opalescens*. To anyone with a correct eye for outline, the pattern of the primaries in these local forms is absolutely different, apart from all minor differences of colouring; but the training of a lifetime is insufficient to enable some men to appreciate the most marked modifications of outline.

42. *TERACOLUS INCRETUS* Butl.

Intermediate phase.—♀, Athi Valley, 4000 feet, 16th December; ♂ ♀, Athi escarpment, eastern side, Kitwi, 18th December; ♀, Msokani, 20th December, 1898; ♂, Tana River, 3800 feet, 16th January, 1899.

Dry phase.—♂, ♀ ♀, Tana River, 3800 feet, 16th January, 1899.

Mr. Crawshay says of the male—“ Very plentiful just here (Athi Valley): this and the following taken in numbers when playing together, with one stroke of the net.” Of the female he says—“ An insect I have never before taken, and which, until settled, I imagined to be another, rather common just here—plain sulphur with an orange tip.” In this conjecture Mr. Crawshay was quite correct.

43. *TERACOLUS XANTHUS*, var. *METAGONE* Holl.

♂, Tana River, 3800 feet, 16th January; ♀, Neugia, Kitwi, 30th January, 1899.

♂. “ Fairly common, frequents the more open country, dry and desert-like, and covered with thorny scrub ” (*R. C.*).

This is distinctly a dry-season phase, and differs from *T. xanthus* var. *comptus* in the entire absence of the internal grey streak on the primaries.

44. *TERACOLUS ANTEVIPPE* Boisd.

♂, var. *subvenosus*, Kitwi, 18th January, 1899.

45. *TERACOLUS GAVISA* Wallgr.

♀, Kitwi, 19th January, 1899.

46. *TERACOLUS CALLIDIA* Grose-Smith.

♂ ♂, *Intermediate and dry phases*.—Tana River, 3800 feet, 16th January, 1899.

“ Fairly plentiful; but, if once missed, by no means an easy insect to take ” (*R. C.*).

The typical orange form of this species is by no means too well represented in the Museum series,

47. *TERACOLUS CATACHRYSOPS* Butl.

♂, Msokani, Kitwi, 20th December, 1898.

"Fairly common, but not an easy insect to take: a fast flier and a desperate doubler" (*R. C.*).

The single example sent belongs to the wet phase, which differs from the dry phase in the better defined, browner, and much darker bands on the under surface. This character, apart from other differences, amply serves to prove its entire distinctness from *T. mutans* of Southern Africa, whilst at the same time it indicates some relationship to *T. protomedia*.

48. *BELENOIS SEVERINA*, Cram.

Tana River, 4th January, 1899.

Rather a curious male, having a wet aspect on the upper surface, but the under surface characteristic of a late intermediate phase, the black veins being feebly indicated.

49. *BELENOIS MESENTINA* Cram.

♀, Nthatha Hill, Kitwi, 4700 feet, 31st December, 1898;
♂, Neugia, 30th January, 1899.

50. *BELENOIS WESTWOODI* Wllgr.

Tana River, 4th January, 1899.

51. *PINACOPTERYX ASTARTE*, sp. n. (Plate LXX. figs. 6, 7.)

The wet phase of this species, of which we received a male from Fwambo in 1897, is not very unlike that sex of *P. falkensteinii*, but has the costal margin of the primaries shorter and the outer border more interrupted, represented only by spots at the extremities of the first and second median branches; the costal margin of the secondaries below is deep orange. Expanse of wings 60 millim.

The intermediate phase, of which (as well as of the male dry phase) we received sexes in 1889 from Tanganyika, differs in the reduction of the width of the outer border in the male and the paler under surface. The female is bright orange, redder at the base; the primaries with a marginal series of rather large greyish spots becoming black externally, and a similar smaller spot on the disk beyond the middle of the second median interspace; the secondaries with smaller black marginal spots at the extremities of the nervures: below, all the spots are small and black, and there are five tiny squamose spots across the disk of secondaries parallel to the outer margin: body blackish above, whitish below. Expanse of wings, ♂ 63 millim., ♀ 61 millim.

The dry phase (now sent by Mr. Crawshaw) is smaller; the marginal blackish border of the male primaries is reduced to more or less connected spots, the number of black spots on the secondaries is reduced, and on the under surface those of the primaries are absent, whilst the orange costal border of the secondaries is

reduced to less than half its width. The female differs in the same way, but retains its vivid orange colouring. Expanse of wings, ♂ 56-60 millim., ♀ 54 millim.

♂ ♂, ♀, Tana River, 3800 feet, 4th & 16th January, 1899.

Of the male Mr. Crawshay writes—"Plentiful on a bush with a red flower, where I could have taken any number in season; but nowhere else have I seen this insect." This is probably the Eastern representative of *P. pigea*. [An allied species of *Pinacopteryx* common in the same country, but hitherto identified with *P. orbona*, is described below¹.]

52. PINACOPTERYX SPILLERI Staud.

♂ ♂, Tana River, 3800 feet, 4th & 16th January, 1899.

In Staudinger's figure the under surface of the secondaries is represented as unspotted; this is the case with the single example taken on the 4th January; all the others have a series of grey spots across the disk; the tint of these wings below varies from sulphur- to butter-yellow.

From Mr. Crawshay's note it appears that this was taken in company with *P. astarte* on the same red-flowered bush on the river's bank whilst he waited for his men to find a crossing to the other side.

53. PINACOPTERYX GERDA Grose-Smith & Kirby.

Intermediate phase.—♀ (shattered), Muthambi River, 4500 feet, Ndya, 10th January; ♂ (perfect), Tana River, 16th January, 1899.

Dry phase.—♂ (shattered), Muthambi River, 6th January; ♀ (perfect), 11th January, 1899.

Of the first female Mr. Crawshay writes—"Oblong ova, of a greenish-yellow colour"; of the second one—"A perfect specimen at last! The first I saw of this species was on the wet mud of the Tana River, where—when waiting for my net—it was devoured by a dragonfly; the second and third which I took are both rags. Greenish-white spike-shaped eggs."

Professor Aurivillius questions the distinctness of this species from *P. simana*, and suggests that the latter may be a seasonal

¹ PINACOPTERYX VIDUA, sp. n. (Plate LXX. figs. 8, 9.)

Allied to *P. orbona*, which it represents in Eastern and Northern Africa: it has well-defined seasonal phases, the wet phase being most like *P. orbona*; the male, however, has a narrower marginal border to the primaries; the apex of these wings and the secondaries creamy on the under surface, instead of white: the female shows far less grey basal shading, not filling the discoidal cell; the discal spots are wanting from the secondaries, and the orange from the under surface of the primaries (which is characteristic of *P. larima*, the female of *P. orbona* in my opinion, not of *Belenois thyia*). Expanse of wings, ♂ 45-46 millim., ♀ 46-50 millim.

♂ ♂, White Nile, Fôda (*Emin*), and British East Africa (*Gregory*); ♀ ♀, Wasin, and British East Africa (*Gregory*).

Formerly I referred this species to *P. ortygna* (cf. *P. Z. S.* 1888, p. 76), see Aurivillius, *Rhop. Æthiop.* p. 411. It is certainly not *P. gerda*, which is more nearly related to *P. simana*.

form of *P. charina*. I am quite sure that all collectors of South-African species will dissent from his last suggestion, because the wet, intermediate, and dry phases of *P. charina* are well known, and (apart from size) can readily be distinguished from *P. simana* by the uniform character of their upper surface at all seasons, and by the absence of the black veins on the upper surface and the black discal spot on the under surface of the male primaries. We do not possess the wet phase of *P. gerda*, in which the outer border attains to a width of 4 millimetres; but even the intermediate phase has a wider border than the wet phase of *P. simana*; and in all the specimens now received the grey at the base of the primaries is rather more diffused and the black veins are obliterated almost to the outer margin, bringing the species nearer to *P. charina* excepting for the black discal spot of the male and the absence of the dense speckling which characterizes the dry phase of the Natal species; the primaries in *P. gerda* are somewhat shorter, and therefore less acutely triangular, than those of *P. simana*.

Before leaving *Pinacopteryx* it is perhaps as well to point out that "*Ixias venatus*" proves to be a female of a species undoubtedly referable to this genus, and apparently most nearly related to *P. liliana*; it certainly has nothing to do with *Belenois*.

54. *HERPÆNIA MELANARGE*, var. *ITERATA* Butl.

♀, Kitwi, 18th January, 1899.

"Dark yellow ova" (*R. C.*).

55. *PAPILIO ANTHEUS*, var. *UTUBA* Hamps.

Tana River, 3800 feet, 3rd January, 1899.

"This, a poor specimen rather, is one of only two seen here in as many days" (*R. C.*).

56. *PAPILIO NIREUS* Linn.

♀, Undulating plains N. of Tana River, 4200 feet, 14th January, 1899.

"By no means plentiful; the second or third only I have seen" (*R. C.*).

HESPERIIDÆ.

57. *SARANGESA ELIMINATA* Holl.

E. of Athi R., some 4300 feet alt., Kitwi, 18th December, 1898.

"Undulating uplands, timbered with thorny scrub and very dry. This is, I think, an insect very familiar to me in British Central Africa."

Mr. Crawshaw obtained examples of this species at Machako's, in British East Africa, but not in Nyasa-land; he is probably thinking of *S. synestalmenus*.

58. *ERETIS DJÆLÆLÆ* Wlgr.

Clue to exact locality and date lost.

59. *ABANTIS PARADISEA* Butl.

Tana River, 3800 feet, 16th January, 1899.

"Taken on a flowering shrub in one of the hottest places and under one of the hottest suns I have ever experienced" (*R. C.*).

60. *PYRGUS COLOTES* Trim., var.

♂ ♂, Kangonde, 4500 feet, Kitwi, 31st December, 1898.

"Taken when playing together, with a single stroke of the net" (*R. C.*).

These examples are blacker above and browner below than Angolan specimens, but they are spot for spot alike in other respects.

61. *GOMALIA ELMA* Trim.

Muthambi River, 4500 feet, Ndya, 12th January, 1899.

"An insect new to me; taken in the early morning on the flags growing on the river's bank" (*R. C.*).

Mr. Crawshay obtained this insect previously at Kikuyu and Ngongo in July and August, but it is impossible for anybody to remember every obscure little thing that has passed through his hands; it surprises me that Mr. Crawshay remembers so many.

62. *ACLEROS MACKENII* Trim.

♂ ♂, Muthambi River, 10th & 16th January; ♀, 12th January, 1899.

♂. "Quite a new insect to me." ♀? "No doubt the other of the pair which I saw of which I took the one five or six days ago in precisely the same spot—in a cool glade in my camp close to the stream. I have meantime each day visited the same place in the hope of securing the other insect, but without success until now."

♀. "Taken in the early morning on the banks of the stream about twenty yards from the spot where I took the pair" (*R. C.*).

Either the white spots on the primaries are sometimes almost obliterated on the female or the example taken on the 16th January is a male; it, however, has the white border of the secondaries almost entirely restricted to the fringe; the body having been squeezed out I cannot be sure that claspers are present. In my opinion it is a female (as Mr. Crawshay evidently believed), but an aberrant one.

63. *ANDRONYMUS PHILANDER* Hopff.

Muthambi River, 11th January, 1899

64. *PADRAONA ZENO* Trim.

Muthambi River, 3rd January, 1899.

♀. "Containing two rather large yellow spherical ova. The only specimen of this species which I have seen" (*R. C.*).

65. *GEGENES LETTERSTEDTI* Wlgr.

♂, Muthambi River, 7th January, 1899.

66. *BAORIS AURITINCTUS* Butl.

♂, Tana River, 3800 feet, 16th January, 1899.

"Taken in my two hands and then manipulated so as to enable me to transfix it with a thorn, after which I was able to administer the necessary pinch" (*R. C.*).

67. *BAORIS MARANGA*, sp. n. (Plate LXX. fig. 5.)

Not nearly related to anything known to me. Wings deep smoky brown; the male with three unequal streaks of bronze-brown only visible in certain lights, divided by the median vein with its first and second branches; fringe towards anal angle of secondaries and anal tuft paler: wings below slightly more silky than above; primaries with an oblique series of three whitish dots beyond the cell; secondaries with a central patch slightly paler than the ground-colour. Expanse of wings 32 millim.

The female chiefly differs in having five transparent spots in the primaries—three in an oblique series beyond the cell (answering to those on the under surface of the male), and two placed obliquely near the base of the median interspaces; fringe towards anal angle of secondaries and anal tuft whitish. Expanse of wings 34 millim.

Muthambi River, 4500 feet, Ndya, 7th & 10th January, 1899.

Of the male Mr. Crawshay writes, "A new *skipper* to me;" and of the female, "Greenish-white spherical ova."

This species, though unlike any other African species that I have seen, either in collections or plates, and unrecognizable in any published description that I have met with, is perhaps more nearly related to *B. alberti* than to any other known species.

68. *RHOPALOCAMPTA ANCHISES* Gerst.

♀, Bondoni Plains, 5400 feet, 8th December, 1898.

In the present collection Mr. Crawshay unfortunately did not follow his usual plan of putting the exact locality on every envelope, but in some instances referred to a number; thus:—See no. 166, 125, or 146, place and date. Not being at all prepared for this, I took no special note of these collector's numbers until I came to one of the notes, by which time no. 166 had been incorporated and I had lost all clue to it; the others I happily found: this will account for the absence of exact localities and dates for several species, they are all referred to no. 166 (cf. *Acræa cabira*, *Cacyreus lingens*, and *Eretis djelcelæe*).

One other species was obtained by Mr. Crawshay—a possibly new *Ypthima*,—but in such a shattered condition that it is neither fit for description nor for the collection; it is also referred to the unlucky number.

EXPLANATION OF PLATE LXX.

Fig. 1. *Catochrysops peculiaris*, ♀ (intermediate phase), p. 965.2, 3. *Chloroselas azurea*, ♂ ♀, p. 967.4. *Everes kedonga*, ♂, p. 967.5. *Baoris maranga*, ♂, p. 975.6, 7. *Pinacopteryx astarte*, ♂ ♀, p. 971.8, 9. " *vidua*, ♂ ♀, p. 972.

6. On a small Collection of Butterflies from the Nandi District, Uganda Protectorate, Eastern side of Lake Victoria; made by Captain Hobart, of the Grenadier Guards. By ARTHUR G. BUTLER, Ph.D., F.L.S., F.Z.S., &c.

[Received November 1, 1899.]

The collection of which this is an account is small but interesting, the whole of the specimens having been captured by Capt. Hobart on the march. Among them is a *Cymothoe* which appears to be quite new, and of considerable interest from its affinity to Western forms; also *Planema poggei* ♂, and *Acræa leucographa*, an extremely beautiful variation of the Western *Acræa admatha*.

N Y M P H A L I D Æ.

1. AMAURIS ALBIMACULATA Butl.

The white spots on the primaries are reduced in size in all the specimens, but we have similar examples in the Museum collection.

2. CHARAXES CANDIOPE Godt.

One male example.

3. CYMOTHOE HOBARTI, sp. n.

The male above vermilion suffused with carmine, the costal and outer margins narrowly black; primaries with a small apical patch and one subapical spot, sometimes continued indistinctly as a submarginal series; secondaries with well-defined submarginal black spots commencing with a sagittate spot at apex and terminating in an obtusely biangulated linear marking above anal angle; abdominal border pale brown; body bronzy brown, the palpi and under surface of antennal club tawny. The under surface is of a sandy-brownish hue, with fleshy and weak olivaceous changeable tints; the general pattern is that of *C. uselda*, the markings on the basal half being sharply outlined in black; the nearly straight dividing line beyond the middle is dark rich brown in the type, but weakly defined in a second smaller male; the series of A-shaped markings beyond the dividing stripe are alternately pale pearly pink and olivaceous greyish, the outer series of the latter tint uniting into a continuous wavy submarginal line dotted with blackish between the nervures; the femora are whitish. Expanse of wings 42 to 58 millim.

The female nearly resembles that sex of *C. adela*, excepting that the basal area is internally suffused with olivaceous greyish and externally with pale sandy yellowish, the discal series of sagittate markings is weaker in the primaries and very much more so in the secondaries; on the under surface the general appearance is

even closer to that of *C. adela* ♀, but the central stripe is straighter and very dark, with the irregular series of spots which bound it internally white, the outer border (excepting at its extremities) sandy brown, and the discal markings very indistinct. Expanse of wings 64 millim.

Two males and one female were obtained.

4. *PRECIS AURORINA* Butler.

A female (very much shattered) of an example in which the tawny band crossing the wings is nearly half as wide again as usual. If this should be proved to be a constant difference in Uganda specimens, it would be necessary to regard them as representing a distinct local race.

5. *PRECIS CEBRENE* Trimen.

Two males.

6. *PRECIS BOOPIS* Trimen.

One shattered male.

7. *PRECIS GREGORII* Butler.

A pair in poor condition.

8. *CYRESTIS ELEGANS* Bois.

A much shattered specimen.

9. *EURYTELA HYARBA* Fabr.

Two examples.

It is difficult to decide whether these Eastern examples should be assigned to typical *E. hyarba* or var. *angustata*; they seem to vary as regards the width of the white band.

10. *VANESSULA MILCA* Hewits.

An example with an unusually broad tawny band, remarkably resembling the female of *Precis aurorina* in the same collection excepting in size.

11. *PLANEMA LATIFASCIATA* E. M. Sharpe.

Two males.

This is new to us.

12. *ACRÆA ADMATHA*, var. *LEUCOGRAPHIA* Ribbe.

Two males.

This is an extremely pretty local race of the species, which Capt. Hobart assures me was quite common.

PAPILIONIDÆ.

13. *NYCHITONA MEDUSA*, var. *IMMACULATA* Auriv.

One male.

14. *TERIAS SENEGALENSIS*, var. *BISINUATA* Butler.

One female.

15. *BELENOIS MESENTINA* Crainer.

A pair.

16. *GLUTOPHRISSA SABA*, var. *CONTRACTA* Butler.

One male.

17. *LEUCERONIA THALASSINA* Boisd.

One male.

7. Note on the Habit and Mode of Growth of the Corals belonging to the Genus *Pleurocorallium*. By JAMES YATE JOHNSON, C.M.:Z.S.

[Received November 2, 1899.]

In my communication to the Zoological Society on the *Coralliida* of Madeira (P. Z. S. 1899, p. 57) nothing was said as to the cause or meaning of the peculiar habit and mode of growth of the four known species of *Pleurocorallium*. Whilst the species of *Corallium*, such as the red coral of the Mediterranean, branch in all directions and put forth their polype-cells on all sides of the branches, the *Pleurocorallia* ramify more or less in one plane and their polype-cells are confined to one face of the branches.

When Dr. Gray first alluded to the matter (P. Z. S. 1867, p. 125) he said, "I have no doubt that it (the coral under description) grows out horizontally from the rocks, and that they (the polype-cells) arise from the upper surface of the branches." This would appear to have been only a conjecture; but in his Catalogue of Lithophytes or Stony Corals in the British Museum (p. 24) he went further, and described them as "growing horizontally from the sides of rocks," without citing any authority for the statement. The facts about to be mentioned seem to throw doubt on the correctness of Dr. Gray's view, and to suggest another conclusion.

In describing the only known specimen of *Pleurocorallium maderense* (*loc. cit.*), I stated that some zoophytes of rare occurrence at Madeira were growing parasitically upon it. Two of these were branched specimens belonging to the genera *Suberea* and *Stenella*, and they were seated on different parts of their host at a distance from each other. The point to which I wish to draw attention is, that both these Alcyonarians had grown in the plane of the host. If that had extended horizontally, they too had extended horizontally. But can we suppose it possible that they would take in

growing any direction other than a vertical one, whatever might be the position of the coral on which the embryos had settled? May we not hold it as certain that they had grown upright, that is towards the surface of the sea? If so, the *Pleurocorallium* must have taken the same direction.

Another piece of evidence having the same bearing is afforded by some specimens of a simple Madreporarian coral (*Desmophyllum*). All four examples were attached by their bases to the front of the coral, but two of them had twisted themselves round and had pushed their calyces between its branches to the other, that is the posterior side. It is not probable that the *Desmophylla* would have acted so if the supporting coral had possessed a horizontal position, because their calyces would then have been directed downward to the bed of the sea.

Again, it is observable that in a specimen of another species of *Pleurocorallium* in my possession where a small branch by some accident had been broken off the living coral, it had fallen upon the spreading base, to which it had in course of time been made to adhere by an extension of the growing cœnenchyma of the base. It is not easy to understand how the fractured branch could have lodged upon the base in the manner it has done unless the coral had been upright.

In view of these facts, I do not see how we can adopt Dr. Gray's hypothesis, however plausible it may appear at first sight, or come to any other conclusion than that these corals assume in their growth a vertical, not a horizontal position. But if this is so, what is the meaning of the fact that the polype-cells are confined to one face of the branches? If we suppose that the habitat of the corals is in that part of the sea's bed where a constant current is flowing, it is clear that it would be more beneficial to the colonies if all their polypes were turned towards the direction from which their food comes, than if half of the polypes were turned in the opposite direction. A colony will obviously obtain the largest possible supply of nutriment when all its members face the current that carries it, we may say, into their mouths.

To summarize what has been said: in order to account for the fan-like mode of growth of the *Pleurocorallia* and the unilaterality of the polype-cells, Dr. Gray maintained that the corals grow in a horizontal position. As such a position is not easily reconciled with the facts above stated, the suggestion is now put forward that the corals grow upright in the path of a submarine current with all their polypes opposed to the on-coming stream.

8. Further Notes on the Moulting of the King Penguin (*Aptenodytes pennanti*) living in the Society's Gardens. By W. E. DE WINTON, F.Z.S.

[Received November 24, 1899.]

Some observations on the moulting of the King Penguin were offered by me in November 1898 (see P. Z. S. 1898, p. 900). I am now able to supplement those notes by further important particulars, the same bird, still alive and well in the Society's Gardens, having again gone successfully through the moulting.

The dates given for last year were again closely adhered to, and the succession of changes were made pretty much in the same order; but I am able now to explain fully the very strange appearance of the old feathers, which were then likened to withered leaves. The coloured portion of the sheath of the bill was shed as before.

An examination of the freshly moulted feathers showed that the bases were unlike those of the feathers of any other bird so far as I am aware.

Feathers of *Aptenodytes pennanti*.

- A. Part of new feather of flipper, with old feather still attached.
- B. Moulted breast-feather, showing flexible sheath attached to its base.

The quill does not end sharply with a contracted base, but is shed with a flexible sheath attached to it (fig. B). This sheath, which is a continuation of the outer coat of the quill, is, in the body-feathers, as long as the naked part of the shaft. On handling the bird it was found that its enlarged puffed-out appearance at the beginning of the moulting arose from the old plumage being actually raised, and now adhering only to the new feathers that were growing into the

J. Smit del et lith.

Minterr. Dres. imp.

SKIN OF SMITHEMAN'S KOB.
(*Cobus smithemani*)

bases of the old ones, the thin sheaths attached to the bases of the feathers being occupied by the points of the new feathers.

All the feathers so raised had lost most of their original colour, or lost it entirely, the yellow feathers of the neck having bleached white, and the slate-blue feathers of the back and flippers being dull brown or drab except at the extreme tips.

The greater part of the old plumage is removed by the bird's bill as soon as the new feathers upon which it is raised are sufficiently developed to form a covering, but many feathers upon the back and flippers may be left for a longer period (fig. A); these scattered feathers, adhering to the now nearly fully developed new plumage, produce the appearance of small crinkled leaves, which puzzled me so much last year.

The small body-feathers have a large downy aftershaft; the quill-feathers have a naked shaft as long as the plumed portion.

The feathers of this bird are so unlike those of any other bird, in the entire absence of a raised midrib as well as in their umbilical portion, and the nature of the moult seems to open up such interesting questions, that I have placed all the materials possible in the hands of my friend Mr. W. P. Pycraft, who I hope will shortly publish the result of investigations which he is about to make upon these specimens.

9. Description of the Skin of an apparently new Kob Antelope from the Neighbourhood of Lake Mweru, with Note on a Skull and Horns of an Antelope of the same Genus. By R. LYDEKKER.

[Received November 20, 1899.]

(Plate LXXI.)

I am indebted to Mr. Rowland Ward, F.Z.S., for the opportunity of exhibiting to the Society this evening the skin of the very handsome and apparently new species of Kob Antelope forming the subject of the drawing (Plate LXXI.). The specimen, which consists of a flat skin, wanting the head, feet, and the greater portion of the tail, was obtained by Mr. F. Smitheman, F.Z.S., in the neighbourhood of Lake Mweru, situated to the south-west of the lower end of Lake Tanganyika. It arrived in England during last summer, and there were hopes that the head would follow; but if the latter was ever despatched at all, it has evidently miscarried.

The skin has the appearance of belonging to an adult animal, and there is every probability that it pertained to a male. On the under surface there are rudimentary mammae, which indicate that the skin must be that either of a male or of a young female. If it were that of a young female, it would indicate that the adult animal was of very large size; but I think it may be pretty confidently assigned to an adult male.

From the general characters of the pelage and its coloration, and

especially from the long shaggy hair on the nape of the neck and the absence of a mane, the specimen may be confidently assigned to the genus *Cobus*. Additional evidence in favour of this reference is afforded by the circumstance that the long hair on the middle line of the back is reversed from a point some distance in advance of the loins to the withers, exactly as in the Puku (*C. vardonii*).

In size the animal to which the skin pertained must evidently have been considerably larger than the species last mentioned, and may have been more nearly comparable in this respect with *C. maria* of the Bahr-el-Ghazal. In colour, the nape, the sides of the buttocks, and thighs are bright chestnut-tawny; the middle line of the back, the hinder portion of the shoulders, and the hind-quarters are the same chestnut-tawny mingled with blackish brown; the fore part of the shoulder, a line on the under surface of the neck, the flanks, and the front surface of the fore-limbs and of the lower part of the hind limbs are of a deep glossy blackish brown, the under-parts being dirty white.

The portion of the skin of the neck remaining, which seems to have been cut off a considerable distance below the head, is suggestive of a comparatively long-necked animal. And if this be a correct inference, it would be natural to expect that the horns were of a comparatively long and slender type. Now, in its dark colour the skin is more like that of *Cobus maria*, of the swamps of the White Nile, which is a species with comparatively long, slender, and doubly curved horns; and it is to that animal, rather than to any other member of the genus *Cobus*, so far as the materials permit of forming an opinion, that I am inclined to consider the form represented by the skin before us most nearly related. Altogether apart from the distance between the White Nile and Lake Mweru, the skin under consideration is broadly distinguished from the male of *Cobus maria* by the absence of the white patch on the withers and the white line down the back of the neck. As regards the female of the latter species, there is a definite statement in the 'Book of Antelopes,' vol. ii. p. 122, that it is similar in all respects to the buck, except for the lack of horns; but in the plate a female is figured without the white saddle and neck-line¹.

As there is no other Antelope of which the skin is known that presents any close resemblance to the specimen under consideration, and since it is certainly distinct from the male of *C. maria* (being itself probably a male specimen), I take leave to regard it as representing a new species, for which I propose the name of *Cobus smithemani*; the skin represented in the figure (Plate LXXI.) being of course the type. The species may be provisionally defined as a large-sized Kob, differing from every other species of the genus except *C. maria*, and distinguished from the male of the latter by the absence of the white line down the back of the neck and the patch of the same colour on the withers, in which region the present species is chestnut.

¹ From a specimen that has recently come under my notice, the white neck-line does not seem to be constant even in the bucks of *C. maria*.

I may now mention another piece of evidence in favour of the relationship of this *Cobus smithemani* to *C. maria*. A correspondent of Mr. Rowland Ward writes that in the neighbourhood of Lake Mweru he has seen an Antelope very like a Situtunga, with similarly elongated hoofs, but with horns more like those of a Lechwi, although longer and slighter. Such a description would admirably fit the present species, if, as I suggest, it be more nearly allied to *C. maria* than to the Puku. Supposing it to have elongated hoofs, I should not regard such difference as, at the most, of more than subgeneric importance.

Skull and horns of Loder's Puku (*Cobus vardoni loderi*).

Before concluding this communication, I may call attention to the skull and horns of a Puku-like Antelope in the collection of Sir E. G. Loder, which I first thought might belong to the same species as the above skin. This skull and horns, of which the locality is unknown, are shown in the accompanying drawing. The

specimen is essentially of the Puku type, but broadly distinguished by the circumstance that while the skull itself is slightly shorter than that of an average-sized Puku skull in the British Museum, the horns are very much longer and stouter. In the Puku skull the length from the fronto-parietal suture to the tip of the nasals, measured in a straight line, is 8.5 inches, while in Sir E. Loder's specimen the corresponding dimension is but 8 inches. In the present specimen the horns have more ridges (17) and relatively shorter tips than any Puku horns I have seen; they measure 20.3 inches along the front curve, 8.0 inches in basal circumference, and 8.1 inches between the tips. Now the only horns assigned to the Puku with which I am acquainted that have anything like these dimensions are a pair obtained by Mr. Smitheman from the Luswesi Valley, in the neighbourhood of Lake Bangweolo, which lies S.S.E. of Lake Mweru; these horns measuring $20\frac{1}{4}$ inches along the curve, $8\frac{1}{2}$ in basal girth, and $12\frac{1}{4}$ from tip to tip¹.

The wide interval between the tips I consider of no importance, but in other respects these horns agree very closely as regards measurements with Sir E. Loder's specimen. And they differ from the next specimen of Puku horns ($19\frac{1}{8}$ in.) in Mr. Rowland Ward's list² by the much greater girth, the basal circumference of the latter being $6\frac{3}{4}$ inches.

Accordingly, so far as horn-measurements alone are concerned, there would seem a probability that Mr. Smitheman's Lake Bangweolo skull may be specifically identical with Sir E. Loder's specimen. And if this be so, there arises the question whether both are not referable to *C. smithemani*. But if the evidence of the correspondent quoted above as to the Lechwi-like character of the horns of the Antelope presumed to be identical with the latter be reliable, this can hardly be the case. It must also be remembered that Lake Bangweolo is a considerable distance from Lake Mweru, so that each district (in spite of the fact that the true Puku and Lechwi extend from the Chobi-Zambesi Valley to Lake Mweru) may have its own particular species or race of Kob.

The matter is one of great difficulty, and I may be accused of rashness in what I propose to do, which is to consider, for the present, Sir E. Loder's specimen as typifying a large-horned race of Puku to be known as *Cobus vardoni loderi*, until it can be either proved to be the same as *C. smithemani* or entitled to rank as a species by itself. Whether Mr. Smitheman's large Puku head from Lake Bangweolo belongs to the same form may be left an open question.

¹ See Rowland Ward, 'Records of Big Game,' p. 189 (1899).

² *Loc. cit.*