in one case the patient became so blind that vision was reduced to perception of shadows. In neither of these cases did any therapeutic measure do more than relieve some of the symptoms.

In a more recent instance the disease was confined to lower culdesacs, but involved both the ocular and palpebral conjunctivae, resembling Koerber's case briefly outlined by Dr. Weidler.

Looking back on my slight experience of ocular pemphigus, I have noticed that the disease is frequently mistaken for and treated as a manifestation of trachoma and this is an error that in some instances may easily be made.

Dr. H. W. WOODRUFF, Joliet, Ill.: I operated for complete atrophy of the conjunctiva in a case which I attributed to pemphigus. There might be some doubt as to the diagnosis, but it had reached the stage of complete atrophy so that there was no longer any culdesac. It may be that trachoma may reach that stage after a long time, but I had never seen such a case and attributed it to pemphigus. I entirely relieved the contraction by skin-grafting after the method of using a plate of block tin with the skin graft placed over the tin, doing it in two operations; that is, restoring the lower culdesac first and afterward the upper one. This gave the patient some mobility of the eye, but the clearness of the cornea was not restored.

Dr. E. L. JONES, Cumberland, Md.: I saw a very severe and typical case four or five years ago in which Dr. Price, of Frostburg, had made the diagnosis. It was in an elderly woman already affected for several years with pemphigus in the vagina. It had now affected the eye which looked as though it had been scalded, with white blisters. It gradually extended until the entire conjunctiva was destroyed, also the cornea, and everything was matted and massed together. In the course of time the other eye was affected similarly. The disease lasted ten years or more. The mouth was invaded by white blebs of gelatinous consistency, followed by atrophy and stricture of the esophagus with intense suffering, and for years the patient could only swallow milk in small sips. All sorts of treatment were tried but nothing availed. The woman has recently died.

Dr. JOSEPH S. LICHTENBERG, Kansas City, Mo.: I had a case exactly parallel with Dr. Weidler's. The history was indefinite; it could not be determined whether it was trachoma or pemphigus but from the age of the patient and the history it was concluded that it was pemphigus. In this case I restored all four culdesacs with mucous membrane grafts taken from the mucous and submucous tissue of the lips on the Wolff principle and not on the Thiersch principle. This case is now of four or five years' duration and still holds good.

Dr. WALTER E. LAMBERT, New York: I was called by a general practitioner to see a patient on account of the very serious involvement of both eyes in a case diagnosed as acute septic pemphigus, resulting from a streptococcus infection after a submucous operation in the nose. The patient had a most extensive eruption, vesicular and pustular, involving the mucous membrane of the mouth, throat and eyes, and the skin over the entire body. There was quite a high temperature and the suffering was intense. The attending physician was greatly perplexed, and at first, thinking it was a case of small-pox, had called in a health officer. Dr. George T. Elliot saw the case in consultation, and made a diagnosis of acute septic pemphigus. The patient died from general septic infection three or four days later.

Dr. WALTER BAER WEIDLER, New York: When this patient was received in the clinic the diagnosis made was trachoma, but Dr. Thomson seeing the patient immediately suggested the diagnosis of pemphigus, and thorough study of the case proved the correctness of that diagnosis. The patient was presented at the Academy of Medicine and at the ophthalmologic and the dermatologic sections and there were some who doubted the correctness of the diagnosis, because there had been no general eruption whatsoever, but there were lesions which we were able to demonstrate in the nose. Both eyes were completely sealed together at the palpebral margins, and on the patient's last visit to the clinic there was a general eruption of pemphigus seen on the abdominal walls.

ORBITAL CELLULITIS FROM DISEASE OF THE SUPERIOR MAXILLA IN CHILDREN

WILLIAM CAMPBELL POSEY, M.D.
PHILADELPHIA

Orbital cellulitis in children is not very uncommon, and I should hesitate to ask the Section to devote any time to its consideration were it not for the fact that I believe the sublying cause in many of the cases which come under our observation is permitted to remain undetermined, and that we proceed with surgical and therapeutic measures without trying to analyze thoroughly the agency in each particular case which may have given rise to the cellulitis. It shall be my endeavor, therefore, to elucidate, as far as I am able, the various conditions which may operate as etiologic factors in this affection, and to illustrate one phase of the subject I shall, without further preamble, recite the histories of two cases which I have had under my care at the Children's Hospital during the past few years.

CASE 1.—*History*.—F. C., female, aged 1 year, came to the hospital Dec. 15, 1906, on account of a deformity of the left lower eyelid and a discharging fistula in the same location. The history as recorded by the resident physician, Dr. Vail, is as follows: The patient was the first child; no difficulty at birth; breast-fed; vaccinated when 3 months old by the family physician; parents living and well. When the child was 8 months old the left axilla swelled and was opened, pus escaping. Pertussis occurred one month later, followed by pneumonia; with convalescence after two months. A month later the left eye swelled and the lids shut tight, much pus running from the eye. About the same time an abscess in the cheek opened spontaneously by way of the mouth.

Examination.—At admission, the head was found well formed, the left lower lid ectropionized, with a discharging sinus at the lower outer angle of the orbit. Examination of the nose and throat was negative; vaginal smear negative.

Treatment and Course of Disease.—Under ether I passed a probe into the orbital sinus and dead bone was encountered on the floor of the orbit. This was removed by curet and dissection, an incision having first been made along the lower rim of the orbit. Part of a newly-formed but decayed tooth was removed from the upper left alveolar process. Although the left cheek was considerably swollen, no sinus could be detected in the mouth. The orbital wound was drained by a small tube. The wound was dressed daily, copious quantities of pus escaping through the drainage tube at each dressing. At the expiration of six days after the operation the wound failed to drain and orbital cellulitis supervened, the eye being proptosed and the bulbar conjunctiva ecchymotic. Dr. Packard saw case in consultation and concurred in the diagnosis of necrosis of the left superior maxillary bone. The patient was etherized a second time, the orbital wound thoroughly drained and a sinus discovered leading from the dead bone in the floor of the orbit into the cavity of the bone. The bone was incised through the cheek just above the alveolar process, and a drainage-tube passed from the orbit through the bone into the mouth, thereby establishing free drainage. Subsequent recovery was uneventful, the swelling in the orbit and over the superior maxillary gradually subsiding, and after some weeks the sinuses drying up. Tonics and mercurial inunctions were administered throughout the entire course of the disease.

CASE 2.—*History*.—M. Z., aged 2 years, was admitted to the wards Nov. 2, 1911, on account of a marked prominence of the left eye, with swelling of the lids and adjacent cheek, which had been present for some days before admission. The mother stated that the child had been in good health, following an operation for abscess on the chest (empyema) some four months previously, and denied the possibility of the condition being due to a traumatism.

Examination.—Marked proptosis of the left eye was evident with all the attendant signs of orbital cellulitis. The left cheek was also puffed and the palate and alveolar border were found to be indurated and swollen. The left pupil was slightly dilated and an ophthalmoscopic examination revealed some pallor of the optic nerve, with distinct engorgement of the retinal veins. The right eye and side of the face were unaffected. Dr. Packard examined the nose and throat and pronounced them to be normal.

Treatment and Course of Disease.—The case being regarded as one of osteomyelitis of the superior maxillary bone, the patient was etherized and the orbital tissues and the alveolar border were freely incised, giving exit to a large quantity of thick pus. A number of loose teeth were also removed from the affected alveolus. No dead bone was encountered. The orbital wound was drained from its most dependent portion and a dressing applied. At the end of three weeks all the sinuses had closed and the child was discharged from the house. Two weeks later, however, she was readmitted, on account of a painful swelling of the left leg. This was incised and found to be dependent on disease of the tibia. At the end of another week the swelling reappeared in the left upper jaw and there was some proptosis. The patient was again etherized and with the aid and counsel of Dr. Packard the orbital and alveolar tissues were again incised. Several more dead teeth with sequestra were removed from the upper jaw and free drainage established from the orbital and alveolar structures. The floor of the orbit seemed smooth anteriorly, but rough posteriorly, though it was impossible to assert positively that the bone was actually necrosed. Healing slowly followed, the sinuses gradually closing, and the child was discharged at the end of another month as cured.

In neither case was there a history of traumatism, and in neither had the attention of the mother been called to any nasal condition, and a careful examination of the nose by Dr. F. R. Packard, the rhinologist to the hospital, which was made while the inflammation was at its highest, failed to reveal any evidence of nasal or sinus disease. A history of inherited syphilis could not be obtained in either of the two cases and signs of tuberculosis were absent. In each of the cases, however, there was evidence of previous infection of the general system, i. e., axillary abscess and pertussis, followed by pneumonia, in the first case, and empyema in the second, while in the second, staphylococci were found in the discharges. Unfortunately, no culture was made in the first case.

I am led, therefore, to the supposition that the orbital cellulitis in these cases was due to disease of the superior maxillary bone, to an osteomyelitis, doubtless secondary to a general blood infection.

ANATOMIC CONSIDERATIONS

When one considers the structure of the superior maxillary in infancy and early childhood, the reason for its not infrequent involvement in infectious processes is not difficult to understand, for at that age the bone is quite a different structure from what it gradually evolves into later. For example, to quote Logan Turner:¹

"At birth the body of the superior maxillary is made up almost entirely of the alveolar process of the jaw, the sockets of the teeth being almost in contact with the orbital plate of the maxilla forming the floor of the orbital cavity. The antrum exists merely as a slit-like indentation on the outer wall of the nasal chamber, and there is, therefore, at birth no maxillary sinus beneath the orbit." According to Turner, "the growth of the body of the bone, and with it an increase in the size of the face, takes place by the formation of a mass of cancellous or spongy bone between the alveolar process

and the orbital plate of the maxilla. In this way the alveolar process, along with the teeth, becomes separated from the orbital plate. A little later, coincidentally with the increase in the cancellous tissue on the facial and dentary aspects of the bone, a process of absorption apparently takes place on its nasal and orbital surfaces, thus causing an increase in the size of the antrum. This simultaneous process of growth and absorption continues until the eruption of the

Fig. 1.—The left superior maxilla of a new-born child, viewed on its nasal aspect. This and the three following figures show the development of the superior maxillary bone from "The Accessory Sinuses of the Nose," by A. Logan Turner.

Fig. 2.—Vertical transverse section through the right superior maxilla of a new-born child, viewed from behind. The antrum is a mere recess on the nasal aspect of the bone, and the teeth sockets lie almost in contact with the floor of the orbit.

Fig. 3.—Vertical transverse section through the right superior maxilla during the first dentition, on the plane of the second molar tooth. The antrum, viewed from behind, is small, and the body of the maxilla consists mainly of cancellous bone.

Fig. 4.—Vertical transverse section through the right superior maxilla of an adult, viewed from behind, on the plane of the first molar tooth. The section shows that the cancellous bone is almost entirely absorbed and that the maxillary sinus has reached its full dimensions. The fang of the tooth projects into the floor of the sinus.

third permanent molar or wisdom tooth at or about the twenty-fifth year of life, at which time the antrum reaches its complete or adult form."

In infancy, therefore, the cancellous structure of the bone largely predominates, rendering it especially liable to inflammatory processes.

1. Turner, Logan: Accessory Sinuses of the Nose, Edinburgh, 1901, p. 5.

Four illustrations (Figs. 1, 2, 3 and 4) have been reproduced from Turner's admirable book, which will illustrate the changes which have just been described, and demonstrate the manner in which the bone evolves from a mass of cancellous bone to a thin framework, offering attachment to the upper teeth, forming walls for the important spaces which surround it, as well as for the huge antral cavity within its structure.

When the complexity of the mechanism of dentition is considered, it becomes apparent, I think, how easily infection of the alveolar border may be followed by a general inflammatory process of the entire bone. A study of the literature has revealed a number of cases in which such was the case, the inflammation being apparently conveyed to the alveoli by an infected nipple during the act of suckling. The alveolar border once involved, suppurative periostitis soon extends rapidly under the periosteum along the anterior surface of the upper jaw or through the maxillary antrum to the orbit. As a rule, an acute osteomyelitis of the entire superior maxillary follows with all the signs of an active inflammatory process, such as is observed from whatsoever

and if the inflammation be continued, of atrophy. After a variable period of time, but generally rather rapidly, cure may follow by absorption and the affection may disappear without leaving a trace; more usually, however, spontaneous evacuation follows the suppurative process and fistulas persist in the orbit and in the mouth. In consequence of the former, cicatrices affect the lid and ectropion results. At times the entire lower rim of the orbit may slough away, leaving a most disfiguring, indrawn scar (Figs. 6 and 7). As is usually the rule in osteomyelitis, the sequestrum shows but little tendency to separate itself from the living bone for a long time, so that healing is frequently very protracted.

Orbital cellulitis may arise from disease of the superior maxillary in consequence of traumatism; indeed it not infrequently happens that a very slight blow will excite most active symptoms in children in whom the vitality has been lowered by some form of exhausting disease.

Morax⁵ has recently drawn attention to this class of cases and describes how staphylococci circulating in the blood may infect the site of a contusion endog-

Fig. 6.—Case 2, showing sequel of orbital cellulitis; fistula at lower outer rim of orbit, with swelling of the cheek.

Fig. 7.—Tuberculosis of the superior maxillary bone with abscess of the orbit, showing ectropion of upper lid due to cicatricial connections with upper rim of orbit.

cause. The cheek and lids become swollen, the globe is proptosed and its movements restricted. The hard palate is swollen and edematous and the alveolar process tumefied. There is a marked tenderness over the superior maxillary and pain may be complained of. This may be localized or may radiate into the temple, forehead, into the nose and teeth, or may be referred to the entire half of the corresponding side of the face. The pain may be most intense, especially at night, and may be accompanied by vomiting. The inflammation in the orbit may extend along the periosteum posteriorly, and, invading the sphenoidal fissure, may implicate all the nerves of the orbit, both motor and sensory. The Gasserian ganglion may become affected and keratitis neuroparalytica may supervene with or without concomitant symptoms. Phlegmon of the orbit may arise from inflammation of the fatty cellular tissue of the orbit and may be followed by thrombosis of the orbital veins or even of the cavernous sinus.

Not infrequently the optic nerve becomes implicated and the ophthalmoscope may reveal the signs of neuritis,

enously and may give rise to osteomyelitis in cases in which there had been no wound in the integument. Willard⁶ asserts that while the staphylococcus is the most common bacterium present, the pneumococcus and streptococcus, or other organisms are occasionally encountered.

While no history of trauma was obtainable in either of my cases, it is not unlikely that some slight injury may have occasioned and excited the inflammation in the superior maxillary, the bacteria in the blood acting endogenously in giving rise to the general osteomyelitis.

Contusion of the orbit is especially liable to be followed by caries of its walls when the subjects are tuberculous, such as is observed when other bones are injured in this class of cases. Tuberculosis may affect the orbit, however, without the necessity of an injury to excite it, the superior maxillary and the malar being in common with the inferior maxillary the most common seat of tuberculosis of the face.

5. Morax: *Ann. d'oculist.*, March, 1905, p. 208.

6. Willard: *Surgery of Childhood*, 1910.

We are now brought to a consideration of the possibility of orbital cellulitis in the class of cases under discussion being occasioned primarily by an inflammation of the maxillary sinus, and while there are those⁸ who assert that the antrum is too small at such an age to be the seat of an inflammatory process sufficient to excite an osteomyelitis in the body of the bone, it seems to me that the recent studies of Onodi and the large number of clinical observations which have been recorded by trustworthy observers attest that such may be the case, and that we must acknowledge that such an inflammation may not be an infrequent cause of disease of the orbit in children.

All of us will doubtless bear testimony to the debt ophthalmology owes to Onodi in his elucidation of the manner in which the optic nerve may be involved from sphenoiditis and ethmoiditis in adults, and it is my opinion that the observations of this distinguished Hungarian rhinologist in his most recent volume⁹ will prove of equal value to us. I have ventured to reproduce a number of the plates from this book, which show the development of the antrum and trace the growth of the cavity through its entire development.

The reproductions have been reduced in size, but show in coronal, horizontal and longitudinal sections the topographic relations and form of the maxillary antrum from birth to the eighth year.

These investigations prove apparently that the antrum is quite large enough to become the seat of inflammation even in the earliest months of life. To these favoring anatomic conditions the weight of actual clinical experience may now be added.

In an admirable dissertation, Guisez¹⁰ has given the clinical picture of maxillary sinusitis in the infant and attributes the common or acute form to grippal coryza, to involvement of the mucous membrane of the sinus during convalescence from infectious diseases, or, in fine, to any form of nasal infection. On account of the thinness of the anterior wall of the sinus at that age, he asserts that the sinusitis manifests itself principally by the external symptoms, which are, in consequence, much more accentuated in infants than in adults. The cheek swells and is edematous, the lids become involved and the tissues of the orbit are affected. Discharge from the nose is at all times incessant and abundant. Osteomyelitis of the body of the bone follows and caries sets in. In one case of an exceptionally severe nature, which nearly terminated fatally, Guisez was compelled to operate to relieve the symptoms occasioned by thrombosis of the ophthalmic vein.

Comparatively recently Beauvois¹¹ sums up the entire subject in a masterful manner and in addition to other cases cites that of an infant 15 days old, in which orbital inflammation was attended by a copious discharge of pus from the nose, the exophthalmos gradually disappearing as the nasal discharge subsided. Beauvois suggests that in very young infants the leukorrhoeal discharge of the mother may readily infect the nose of the infant, or the contagion may be conveyed through the medium of the bath or even of the air.

Fano,¹² Weissman and Fiocre¹³ and Dujardin¹⁴ have all reported cases of sinusitis in children, the latter in an infant 19 days old.

It has long been a matter of clinical experience that orbital cellulitis may follow scarlet fever, measles, typhoid fever, tonsillitis, erysipelas, small-pox and other infectious diseases, and there is strong evidence that in many of these cases, at least, the inflammation in the superior maxilla follows a primary involvement of the antrum.

Lange,¹⁵ for example, in a paper containing most of the literature on the subject, has reported six cases following scarlet fever, in which an acute empyema of the sinus gave rise to osteomyelitis of the superior maxilla and terminated in caries of the bone with the formation of sequestra, precisely as is so frequently observed in middle-age disease after this form of fever.

Estor¹⁶ is authority for the statement that maxillary sinusitis may also originate in children from a carious tooth, just as it does in adults, the alveolar process becoming infected and the inflammation spreading by continuity of tissue under the periosteum to the antral cavity.

While American literature, which is so full of original treatises on all forms of sinus disease and its treatment, contains but scant reference to the subject, F. Krauss has recently reported a case of orbital involvement from disease of the superior maxilla in a 4-months-old child, in which the inflammation was apparently dependent primarily on antral disease.

As in the diagnosis of disease of the sinuses in adults, much information may be derived from an x-ray study of their condition in children, and those interested will do well to consult the splendid series of radiograms of infants' sinuses recently published by Haike,¹⁷ as well as those by Onodi. In all suspected cases of primary sinusitis, care should be exercised to exclude the presence of a foreign body in the nose and intranasal syphilis.

Syphilis, however, of the superior maxilla is rare in children. Neither Hutchinson nor Keyes refer in either of their text-books to the possibility of this bone being the seat of syphilis in young subjects, though both describe at some length the infiltration and softening between the tables of the skull and in the sutures, and the diffuse periostitis of the frontal and parietal bones, which occur at this age. Indeed, Hutchinson¹⁸ asserts that while children of the age of 6 to 10 to 14 years appear to be prone to suffer from bone disease of a syphilitic nature, and at this period of life multiple periostitis of a very extensive kind is not uncommon, the periostitis of inherited syphilis which leads to supuration is but seldom met with in infancy. When syphilitic periostitis does affect the orbit, both cavities are usually involved and the diagnosis is rendered easy by the evidence of periostitis elsewhere in the body.

Treatment consists in early evacuation of the pus from the orbit and of establishing proper drainage from that cavity and through the body of the superior maxilla if it, too, be the seat of disease.

Prompt healing may follow, and even when necrosis of the bone occurs, almost complete regeneration of the bone will take place provided the periosteum be saved.

8. Axenfeld (*Bacteriology of the Eye*, p. 387), with his vast experience, says that these rare cases of orbital cellulitis in infants cannot be associated with affections of the sinuses, as these latter are either absent or underdeveloped up to the eighth year, and asserts that it is only after the sixth year that a nasal etiology becomes important. The youngest subject he has seen with orbital abscess which certainly originated from a sinus (sphenoidal containing streptococci) was 9 years old.

9. Onodi: *The Accessory Sinuses of the Nose in Children*.

10. Guisez: *Le Pratique des maladies des enfants*, p. 346.

11. Beauvois: *Rec. d'ophth.*, 1908, p. 54.

12. Fano: *Jour. d'ocul. et chir.*, Paris, 1873-1874, I.

13. Weissman and Fiocre: *Ann. d. mal. de l'oreille et du larynx*, 1906.

14. Dujardin: *Jour. d. Sc. méd. de Lille*, Feb. 24, 1888.

15. Lange: *Med. klin.*, Aug. 19, 1906.

16. Estor: *Guide pratique de chirurgie infantile*.

17. Haike: *Arch. f. Laryngol. u. Rhinol.*, 1910, xxiii, 217.

18. Hutchinson: *Text-Book on Syphilis*, p. 389.

Should deformities of the lids occur in consequence of contractions from fistulæ, they should be corrected by plastic operations some months after all acute symptoms have subsided.

2049 Chestnut Street.

ABSTRACT OF DISCUSSION

DR. F. R. PACKARD, Philadelphia: Dr. Posey calls his case a case of orbital cellulitis due to disease of the superior maxillary bone—not due to disease of the maxillary sinus. I remember in the 1910 meeting of the American Otological Society, Dr. William C. Braislin, of Brooklyn, read a paper on the accessory sinuses of the nose in a 5-year-old child, with notes on cases of suppuration of the antrum of Highmore at and before that age. He described one or more cases practically identical with the two reported by Dr. Posey. In the discussion that followed, a great majority took the view that there was no such thing as suppuration of the antrum in a child 1 or 2 years of age; in other words, that the antrum was not sufficiently developed to permit of suppurative disease, in the true sense of that term. As Dr. Posey said, Onodi has pointed out recently, in agreement with other men who work on the accessory sinuses, that this sinus is appreciably well developed in the first year. Onodi has done more than anyone else to show that this sinus is the first one to be developed. It is present at birth and within a month is a fair-sized cavity. At that time also its opening is large and its floor is lower than the ostium. In other words, at a very early age the maxillary antrum possesses the same disagreeable characteristics as in adult life, namely, a large portal of entry for organisms and poor facilities for drainage. The cases, however, which are reported ordinarily as maxillary suppuration in young children have all of them, as far as I can learn, presented characteristic disease of the maxillary bone, and not the ordinary suppurative sinusitis.

I believe that the explanation of these cases is to be found, not only in the entrance of infection between the teeth, as Dr. Posey mentioned, but also in the fact that the child's nose is just as susceptible to inflammation or infection as the adult's and that the large size of the antral opening, which is very much larger in early childhood than in the adult, accounts for some of these infections, and therefore if micro-organisms once get in either through the tooth-channel or the ostium, the large mass of cancellous bone tissue which undergoes subsequent absorption to form the antral cavity is the kind of bone which is susceptible to osteomyelitis and other inflammatory bone troubles. In other words, I think Dr. Posey chose wisely when he termed his cases diseases of the maxillary bone and not of the sinus, although the infection of the bone occurred through the small chamber corresponding to the large antrum which we find in the adult skull.

MEASUREMENT OF FATIGUE OF THE OCULAR MUSCLES

LUCIEN HOWE, M.D.

BUFFALO, N. Y.

I. INTRODUCTION

1. *Object.*—The object of this study was to obtain a record written by the ocular muscles themselves of the fatigue which they develop under various circumstances. This was with the expectation—which has apparently been realized—that the data thus furnished would enable us to separate more clearly the factors which, when grouped together, have been called "eye-strain."

2. *Method.*—The autographic records here referred to have been made by a modification of the ergograph. This is a piece of apparatus well known to physiologists, and described now in all of the text-books on that subject. But as it is important for our purpose to under-

stand the principle on which it depends, its construction and action, it is desirable first to consider briefly the measurement of fatigue in other muscles, in animals and in man.

II. FATIGUE OF MUSCLES OF THE HAND OR LEG AS MEASURED BY THE ERGOGRAPH

More than twenty years ago an Italian, Mosso, measured the fatigue produced when a given group of muscles contracts repeatedly. To the last phalanx of one of the fingers he attached a sort of glove, from which a string extended horizontally over a pulley, and to the other end of which a weight was attached (Fig. 1). Each contraction of the finger required a given amount

Fig. 1.—General view of the ergograph of Angelo Mosso (Arch. f. Physiol., 1890).

Fig. 2.—An ergographic record of fatigue of the muscles of a finger.

of work to lift the weight. The extent and character of these contractions were registered on a receiving drum.

Such tracings made by the leg muscles of the frog, or of some other cold-blooded animal, usually give the most typical pictures (Fig. 2).

Although these tracings are familiar to every student, there is one point about them, not always mentioned in the text-books, that is of clinical importance. It is known among physiologists as the "Treppe," or "staircase."

As long ago as 1865 Ranke said: "Everyone knows that the first twitch of a muscle is not its greatest. With stimuli uniform in strength, the later contractions are stronger than the earlier ones—a phenomenon the reason for which has heretofore been wholly obscure."