cataract was present on each side; that on the left side might have been thought traumatic in origin had not that of the right lens been so obviously otherwise. A slight bilateral divergent strabismus was evidently the result of defective vision.

His articulation of both English and French was slurred and difficult to follow, though no structural defect was found which could account for this symptom.

The sternomastoid muscles were completely atrophic and there was some wasting in the extensor muscles of the forearms. The position of rest of both hands was one of slight flexion, the thumbs being slightly adducted. All movements were present, but it was found that after making a strong, grasping movement with either hand he was unable quickly to extend the fingers again, the muscles involved being apparently in a state of cramp. If the grasping movements were frequently and quickly repeated this cramp became less and less noticeable until, after perhaps half a dozen efforts it would completely disappear. This phenomenon was not present

both sides. In the muscles of the forearms and in the intrinsic hand muscles reactions were strong but enormously prolonged.

There were no pathologic changes in either the blood or the urine. The Wassermann test of the blood-serum and the cerebrospinal fluid was negative.

CONCLUSION

The combination of premature bilateral cataract with atrophy of the temporal, orbicular, masseter, sternomastoid, vasti and anterior tibial muscles, together with a sharply contrasted myotonus in the hands, occurs too frequently to be ignored and most probably points to a deficient hereditary endowment as the approximate cause of the disease.

20 West Fiftieth Street.

POLYPOSIS GASTRICA (POLYADENOMA)*

JESSE S. MYER, M.D.

Associate in Medicine, Washington University, Medical School
ST. LOUIS

Although many instances of the presence of polyps in the gastro-intestinal tract are reported in the literature, they are not altogether common in the stomach. Reports of the existence in the stomach of a sufficient number to justify the term "gastric polyposis" are meager, indeed. In a search of the literature it has been rather surprising to see how lightly the subject of gastric polyps has been touched on in text-books on diseases of the stomach, as well as those on its pathologic anatomy. Indeed, the only text-books on diseases of the stomach which do more than mention the subject are those of Bouvert, Hemmeter and Deaver and Ashhurst.

Chosrojeff finds a dearth of information on benign tumors of the stomach in general, and attributes it to the fact that they present the picture of malignant disease and are thus overlooked. As a result of his investigation, Alfred Tilger in 3,500 necropsies reports but fourteen benign tumors of the stomach. In the Obruchow-Krankenhaus, in 7,500 necropsies, four cases of polyps of the stomach were found, or 0.053 per cent. In Russian hospitals he found the percentage to vary from 0.007 to 0.04 per cent. Versé collected reports of fifty-five cases of polyps of the digestive tract, in four of which the polyps were in the stomach. Ebstein, in 1864, found in 600 necropsies, fourteen cases of stomach polyps, and was able to collect reports of eight others from the earlier literature. Of these twenty-two cases, twelve were of solitary growths, and the others comprised one large polyp with several small ones. In only three cases did Ebstein feel justified in speaking of polyposis; these presented as many as 200 polyps which varied from the size of a pea to that of a bean.

Much of the literature on gastric polyposis has come to us from the French. As early as 1888, Menetrier wrote a very complete study of the subject, and his subdivision of *polyadenomes polypeux* and *polyadenomes en nappe* is still accepted. The earliest reports were made by Morgagni, Andral, Cruveilhier, Vulpian, Leudet, Cornil and Richard. Single polyps of the stomach have been reported by Lyman, Lange, Bennett, Skifossowsky, Hind, Chaput and others.

Wegele, in 1908, reported one case of polyadenoma, and in a search of the recent German and English literature was able to find the treatment of this subject only in the works of Collier, Hanser, Post and Galland.

* Read before the Sixteenth Annual Meeting of the American Gastro-Enterological Association, Washington, D. C., May 6, 1913.

Characteristic facial appearance in myotonia. Incomplete ptosis of both eyelids, marked hollowing of the temporal fossae and flattened masseteric region.

in any other muscle or group of muscles. The abdominal muscles and those of the back and chest were normal. The condition of the spine was normal. The vastus internus and vastus externus of the right thigh were diminished in bulk, though no change was found in the intervening rectus femoris muscle. The knee-jerk on the right side was difficult to obtain. There was a reduction in response directly proportional to the reduction of the muscular mechanism involved. There was almost complete atrophy of the anterior tibial and peroneal muscles with complete bilateral foot-drop as a result. The ankle-jerks and plantar reflexes were absent. The abdominal and arm reflexes were normal.

The gait was of the steppage type peculiar to paralysis of the dorsi-flexors of the feet. There was no sensory change. No reaction to galvanism or faradism was present in the orbicular, temporal, masseter or sternomastoid muscles. There was almost complete loss in the anterior tibial groups of

In French and English literature, Menetrier, Brissard and Norman have each reported one case. Chosrojeff recently reported a case in which he removed six large polyps varying from the size of a pigeon's egg to that of a hen's egg, together with a large number of smaller ones from the size of a pea to that of a hazelnut. Bouvert, in his splendid treatise on the subject, states that he never encountered a case.

The etiology of polyadenoma, or gastric polyposis, is as obscure as is that of other growths. Chronic gastritis, however, is accepted by all as a factor in its development. Kaufmann in his text-book on pathologic anatomy states that they develop either from a basis of chronic hypertrophic gastritis, or independently. Menetrier likewise considers chronic gastritis as an important etiologic factor, though not in all cases. He maintains that the condition usually occurs in advanced age, that atheroma of the blood-vessels is always present, and that the changes occur in the mucosa as a result of the involvement of its nutrition. In all of the cases reported, except the one by Norman, in which the patient was 34, and one by Chosrojeff, in which the patient was 36, the patients were over 59 years of age. Sex apparently played no rôle.

Macroscopically, the polyps in previous reports were small, pedicled, varying in size from that of a lentil to that of a pea, gray or reddish, depending on the blood-supply, of a soft consistency and never adherent to each other. The polyps sometimes numbered several hundred and were all about the same size, as if they developed at the same time, and after reaching a certain size stopped growing. They may develop from any part of the stomach mucosa. Brissard thinks that they occur with greatest frequency in the large culdesac, while another writer thinks that they develop chiefly from the pepsin-producing glands. Menetrier recognizes two anatomic forms, according to whether the hypertrophy and hyperplasia involves the excretory or deeper part of the tubular glands. In the first type lobulation is more apparent and cysts more common, owing to obstruction of the excretory ducts by the connective tissue. If they develop in the deeper parts of the glands, the polyps have a more uniform appearance, and tubulation is less pronounced. A mixed type may also occur. The portion of mucous membrane of the stomach which is not involved in the polyp formation usually shows the macroscopic characteristics of a chronic gastritis. Enlarged lymph-nodes are often encountered.

Polyadenomes en nappe (Menetrier) is a very rare condition, only three authentic cases having been reported, one by Andral and two by Menetrier. In this condition, instead of the hypertrophy and hyperplasia being limited to a part, it involves equally the entire mucous membrane in an area, so that the membrane

develops in large plaques and not as polyps. The involved area is usually from two to five times the normal thickness of the mucous membrane. The consistency is as soft as that of normal mucous membrane; the membrane develops in large folds, one pressing on another, and forms deep furrows. It develops parallel to the long axis of the stomach, and a sharp line of distinction is always present between the normal mucosa and the diseased area. The pylorus is never involved.

So far as my investigations go, there has been but one other case, besides the one here reported, in which a diagnosis was possible prior to operation or necropsy, namely, that of Chosrojeff.

History.—Mr. X. consulted me in December, 1904, on account of indigestion. There was nothing of importance in the fam-

Fig. 1.—Roentgenogram of the stomach, taken about fifteen minutes after the ingestion of an ounce and a half of bismuth in a pint of sour milk, with the patient in the standing posture.

ily history. As a child he had always been in perfect health. He never drank to excess, but had smoked from ten to twelve cigars daily for a number of years. His wife had given birth to one living child, which died in infancy and was said to have been syphilitic, and had two miscarriages, two and four years after marriage. Twenty years previously the patient had an abrasion on the right hand, which persisted for three months, and was said to have been a primary syphilitic lesion; shortly after this a small sore appeared on the penis, which existed but a few days, and was pronounced a soft chancre. There were no secondary manifestations of lues.

In 1891 a cloudiness of the vision of the left eye suddenly developed which was attributed to grip. The oculist questioned the patient carefully with reference to syphilis. Prior to this periodical indigestion of a mild character had occurred for several years. The patient also experienced "nervous

attacks" in which he became greatly depressed. When he consulted me in December, 1904, he was complaining of indigestion, characterized by belching and peculiar pressure sensations which seemed to "radiate from the stomach upward through the chest and arms." These symptoms seemed to bear no relationship to the character of the food taken, but usually occurred one or two hours after meals, unaccompanied by pain or vomiting. At this time 26 pounds in weight had been lost within a period of four months. The appetite was not good. The bowels were regular, but not free, sometimes compelling the use of enemas. At times there was palpitation, which the patient attributed to "gas in the stomach." There was no dyspnea, edema or any nocturnal urination. From time to time dizziness was experienced on sudden change of position.

dilated vessels. There was no tumefaction, though the epigastrium bulged slightly and presented slight resistance to pressure. Inflation of the stomach revealed no enlargement or displacement. There were a few enlarged lymph-nodes in the left anterior cervical region and also in the groins. On the dorsum of the penis was a small non-indurated scar and a scar was also on the left hand, the result of the lesion previously mentioned. The urine was negative. Hemoglobin, 67 per cent.

After a test-breakfast removed Dec. 14, 1904, 1½ ounces of slightly acid contents were recovered, with free and combined hydrochloric acid entirely negative. Lactic acid was mildly positive. There was a slight increase in mucus. Microscopically, nothing was abnormal. Test-breakfasts taken Dec. 16, 18, 20 and 24, 1904, and Jan. 10, 13 and 24, 1905, constantly revealed normal motility with complete achylia gastrica and a slight increase of stomach mucus. The diagnosis of chronic, atrophic gastritis was entered at this time into the records.

Treatment and Course.—This was prior to the days of the Wassermann reaction, so in view of the history of the case and the presence of abnormal fundus findings as reported by the oculist, iodids were given, which were tolerated so badly that the patient soon discontinued them. He took an extended trip, and on his return home stated that he felt fairly comfortable. I did not see him again for a period of five years, during which time the general condition was such that he felt no need of consulting a physician.

The patient returned in February, 1909, complaining of pronounced gastric symptoms, characterized by intense pressure in the epigastrium from one-half to one hour after meals, followed by "gas pains," three or four hours later. He was often compelled to get up during the night and walk the floor until he could free himself of the gas accumulation.

Physical examination at this time revealed no particular variation from the previous examination, except, perhaps, that the general nutrition was much improved. There was, however, a marked change in the gastric contents, which still contained no hydrochloric acid and no pepsin, but abounded in large quantities of mucus. Many analyses were made between February and October, 1909, with practically the same findings. The mucus was much like finely divided, slightly coagulated white of egg. It was practically impossible to lavage the stomach long enough to obtain wash-water free from mucus. Microscopically, there was an increase in yeast, though there was no branching. Almost invariably

there were a few red corpuscles, many undigested and partially digested leukocytes, many long and short bacilli and at times traces of pepsin, varying from 14 per cent. to 40 per cent. (Hammerschlag). There was no food retention. Numerous examinations of the feces revealed no blood.

The patient obtained his greatest relief through lavage, which was given now at frequent intervals for several months. The quantity of mucus that could be obtained from the fasting stomach was simply amazing. Not infrequently the wash-water was tinged with blood toward the end of the process of lavage. This, naturally, was assumed to be due to the irritation of the diseased mucosa by the tube. In view of the

Fig. 2.—The stomach removed at necropsy showing the entire mucosa, with the exception of a small area near the cardiac orifice, literally covered with polyps; a, a large grape-like bunch of polyps, a portion of which had invaginated the pylorus.

Examination.—The patient was a man of medium height, of a florid complexion, weighing 178 pounds, still well nourished though showing evidence of loss of weight. The pupils reacted normally to light and accommodation. The patellar reflexes and the sensorium were normal. There was a low systolic murmur at the apex of the heart, not transmitted. There was no enlargement of the heart. The lungs were negative. The liver was palpable two fingers' breadth below the free margin of the ribs on deep inspiration and was of normal contour and consistency. There was no pain on pressure in the right iliac, right hypochondriac or epigastric region. At the diaphragmatic insertion were numerous small

fact that the ocular conditions were gradually getting worse, and on the assumption that the gastritis might be of luetic origin, the patient was given mercury injections and iodolbin, which were very well tolerated.

During the latter part of 1909, while the stomach was being lavaged, a small polyp about the size of a pea was expressed through the tube. It was perfectly round, smooth and unaffected by digestion, and showed definitely where the pedicle had separated. A microscopic examination showed it to be a simple benign adenoma. This naturally led to the supposition that there might be other small polyps in the stomach.

In 1910 the patient complained frequently of vertigo and this was attributed to the eye condition. There were also occasional attacks of diarrhea. A blood examination at this time revealed hemoglobin, 80 per cent.; leukocytes, 21,800; erythrocytes, 6,192,000; polymorphs, 67 per cent.; large mononuclears, 16 per cent.; lymphocytes, 12 per cent.; eosinophils, 5 per cent., and red cells normal in appearance.

Test-breakfasts examined from time to time in 1910 revealed perfect motility, immense quantities of mucus containing many leukocytes, a total absence of hydrochloric acid and a mild degree of peptogenic power.

In the summer of 1911, while up North fishing, the patient developed an edema of the extremities and a rather marked degree of dyspnea, due to a relative insufficiency of the mitral valve, the result of overexertion. The condition improved under rest and digitalis, and when I saw him again in August he had a rather marked systolic murmur and intermittent heart-action, with an accentuation of the second pulmonic sound.

March 5, 1912, while in a hotel rotunda, the patient was overcome by a feeling of faintness and vertigo and fell unconscious. Shortly after this he vomited large amounts of blood, much of which was clotted. Before he could be removed to the hospital he had a large, perfectly black bowel-movement. Those present at the time think that he vomited at least a quart of blood. As soon as possible the patient was taken to the hospital. Temperature, 98.4; respiration, 20; pulse 120. There was no further vomiting, but the stools contained very large quantities of changed blood.

On the first examination at the hospital there was visible peristalsis and a palpable resistance in the pyloric region, much resembling the consistency of liver. On the following day with the hand resting lightly over the pyloric region to see if the peristalsis could be palpated, one could feel periodically the stiffening of the pylorus, and this was immediately followed by most peculiar movements under the hand, which resembled the movements of the small parts of a fetus. By stethoscope the movements could be heard as a distant rumbling noise followed by a squirting sound, as of a mixture of fluid and gas. The patient stated that the sensation could be experienced by fixing the mind on it, and, without any suggestion on my part, he described it "as the opening and closing of a valve." In view of this peculiar phenomenon, together with the previous findings of a small polyp in the washings, I concluded that a pedicled polyp had invaginated the pylorus, extended through into the duodenum and that either through laceration or the separation of a portion of it hemorrhage had resulted. The stools were watched carefully, and within three days a portion of a polyp about the size of a walnut was found, irregular in outline, with the surface partially digested. Sections were made which revealed the typical structure of a gastric polyp. A soft resistance could still be felt at the pylorus, but the peculiar movements described above became less apparent, though they remained more or less so throughout the early convalescence.

The treatment consisted in absolute rest, the application of ice-bags to the epigastrium, calcium chlorid by rectum, morphin hypodermically and fluids by rectum during the first forty-eight hours. Fluids were given by mouth on the third day in small quantities which were gradually increased until about the sixth day, when liberal amounts were taken.

Blood was apparent macroscopically for about ten days in the stools, but gradually disappeared, and within three weeks after admission to the hospital there were no chemical traces of blood to be found.

The patient complained more or less constantly of "gas in the stomach" and considerable distress, though of very little actual pain. The pulse remained around 90, until about the twelfth day, when it began to resume the normal. On the twenty-sixth day he sat up and was taking liberal quantities of soft food. At no time throughout his convalescence was there any evidence of food retention. There was no vomiting, no eructations, and in spite of the liberal quantities that were administered in an effort to build up the system as rapidly as possible, there was no stagnation.

March 31 his weight, which had usually been in the neighborhood of from 165 to 170, was 137½ pounds. Following the primary hemorrhage the hemoglobin was 35 per cent. A month later the hemoglobin ranged from 50 to 60 per cent., the erythrocytes 4,264,000. About this time the tertian type of plasmodium was found when a routine examination of the blood was made, and before treatment could exert sufficient influence, the temperature suddenly went up to 101, the patient had a severe chill, and the hemoglobin dropped to 40 per cent. Under quinin the plasmodia soon disappeared. This was during a month when we have no mosquitoes in St. Louis, and while there was no history of malarial infection, the plasmodia were evidently latent in the bone marrow and had probably been liberated as a result of the rapid activity brought about through the acute hemorrhage. By the early part of June the patient's condition was excellent. The hemoglobin was now 84 per cent., the erythrocyte count 6,488,000, and the blood-findings in general practically normal. The weight was now 158½ pounds, a gain of more than 20 pounds from the first weighing one month after admission to the hospital.

The fluoroscopic and roentgenographic examinations of the stomach revealed an almost total obliteration of the pars pylorica and the median portion of the fundus. The bismuth mass formed a shadow to the left of the vertebral column, showing a very irregular and indefinite outline. The bismuth in trickling through the interstices showed a very dim shadow of the right half of the stomach. The mottling of the bismuth shadow in this portion of the stomach suggested irregular masses in the stomach wall (Fig. 1). The length of time required for the stomach to empty itself was not determined, but there was apparently no retardation, as is evidenced by the bismuth shadow in the small intestines within fifteen minutes after the ingestion of the bismuth.

Though immediate operation following the hemorrhage was considered, the patient seemed such a poor risk and improved so rapidly on rest, diet and iron medication that it was decided to defer it as long as improvement continued. Early in June he felt quite himself again. On the basis of the previous literature and the palpatory findings following the patient's hemorrhage, it was thought probable that there was one large polyp attached near the pylorus and extending through it, and that there were probably a number of smaller ones grouped about the large one. It was hoped, therefore, that a gastrotomy could be done and the polyps removed individually or by a resection of the polyp-bearing area.

Operation and Result.—June 17 a median incision was made by Dr. Max W. Myer, a small amount of free fluid escaping. The liver presented itself, somewhat enlarged, but soft. Adhesions were found at the greater curvature about 3 inches from the pylorus, which when detached revealed an area of small, dilated vessels or varicosities on the peritoneal surface. There was some contracture at this point, which presented an hour-glass appearance. Palpation of the stomach revealed a large, soft, putty-like tumor mass easily the size of a man's fist, occupying the median portion of the stomach lumen. Further palpation revealed similar smaller masses filling more or less the lumen of the stomach up to the cardia. Only a small portion of the pyloric end seemed free from involvement. Resection seemed out of the question. The stomach was opened over the large tumor mass by an incision parallel to the greater curvature, the pyloric and the cardiac ends of the stomach first being clamped with rubber-covered clamps. When the stomach was opened a reddish-brown papillomatous growth began to deliver itself, and a tumor larger than a man's fist with much the appearance of a bunch of grapes was gradually delivered, the pedicle of which corresponded to the area of

contracture noted on the greater curvature before the stomach was opened. The pedicle was half the length of the index-finger. In addition to this, the mucous membrane of the stomach from the cardia to within an inch of the pylorus was literally covered with small pedicled polypi, varying from the size of a large pea to that of a hazelnut. In view of the extensive involvement of the mucous membrane, a complete resection of the stomach was practically impossible. In the meantime, the patient's condition became very unsatisfactory. It was decided, therefore, to remove the large papillomatous growth only. After this was done, an attempt was made to close the incision in the stomach, first overwhipping the mucous membrane with catgut. The stomach was closed with several layers of chromic gut, buttonhole stitch and finally with a layer of silk, stitching the omentum to the suture line.

The patient's condition when he left the table was very bad indeed. Shortly after returning to bed he vomited a considerable amount of blood which included several small polypi. There was constant oozing of blood throughout the day, either from the incision or from the pedicles of the detached polypi, and in spite of every effort the patient died about twelve hours after the operation.

Necropsy.—This revealed little of interest outside of the stomach (Fig. 2), which, together with a portion of the duodenum and of the esophagus, was removed, and contained a large quantity of clotted blood. There were enlarged lymph-nodes in the gastrophatic omentum and the mesentery. Intestinal polypi were not present.

I am greatly indebted to Dr. E. L. Opie for the following notes with the results of his macroscopic and microscopic findings.

Gross Examination.—Specimen consists of a portion of the stomach about 17 cm. long and 14 cm. wide. The greater part of the stomach wall is studded with polyp-like growths from 0.5 to 2 cm. in diameter. The mucous membrane between these masses is thrown into deep folds. There is an irregular mass about 12 by 6 cm. in diameter, resembling to some extent a bunch of grapes arising from the mucosa on the lesser curvature (?) about 5 cm. from the pyloric orifice. The nodules forming this mass are similar to those coming directly from the mucous membrane.

Microscopic Examination.—A section of a large polyp-like mass 1.5 cm. in diameter shows that it is attached to the underlying stomach wall by a pedicle 0.5 cm. across. The muscular coat of the stomach forms a regular layer below the polyp. The polyp consists of greatly hypertrophied mucous membrane, the muscularis mucosa and a loose connective-tissue core continuous with the submucosa and containing fairly large arteries and veins. The muscularis mucosa can be traced as a thick, doubtless hypertrophied layer at the base of the mucous membrane separating it from the core derived from the submucosa. On section the polyp exhibits three lobules, each of which contains an extension of the core, here consisting of connective tissue and muscularis mucosa. The hypertrophied mucous membrane which forms the main mass of the polyp consists of immensely hypertrophied glands, often with irregularly widely dilated lumina. Near the surface the dilatation and irregularity of the glands is greater than that deeper down. Here the gland consists of unusually high columnar cells with nuclei near the base. A very large proportion of the cells have the characters of goblet-cells and are obviously secreting mucus in large quantity into the dilated lumen of the gland. At the base of the mucosa in contact with the connective-tissue core, small glands resembling the pyloric glands of the stomach are abundant. The interstitial tissue between the glands contains cells in great number. These are in part lymphoid cells, in greater part plasma cells; eosinophils occur in immense number throughout the tissue, and polynuclear leukocytes in places are fairly abundant. The connective tissue forming the core of the polyp is the site of acute inflammation. Polynuclear leukocytes occur in large numbers and scattered among them in much smaller number are eosinophil cells.

Section through the stomach wall near the pylorus at a point where the mucous membrane is greatly thickened, although there is no definite polyp formation, shows hyper-

trophy and dilatation of the glands, inflammatory changes in the mucous membrane, hypertrophy of the muscularis mucosa, hemorrhage into the submucosa probably traumatic, and hypertrophy of the mucosa.

At the base of the mucosa are a number of widely dilated glands having the appearance of small cysts. The interstitial tissue of the mucosa contains an immense number of plasma cells, among which eosinophil cells are thickly scattered. Lymphoid and plasma-cells form collections of considerable size, particularly below the surface of the mucous membrane. The submucosa shows a mild inflammatory reaction containing scattered polynuclear leukocytes and eosinophil cells.

A section from the cardiac end of the stomach shows no hypertrophy of the mucous membrane. The glands appear to be displaced in part by accumulations of cells in the interstitial tissue which are most abundant immediately below the surface. Here lymphoid and plasma-cells are collected in immense number. Eosinophil cells are very abundant and form in places conspicuous collections. Parietal cells occur in scant number. Several cysts of small size occur and in some of these polynuclear leukocytes are fairly abundant.

The lesion corresponds with what is described as gastritis polyposa.

There is throughout the stomach both within the substance of the polyp and elsewhere evidence of chronic gastritis in marked degree. The relation of polyp formation to gastritis cannot be defined with certainty. On the one hand, the polyp may be regarded as a hypertrophy of the mucous membrane brought about by irritation associated with inflammation. On the other hand, the polyp may be regarded as a tumor or adenoma formed by proliferation of the glandular tissue. Localized hypertrophy of the mucous membrane will lead to the formation of the pedunculated polyp, the connective-tissue core being mechanically drawn into the nodule.

CONCLUSIONS

1. Though the diagnosis was made possible in this case through the presence of a small polyp in the wash-water during lavage, which also occurred in Chosrojeff's case, and the presence of a large polyp in the feces following the hemorrhage, it would seem that at least a probable diagnosis might be made in future cases without this conclusive finding.

2. The roentgenographic and fluoroscopic examinations in a case as extensive as this should always be helpful. I have not seen a similar picture in other conditions in my own experience, or in the records presented by others. The mottled appearance of the entire right half of the stomach, as though the bismuth were trickling through and around numerous masses, together with the irregular and indefinite outline of the stomach, could be produced only by such a condition as here described, or a most extensive malignant disease which would readily be differentiated by other means.

3. Achylia gastrica, also observed in Wegele's and Chosrojeff's cases, together with unusual mucus production (Chosrojeff), should always arouse suspicion. Ordinarily in achylia gastrica mucus is not encountered in the wash-water either in the large quantities here described or with the peculiar egg-white character, such as one would expect in the great multiplication of goblet-cells.

4. The repeated presence of fresh blood microscopically in gastric contents removed with care, or in the wash-water, is indicative of a redundant, vulnerable condition of the mucosa in which bits of tissue are readily removed by the tube. I have observed this more often in chronic gastritis than in circumscribed lesions such as ulcer or carcinoma. This should at least lead one to think of the possibility of polyposis. In this case, as in those of Wegele and of Chosrojeff, the clinical picture was that of a chronic catarrhal gastritis, with, perhaps,

fewer symptoms than are ordinarily manifested by gastritis patients.

5. In severe, acute, gastric hemorrhage in a patient with achylia gastrica, abnormal mucous production and normal or increased gastric motility, polyposis is more than probable.

6. Invagination of the pylorus by a polyp could hardly be mistaken for any other condition, after one has experienced the peculiar palpatory findings described in this case.

7. The etiology, of course, is very obscure, but it would seem probable that syphilitic gastritis was here the underlying cause.

Wall Building.

GROWTH AND EFFICIENCY OF HOSPITALS *

H. B. HOWARD, M.D.
BOSTON

My excuse for taking up any of your time is that I am appointed chairman of this section, and it seems to be one of the duties of the chairman to open the session with a few remarks. I can assure you that nothing that is of interest to general hospitals is foreign to my sympathy.

This section was formed, as most of you know, last year, and if its work shall prove to be of benefit to the general hospitals, it will grow and have its excuse for existing.

The general hospital is becoming an important factor in most communities, not only of this country, but also of the civilized world. It is almost startling to see what has been accomplished in the multiplication of these hospitals during the last few years.

The first hospitals in this country were the New York Hospital, the Pennsylvania Hospital and the Massachusetts General Hospital. It was nearly two centuries after the first white man landed on the shores of Massachusetts Bay that the first of these was built. The Massachusetts General Hospital was the third. Although the corporation was formed in 1811, the building was not ready to receive its first patient until 1820.

Few municipalities of any size now exist without their city hospitals. If there is not a city hospital, you will find a large private institution within its confines that is devoted to the public interest. I believe that it is not an exaggeration to say that there are twice as many hospitals in existence to-day as there were three years ago. If this is thought to be an exaggeration, you will be convinced to the contrary by talking with some of the firms that make equipment for general hospitals. You will find that all of these firms have been so loaded down with orders for equipment during the last three years that it has been impossible to keep pace with them. This being the condition, it is fortunate that the American Medical Association has formed a section that is devoted to this subject alone.

With this rapid growth, many of the hospitals are undoubtedly in a more or less crude condition, and meetings of this kind tend to perfect, refine and develop those that have been hastily constructed and perhaps carelessly organized into more healthy and common-sense institutions. No hospital can hope to grow in size or reputation unless it has a well-organized staff

and keeps the interests of its patients well in the foreground.

Too much cannot be said, and too much emphasis cannot be placed, on treating the applicants for admission to the hospital in the most dignified, gentle and courteous manner. It should be part of the education which the hospital owes to the community to impress all who knock at its doors with its sympathy and kindness toward them and their maladies. I know that it is a hard thing when you cannot accept patients and relieve them, so to conduct yourself toward them that they realize that you regret your inability to be of service to them; but this is one of the vital points to be insisted on at the very outset when any hospital throws open its doors to the public.

The care of the patient may be prompt, may be scientific and may be most effective, but a hospital will grow slowly in reputation if its promptness and efficiency in treatment and diagnosis are not carried forward with an attitude of perfect courtesy toward its patients.

The three things that our hospitals should stand for are (1) the care of the patients, (2) scientific investigation and (3) the education of physicians, nurses, orderlies, every one within its walls, and through them the community at large, concerning the various maladies that are brought within its doors.

I can well remember, when I was a student visiting the outpatient departments in the city of Boston, being impressed with the helplessness of the physician to relieve the various patients that applied. At that time there was not so much thought taken, or so much effort made, to give advice that the patient was able to carry out. A patient who after a careful examination was found to have tuberculosis was given the prescription, emulsion of cod-liver oil and told to spend his winters in the South. The fact that a man was a day-laborer and had a large family dependent on him for support and not a dollar at his command for traveling to the South apparently did not have much effect on the physician. I have seen a patient without a cent in his pocket loaded down with prescriptions which would have cost him two or three dollars to have filled.

When one of these patients applies to-day at our outpatient departments, first, a careful physical examination is made and conditions are noted; then, there is an investigation as to his ability to apply the treatment. Someone visits his home to see if it is possible for the patient to sleep outdoors. If it is not, some society produces him a sleeping-porch. If he cannot pay for the medicines that are needed, the hospital, as a rule, furnishes these medicines free. If he needs a few months in a tuberculosis sanatorium to begin with, this is provided for, and a social worker interests the various charitable organizations to look out for the family during this sojourn. When he is returned to the community, the same careful investigation is followed up. He is gathered into a class of tuberculous patients who meet evenings to get their directions and encouragement from the physician. So that in a short time this family and the neighbors have all become educated in what it is proper to do, not only to help a tuberculous patient to recover, but also to prevent others in the neighborhood from contracting the disease. The sputum is carefully looked after and destroyed, and the whole family is enlightened as to the dangers of carelessness in the conducting of such a case.

This progress has been made largely, if not entirely, through the hands of the general hospital. I have men-

* Chairman's Address before the Section on Hospitals of the American Medical Association, at the Sixty-Fourth Annual Session, held at Minneapolis, June, 1913.