the tongue was dry, brown, and hard; the urine, obtained by catheter (6 drachms), showed a large amount of albumin, with numerous hyaline and granular casts. He was treated with ordinary uræmic therapeutics and rapidly regained consciousness. In two weeks' time he had improved so much that he required no special treatment or attention. When first examined he showed Kernig's sign on the right and left sides, the greatest angle of extension being about 110° . On the eighth day of observation this sign could not be elicited on either side. This patient was only examined for the sign when he was lying in bed, the thigh being flexed upon the abdomen at an angle of 90° .

CASES IV. and V.—The two hemiplegics referred to were both cases in which the mentality was below par. They were both rather typical cases of right-sided paralysis, which condition had lasted in one case for over twenty years, and in the other for about fifteen years. The mode of onset as well as the subsequent course of the affection could not be ascertained. They were examined, both when sitting up and in a recumbent posture, and Kernig's sign was found present on the right side in each case. In one case the greatest angle of extension was 115° , and in the other 120° . In both cases when the thigh was not flexed the leg could be extended to an angle of 180° .

In concluding this report of 100 cases it is essential to state that in no positive case was there any joint involvement either osseous or tendinous; also that in all these cases the legs could be fully extended, and when the thigh was flexed the tendons became hard and tense. I have called no case positive in which the angle was more than 120° . The only case on this list in which the presence of Kernig's sign might have proved misleading by resembling meningitis was Case I., in which there were delirium, elevated temperature, rapid pulse, a red eruption over the chest and abdomen, with hyperæsthesia of the arms and legs. A leucocytic count of 5600, with positive Widal reaction, and two persons ill with typhoid fever in her home at the same time, made the diagnosis clear.

THREE CASES OF MENINGITIS IN WHICH KERNIG'S SIGN WAS PERSISTENTLY ABSENT.

By F. S. CLARK, M.D.,
RESIDENT PHYSICIAN, PHILADELPHIA HOSPITAL.

(From the service of Drs. Hughes and Salinger, of the Philadelphia Hospital.)

CASE I.—D. C., Italian laborer, aged twenty-two years, had a severe chill associated with pain in the abdomen and headache. In a short time he became unconscious, and when admitted to the hospital one day after the onset, his eyes were open and staring, the pupils were dilated, the right slightly larger than the left. Both reacted to light and accommodation. There was some rigidity of the muscles of the neck which increased to marked retraction of the head as his symptoms progressed. Photophobia and hyperæsthesia were prominent

symptoms. There were herpes labialis, and sordes on the teeth, but no eruption on the skin. The knee-jerks were normal; Kernig's sign could not be elicited in either leg in the recumbent or erect postures. Lumbar puncture was performed on two occasions, and a turbid fluid containing pus cells withdrawn. Microscopical examination of this fluid for bacteria was negative. There was a leucocytosis of 15,000, and a slight albuminuria with a few hyaline casts. The patient died on the fifth day without regaining consciousness. At the autopsy there was venous congestion of the meninges, and there was an exudate upon the under surface of the cerebellum and on the thoracic portion of the cord. The dura mater was adherent to the skull.

CASE II.—A. H., a salesman, aged twenty-three years, was suddenly attacked with vertigo associated with faintness and nausea. This lasted about five minutes. Similar attacks recurred at intervals, and he also complained of nose-bleed. Three weeks later when he was admitted to the hospital it was noted that his vision was impaired. He complained of dizziness; the gait was staggering and hesitating; the eye-grounds were reported to be characteristic of inherited syphilis, showing extensive exudative disseminated retinochoroiditis. Potassium iodide, however, produced no effect. Speech was hesitating; there was occasional vomiting; the knee-jerks were increased, and patellar clonus was present on the right side, and ankle clonus on both sides. Babinski's sign and Kernig's sign were not present. There was no retraction of the head, no loss of consciousness, but speech became more and more impaired, and the patient died in the fourth week of the disease. The autopsy showed the presence of a subacute tuberculous leptomeningitis. The membranes at the base of the brain were very adherent, and were separated with considerable difficulty. A few old calcified tubercles were found in the lung, and the right lobe of the liver showed some old healed tubercles.

CASE III.—T. B., an Englishman, aged forty-eight years, a tailor by occupation. Two days before admission he had become violent and had destroyed the furniture of the house. On admission he was unconscious, delirious, and there was continual twitching of the hands and feet. The thoracic and abdominal organs were normal. Four days after the onset of the disease he developed marked retraction of the head, but the reflexes were normal. Babinski's sign was not present and Kernig's could not be elicited. Lumbar puncture was performed on two occasions, and the tubercle bacillus was found in the clear fluid withdrawn. There was slight albuminuria and a few casts. The patient died on the seventh day, and at the autopsy miliary tubercles were found in the pia mater, and tuberculosis of the left lung, the liver, the spleen, and the kidneys.

Briefly summarized, there is one case of acute cerebro-spinal leptomeningitis whose bacterial nature was not determined, and two cases of tuberculous meningitis. In all of these cases Kernig's sign was repeatedly tested and was never present, no difficulty being experienced in any instance in extending the leg beyond an angle of 150° . This extension did not cause undue prominence of the hamstring tendons, nor apparently hurt the patient. The cases confirm the opinion that Kernig's sign is especially unreliable in tuberculous meningitis.