the atrophy of the right already spoken of. The left corpus is atrophied as the result of direct pressure by the aneurysm. Especially is this so in its caudal portion. The greater part of the left corpus restiforme had undergone a secondary degeneration as the result of the necrosis in the left cerebellar hemisphere. The changes in the cerebellum were most pronounced in the left hemisphere and peduncle. The pathologic changes of this case are described in great detail, the more important ones have only been mentioned in this abstract. The pathogenesis of these aneurysms of the vertebral artery are far from being understood. In this case the modification found in the medulla, the cerebellum, and the suboptical region are particularly important. The literature of the subject is carefully considered in the article, and the plates accompanying it add much to the understanding of the anatomical description.

SCHWAB.

THERAPY.

- 120 TRAITEMENT DE L'EPILEPSIE (Treatment of Epilepsy). Rommé (La Presse médicale, Mar. 21, 1900).

The author has followed de Fleury's method of caring for epileptics for eight years. The arterial pressure is regularly observed and measured, and urinary toxicity tested at stated intervals. The epileptic, being a depressed individual, is subject to dyspepsia and sluggish nutrition, therefore whatever affects arterial pressure may precipitate an attack. Combined bromides lower excitability of the cerebral cortex. Nocturnal cases should be given a single large dose at bedtime. Hydrotherapy, salt baths, massage, etc., with the injection of small quantities of normal salt-solution at various intervals, have been found of great benefit. No water is allowed with meals and the diet should seek to prevent the possibility of gastro-intestinal fermentation. To eliminate toxic substances, moderate exercise, purgatives, diuretics, and diaphoretics, should be employed. The author recommends quiet life in country villages for the epileptic.

CLARK.

- 121 TREPHINING FOR TRAUMATIC EPILEPSY. Lamboth (La Press med. Belges, Jan. 28, 1900).

Lamboth relates the following case: Youth of twenty-one years of age had severe epileptic convulsions for eight years. Thickening of left parietal, surface size of palm, developed toward cranium and much lessening its cavity. Trauma originated in a fracture *in situ*, possible separation of two tables having been followed by proliferation of diploë, or perhaps more probably an infectious diploëitis had occurred; ophthalmoscopic evidences of compressions absent. Patient had fallen from a height and struck his head; had been a typical epileptic for thirteen years, but development and intelligence had not suffered. Craniectomy was performed and a piece of bone 10 Cc. by 3 Cc. removed. It is stated that his attacks became much changed in type and immediately after the operation occurred at night only.

CLARK.

- 122 HEMI-CRANIECTOMY FOR EPILEPSY. Lamprasi (Annali di nevrolgia, 1900, XVII, fasc. VI, p. 414).

Lamprasi describes the following case: A youth, aged twenty years; epileptic. Left parieto-frontal region much depressed; attacks of convulsions typical, but without aura or monospasm; no mention of trauma. It was decided to remove one-half of the skull, and a horse-shoe incision was made from front of ear to rear of same; an osseous pedicle was left. Forty days after the operation patient had continued free from seizures.

CLARK.