attack, when the cystitis is well marked and he gets recurrent attacks of pyelonephritis, drainage of the bladder, by obviating the necessity of bladder washing and by abolishing intravesical tension, should prevent the recurrence of the ascending infection.

In some cases the Hamilton-Irving apparatus cannot be used, on account of the extreme flexion of the legs, and the spasmodic contraction of the abdominal and thigh muscles, set up by pressure on the abdomen.

During the first week or so, there is leakage alongside the tube in the suprapubic wound. The use of a suction apparatus such as that of Cathcart is of great value in keeping the patient dry in these cases. After this the wound contracts down on the tube, and a rubber plaque with a rubber cannula to carry the tube is strapped on over a small square of boric lint. This is retained in position by a rubber or canvas waist-band. In many cases it is sufficient to keep a No. 12 rubber catheter in the suprapubic wound and connect it by rubber tubing passing over the abdomen into a bottle lying at the side, in the hollow of the loin. A piece of adhesive plaster will keep the tube from slipping, and no leakage occurs alongside.

References.—1. Gaskell: The Involuntary Nervous System, 1916. 2. Müller: Deutsche Zeitschr. f. Nervenheilk., 1901, S. 86. 3. Bechterew: Die Funktionen der Nervencentra, i., 1903. 4. Justschenko: Verhandl. d. wissen.-ch. Versamml. d. Klinik für Nerven- u. Geisteskr., St. Petersburg, 1897. 5. Thomson Walker: Journal of Anatomy and Physiology, vol. xl. (third ser., vol. i.), April, 1906, p. 189. 6. Thomson Walker: Deuxième Congrès de l'Association Internationale d'Urologie, London, 1911. 7. Corner: Annals of Surgery, 1901, No. 34, p. 455. 8. Thomson Walker: Brit. Med. Jour., Sept. 13th, 1913, and Edinburgh Medical Journal, June, 1914. 9. Guyon et Albarran: Archiv. de Méd. Expér., 1897.

A METHOD OF CONCENTRATING ENTAMOEBA CYSTS IN STOOLS.*

BY J. W. CROPPER, M.B., M.Sc. LIVERP.;

AND

R. W. HAROLD ROW, B.Sc., F.L.S., F.Z.S.,

ASSISTANT LECTURER AND DEMONSTRATOR IN ZOOLOGY IN THE UNIVERSITY OF LONDON, KING'S COLLEGE.

(From the Laboratories of the Royal Society of Medicine.)

Introductory.

THE following investigations have been carried out on behalf of the Medical Research Committee and with the assistance of the Durning-Lawrence Fund, under the directorship of Lieutenant-Colonel Sir Ronald Ross.

One of the principal objects of our experiments has been the cultivation of the causative organism of amoebic dysentery, and, as is well known, the difficulties in the way of the accomplishment of this have hitherto proved insuperable, almost all observers now being agreed that the amoebæ which have been successfully grown are distinct from both *E. coli* and *E. histolytica*; in fact, the genus created by Chatton and Lalung-Bonnaire,¹ *Vahlkampfia*, has been frequently used, especially for the culturable intestinal forms, to emphasise their distinction from these two species. In a recent communication Penfold, Woodcock, and Drew² have described a method by which they have succeeded in causing the excystation of *Entamoeba histolytica*, but they also state that they have been unable to keep the excysted amoebæ alive for more than a comparatively short time, 42 to 44 hours as a maximum. This failure to obtain a successful culture they ascribe to the rapid growth of bacteria in their preparations, derived from the faecal matter not got rid of in the washings to which the cysts were subjected. The injurious influence of the products of putrefactive bacteria has, indeed, been already shown experimentally³ with cultures of amoebæ of the *imax* type. We have therefore since then devoted a considerable amount of time to the attempt to devise a method of completely separating the cysts from the faecal matter contained in the stools.

Such a separation as we have just mentioned has not, so far as we are aware, been accomplished by anyone in the case of amoeba cysts, but a modification of Bass's method of concentration described by Cochran⁴ has very nearly achieved it for helminth ova. This method consists in the preliminary emulsification of the faeces in water, and then centrifugalisation, after which the supernatant water is poured off, and the centrifuge tubes filled up with a 42.5 per cent. solution of calcium chloride in water, well shaken and re-centrifuged, when the eggs rise to the top of the tubes, away from the faeces and almost free from contamination.

On trying this method with *Entamoeba coli* cysts, we found that the cysts are far too fragile to withstand the osmotic currents set up by the concentrated solution, and incontinently crumple up; and a number of other chemical substances, such as syrup, glycerine, and albumin, which we have tried, have given precisely the same results, so that we feel that there is probably no specific gravity method of differentiation between cysts and faeces which will avail for entamoebæ, nor has any other method that we have yet found enabled us to achieve the complete separation of cysts and debris. We have, however, obtained a very considerable measure of success by the methods detailed below, and we are now able to obtain the cysts practically free from bacteria, and from the major part of the faeces.

The principal part of the experiments which we have carried out in investigating this method has been done with faeces infected with *E. coli* cysts, of which a constant supply has been available to us, but a sufficient series of determinations has been made with *histolytica*-infected material and with *Lambli*a infections, to show that it is equally applicable to these species, and we see no reason to suppose that any protozoal cysts would not be equally easily concentrated by it.

The technique of the method used to obtain a maximum concentration of the cysts is given first; subsequently we describe a method by which we have obtained a relative concentration without the use of any chemicals (which might possibly injure the cysts for cultivation purposes); and, finally, we describe a counting method which was devised to confirm the results of our concentration experiments.

I.—The Maximum Concentration of the Cysts for Diagnosis.

A lump of faeces, which should be at least 1 gramme in weight, is shaken up with about 30 c.c. of normal saline (0.8 per cent. solution of NaCl) per gramme of faeces for a sufficient time to disintegrate the mass into individual particles, and thereby form an emulsion which will only settle very slowly. This is best done on a mechanical shaker in a large flask or bottle of a capacity of at least four times the amount of fluid to be shaken, and we have found it necessary, in order to obtain the best results, to continue the shaking for a minimum of half an hour.

The emulsion is then poured into a separating funnel and shaken up, by hand, for half a minute with 10 to 20 per cent. of its volume of ether (ordinary methylated ether is perfectly suitable for this), after which the mixture is allowed to stand for a minute or two in the funnel until the two liquids have separated. The faecal debris absorbs ether and consequently becomes lighter than water, and, when separation is complete, lies in a mass at the top of the saline, immediately below the excess of ether. The cysts are not affected by the ether, and consequently remain in the saline beneath.

The saline fluid is then drawn off from the separating funnel, and is centrifuged at a slow speed for two or three minutes, the precise time required necessarily depending upon the rate of centrifugalisation and the size of the tubes. By this means the cysts will be brought down to the bottom of the tubes with the comparatively slight amount of faecal matter not taken up by the ether.

For ordinary diagnostic purposes the concentration now effected, which should be some 15 times as rich in cysts as the original material, will be found adequate, but, if desired, a still greater proportion of cysts can be obtained by decanting the supernatant liquid and filling up the centrifuge tubes with normal saline, shaking thoroughly, and submitting the tubes to fractional centrifugalisation, the material brought down in the first 10 seconds being discarded,

⁴ The Concentration of Helminth Ova from Faeces, China Medical Journal, vol. xxix., 1915.

* Read before the Section of Pathology of the Royal Society of Medicine, Nov. 14th, 1916.

¹ Amibe imax (*Vahlkampfia*, n. gen.) dans l'intestin humain. Son importance pour l'interprétation des amibes de culture. Bull. Soc. Path. Exot., Paris. Vol. v., 1912.

² The Excystation of *Entamoeba histolytica* (*Tetragena*) as an Indication of the Vitality of the Cysts, Brit. Med. Jour., May 20th, 1916.

³ Cropper, J. W., and Drew, A. H.: Researches into Induced Cell-reproduction in Amoebæ, McFadden Researches, vol. iv., April, 1914, John Murray, London.

and the whole time of centrifuging curtailed by half. By repeating this process two or three times the maximum concentration is obtained, and the débris still remaining consists almost entirely of particles of practically the same size as the cysts themselves.

The above is a concise account of the method and its results. We append a series of notes on the various processes.

It is desirable, where possible, to take a fairly large mass of faeces for concentrating, since there appears to be considerable variation in the number of cysts present in different parts of the same stool, and one of the special advantages of a concentration method is to minimise the risk of missing cysts not equally distributed.

The period given for mechanical shaking may seem extraordinarily long, but our experience is that unless the various particles of faecal matter are thoroughly washed off from the cysts the latter are removed with their adhering particles by the ether, and consequently do not appear in the final deposit. In fact, test experiments differing only in the length of time the faeces were shaken with saline showed at least twice as great a proportion of cysts in the final deposit after 30 minutes' shaking as after 15 minutes; and while in the latter case examination of the material removed by the ether showed a considerable proportion of cysts, they were almost entirely absent from it in the material shaken for 30 minutes. This length of time may, however, need to be modified in accordance with the character of the stool to be examined; our own investigations have been principally made on formed or sub-formed stools.

We have employed tap water, distilled water, and normal saline for emulsifying the faeces, and we think that saline is preferable. Whatever the method used, we have very frequently found that a considerable number of cysts have shown what we regard as osmotic collapse, their protoplasmic contents being withdrawn from the cyst wall and collected into a spherical mass about half the diameter of the cyst, but we are unable at present to determine the precise causes which have produced this result, as both the occurrence of the phenomenon is irregular and the proportion of cysts affected has varied very considerably under the same conditions. We find, however, that there tends to be a smaller proportion of cysts showing this effect in emulsions for which normal saline has been used than in tap water or distilled water emulsions from the same faeces and prepared concurrently under identical conditions. The ether employed subsequent to emulsification may be to some extent responsible, but we have more than once observed collapsed cysts in material which has not been treated with ether. In our experience this collapse does not interfere with the identification of the cysts, and we have never seen it save in *E. coli*.

We find that the proportion of the material removed by the ether varies considerably with the character of the stool used, but with an ordinary formed stool at least 90 per cent. of the weight of faeces employed is extracted. To give a concrete example, 10 grammes of faeces were emulsified, treated with ether, and centrifuged in tubes of known weight, and the deposit, after pipetting off the supernatant fluid, weighed 0.595 gramme.

Should the faeces contain fruit pips or other large heavy particles not removed by the ether it is advantageous to transfer the fluid, after treatment with ether but before centrifuging, to a tall test-tube or similar vessel and to permit the heavy particles to settle for a minute before decanting into the centrifuge tubes. The cysts themselves settle very slowly, and it has been found that the proportion which fall to the bottom with the large particles in one minute is negligible.

The time which we have found most suitable for centrifuging with 15 c.c. tubes is either $1\frac{1}{2}$ minutes at 1200 revolutions per minute or $2\frac{1}{2}$ minutes at 600 revolutions per minute. This will bring down practically all the entamoeba cysts from the fluid, but only a small proportion of the bacteria present come down in this length of time. In the case of lamblia a more thorough centrifuging is needed to bring down all the cysts.

The special advantages which this method possesses for diagnosis are obvious. The cysts are much more easily noticed and far more easily identified in the concentrated material than in preparations from untreated faeces. The saving of time spent in the examination of scanty or doubtful infections is considerable, and the concentrate from a comparatively large mass of stool can be examined

in a short time: there is, further, the added advantage that the possible errors attendant upon the examination of a single loopful of faeces are almost wholly eliminated. This is especially useful in the case of suspected infections in which the ordinary preliminary examination has given a negative result, and we think the method will also prove to be of great value for the final examination of carrier and of apparently cured cases before discharge.

An illustration is given (Fig. 1) of the appearance under the low power of a field from a concentration prepared in this manner. A count of the original faeces, from a smear made in the ordinary way without concentration, gave 48 cysts in 117 fields under the same lens; after concentration there was hardly a field to be found in which there were not a dozen cysts, and the average number per field was between 16 and 17.

II.—The Relative Concentration of Cysts for Cultivation.

We feel that there are distinct disadvantages in the use of ether for treating material to be employed for cultural purposes, as the prolonged contact of the cysts with ether and with ether-saturated water may easily have some deleterious effect, and we have consequently devised a method of concentration which does not require chemicals. The first attempt to effect this was made by filtering an emulsion of faeces in water or normal saline through very fine silk cloth, some varieties of which show microscopically a mesh which is but little larger than an entamoeba cyst. It was hoped in this way to remove all particles larger than the cysts, but in practice the meshes became clogged so rapidly with débris that no filtrate was obtainable, even with the aid of a pressure pump. The smallest mesh which can be employed satisfactorily for the filtration of faecal emulsions is about 40μ in diameter, and the method by itself is of little value for concentration purposes. Combined, however, with subsequent centrifugation it forms a ready means of obtaining a large proportion of the cysts free from all particles save those which approximate in size to that of the cysts, the larger fragments being held back in the deposit of the silk and the minuter débris remaining suspended in the tubes after centrifuging.

It was found, however, that no modification of this method would give so large a proportion of cysts to débris as the ether method described above, and estimations of the number of cysts per gramme of stool, as determined from the deposit in the tubes by our method of counting, showed that they were considerably less than the actual number present in the untreated faeces, a considerable number having been lost during the treatment, so that the concentration was only a "relative" one—i.e., with respect to the nature of the accompanying débris.

As finally adopted, our method follows in most respects that used by Penfold, Woodcock, and Drew, but as it differs in certain points of importance we think it desirable to give it in full, especially in order to emphasise the necessity of these or similar measures when undertaking any routine work in an endeavour to cultivate entamoebæ.

Ten grammes of faeces are shaken with 100 c.c. of normal saline in a bottle or flask on a mechanical shaker for 5 minutes to obtain a uniform emulsion. This is poured on to a layer of fine silk whose mesh is of the size mentioned above, and which is stretched on a tambour (such as is used by milliners), and the emulsion is gently and continuously stirred with a glass rod to prevent the clogging of the meshes of the silk. The residue left on the filter consists chiefly of large lumps and stringy matter, and is discarded. It contains only a small percentage of cysts, which can be still further reduced, if required, by washing. The filtrate (or a portion of it) is then centrifuged for one minute at a speed of 1200 revolutions per minute, the supernatant liquid poured off, and the volume made up again with normal saline. The tube is then well shaken, and again centrifuged as before. This process is repeated until the supernatant liquid is almost clear. Finally the deposit is shaken with 10 c.c. of normal saline and is allowed to stand for a few minutes. The upper portion is then poured off and is thoroughly centrifuged, and loopfuls of the deposit are used to make hanging-drop preparations for culture experiments. An illustration (Fig. 2) is given of a high-power view of the appearance of cysts and débris in such a preparation.

By this means cysts are obtained which have been thoroughly washed free from toxic bacterial products, acid substances, and other deleterious matter present in the original faeces. The cysts can be found easily with low

powers of the microscope, and are not subsequently lost, even though it is necessary to keep the preparations for several days in the incubator. *Amoeba limax* cysts submitted to this process have been subsequently cultivated without difficulty, and were evidently unaffected either by the shaking or the centrifugalisation.

It must be noted that the length of time, and the speed, of centrifuging considerably affect the percentage of cysts in the final deposit; and we have selected one minute at 1200 revolutions per minute as giving the best practical results. Apparently about two minutes is required to carry down all the cysts, but a much greater proportion of the finer particles of faecal matter and bacteria accompanies them than when centrifugalisation is confined to the shorter time. An additional method of removing any large particles that may still be present is afforded by centrifuging for 15 seconds at 600 revolutions per minute, decanting the fluid into other centrifuge tubes and re-centrifuging for one minute at 1200 revolutions per minute; the larger particles are brought down during the preliminary 15 seconds with but a very small proportion of the cysts, and this process, repeated four times, results in the larger debris containing only one-tenth of the proportion of cysts present in the original faeces.

We have not been able to separate the cysts from debris to any further extent except by picking off a single cyst in a capillary tube, which is a comparatively simple matter in the case of *E. coli*, and has been done repeatedly in our experiments.

III.—Method for Counting Entamoeba Cysts in Stools.

In THE LANCET of June 10 h, 1916. Miss Annie Porter⁵ described a method of counting lamblia cysts in stools by means of a Thoma-Zeiss hæmocytometer. Finding the volume of the chamber too small for counting entamoeba cysts, owing to their comparative scarcity—viz. about 1 cyst per 400 squares in an ordinary degree of infection—we devised the following method, which is described in the stages in which it is carried out.

A uniform emulsion of 10 grammes of stool (selected from several different portions of the bulk) is made in 100 c.c. of normal saline (i.e., 10 per cent. weight ÷ volume) by shaking on a mechanical shaker for 10 minutes. All soft masses are broken up in this way, and the sediment which falls consists only of gritty particles, the so-called "false sand," which microscopic examination shows to contain no cysts.

A capillary pipette is made and is calibrated to deliver 20 c.mm. in a length of about 2 inches. It is provided with

being retained in a pipette made according to the above instructions when the mixture is blown out: in fact, we have not yet found a single cyst in the washings out after use.

Several glass slides, 3 inches by 1½ inches, having been prepared with a ring of vaseline 1 inch in diameter, a measured volume (20 c.mm.) of the emulsion of faeces is drawn into the pipette and is placed in the centre of the ring.

A cover-slip ruled in squares, such as is used with Böttcher's slides, is then allowed to fall gently, with the engraved face lowermost, flat on to the drop of emulsion, and, if necessary, is pressed down to diminish the depth of the layer of liquid. The count is made with a low power. The cysts which happen to be in any portion of the drop which extends beyond the squares are easily counted, and are indicated in the count by the letters A, B, C, &c. The squares themselves are numbered as shown (the reverse way up) in the diagram (Fig. 3).

We have made sufficient experiments to establish the utility of the method for counting entamoeba cysts in faeces, and we see no reason why its employment should not be extended, with such modifications as are found necessary, to the counting of other protozoa, either in their natural surroundings or under different conditions of cultural environment.

It has been pointed out to us by Sir Ronald Ross that in counting methods there are considerable statistical errors, the magnitude of which depends, roughly in an inverse ratio, on the actual number of things counted. Thus it is necessary to count a fairly large number of cysts in order to reduce the error to a reasonable percentage, and the present method has been devised to render this possible. The statistical tables compiled by Ross and Stott⁶ show, for example, that if only 68 cysts are counted the probability is 9 to 1 that the error does not exceed 20 per cent., and that, if the error is required to be within 10 per cent., 271 cysts must be counted. It is advisable, therefore, to use an emulsion for counting purposes as thick as possible, consistent with ability

to recognise the cysts without difficulty. For counting *E. coli* cysts we find a 1 in 10 emulsion the most suitable, and in an average infection we count 100 cysts. With *Entamoeba coli*, as in Miss Porter's cases of *Lamblia* infection, there are considerable daily variations in the number of cysts present in the stools, ranging in a case we have been studying from 1500 to 60,000 per gramme.

Summary.

1. Methods of concentrating entamoeba cysts from faeces are described, suitable for diagnosis and cultivation experiments respectively. 2. A very high concentration, suitable for diagnosis, has been obtained by emulsifying faeces and treating the emulsion with 10 to 20 per cent. of ether. This removes the greater proportion of the faecal matter, while the cysts remain in the saline used for emulsification, and are collected by subsequent

FIG. 3.

Diagram to show the method of counting entamoeba cysts in stools. The edges of the large drop of 20 c.mm. of faecal emulsion are indicated by the thick irregular line, and all cysts in this drop are counted with a low power. $\times 1\frac{1}{2}$.

FIG. 1.

Maximum concentration of *E. coli* cysts in stools by the ether method. One field. Low power: $\times 50$.

FIG. 2.

All visible particles and a 4-nucleate cyst of *E. coli* in one field of an emulsion prepared by the method of relative concentration for cultivation experiments. High power: $\times 275$.

a rubber teat. It is well to have several of these pipettes ready to hand, so that successive counts can be made with a dry one without any unnecessary delay. The pipette supplied with Gowers's hæmocytometer, and other similar ones which are on the market, can be used for the purpose, but they have a somewhat narrow bore and frequently become clogged with debris. There is little risk of any cysts

⁵ An Enumerative Study of the Cysts of *Giardia* (*Lamblia*) *intestinalis* in Human Dysenteric Faeces.

⁶ Tables of Statistical Error, *Annals of Tropical Medicine and Parasitology*, vol. v., No. 3, December, 1911.

centrifugalisation with but little accompanying débris. 3. A second concentration method is described which does not necessitate the use of ether, and which is therefore more suitable for preparing cysts for cultivation experiments. This consists of filtering the emulsion through silk, and subsequent fractional centrifugalisation. 4. The difficulties in the way of enumerating entamoeba cysts are discussed, and the method found most suitable, by means of the coverslips used with Böttcher's slides, is described.

A CASE OF SUPPURATIVE MENINGITIS WITH GLYCOSURIA SIMULATING DIABETIC COMA.

BY FRANK E. TAYLOR, M.D., B.S. LOND.,
M.R.C.P. LOND., F.R.C.S. ENG., D.P.H. CAMB.,

PATHOLOGIST AND BACTERIOLOGIST TO THE QUEEN ALEXANDRA
MILITARY HOSPITAL, MILLBANK; LECTURER ON BACTERIOLOGY,
UNIVERSITY OF LONDON, KING'S COLLEGE;

AND

W. H. MCKINSTRY, M.B., C.M. GLASG., D.P.H. LOND.,
TEMPORARY CAPTAIN, R.A.M.C.

ON Oct. 13th, 1916, the following case was admitted into the Queen Alexandra Military Hospital, and was thought worthy of being placed on record on account of several interesting points to be discussed later.

Second Air Mechanic A—, of the R.F.C., aged 36, with four months' service, had been ill for 24 hours, complaining of severe pain in the right side of the chest, cough, and difficulty of breathing. On admission the patient appeared ill, with flushed face and marked dyspnoea. The temperature was 102° F., the pulse-rate was 112, and the respirations were 38 per minute. Examination of the chest revealed diminished expansion of the right lower chest, with stony dullness and absence of tactile vocal fremitus and breath sounds of the right lower lobe. Typical "rusty" sputum was being expectorated, and the case was diagnosed as one of pleurisy and pneumonia. Poultices were applied to the chest. Calomel gr. iii. and pulv. ipecac. co. gr. x. were ordered *statim*.

Oct. 15th: The patient had had a bad night. Pain was intense, so that he could not lie comfortably in bed. The temperature still ranged from 101° to 103°. The pulse was of good quality, with a rate of 100. The respirations were very rapid, rising to 60 per minute. The physical signs in the chest remained the same, no signs of fluid being present. There was no cyanosis. The heart and pericardium were normal. The application of leeches to the right side of the chest caused great relief of the pain almost immediately, and a stimulant expectorant mixture was prescribed. Oct. 16th: Patient appeared to be greatly improved, said he felt quite well, and was very cheerful. The temperature was 102°, pulse 100, and respirations 36. There was very little cough and no expectoration.

Oct. 17th: The ward sister noticed some twitching of the patient's face; he had previously complained of intense headache. He vomited immediately afterwards and became very collapsed with incontinence of faeces. Coma rapidly supervened, with dilated pupils which did not react to light. Conjugate deviation of the eyes was noted first to the left and then to the right. All the superficial reflexes except the conjunctival were lost. Later he became very restless, constantly picking at his face, which wore a perpetual frown. The respirations rose to 64 and presented the appearance of air-hunger. The pulse still remained of good quality, being 100 to the minute. He had retention of urine, and a catheter was passed, drawing off 20 oz. On examination this contained a trace of albumin and a few hyaline and granular casts, as well as a large amount of sugar (not estimated quantitatively), but acetone and diacetic acid could not be detected. No distinctive odour of his breath could be recognised by any of several observers who examined him with this in view. Further, no history suggestive of glycosuria—excessive thirst, wasting, or polyuria—could be elicited on cross-examination of his wife; his health before this attack seemed to have been very good. An attempt made to administer alkalies subcutaneously was unsuccessful on account of the patient's restlessness, and rectal salines met with no better results. He could not take anything by the mouth. His condition remained much the same during the night.

Oct. 18th: The respirations were now 80 and the pulse 144, and inclined to be irregular. His colour had much improved. The pupils were no longer dilated. He appeared to be temporarily paralysed on the left side, and had conjugate deviation of the eyes to the left. Incontinence of urine now

developed, but a sample of urine was collected and found to be quite free from sugar, whilst the quantity of urea present was normal. Venesection was performed, 20 oz. of blood being removed, and two pints of normal saline containing 1 per cent. sodæ bicarb. were injected into the vein. Later the eyes became deviated to the right, the pulse got weaker, Cheyne-Stokes respiration developed, the colour became cyanosed, spasms of the jaw and muscles of both arms were noted, and the patient died at 3 P.M. Just before death the temperature rose to 107°. The temperature chart is here shown.

A post-mortem examination was made 36 hours after death. The body was well nourished and showed no signs of external injury except a hæmatoma about the size of the palm of the hand between the umbilicus and left anterior superior spine of the ilium. Thorax: The whole of the pleura covering the right lung was covered with a layer of semi-membranous purulent exudation about a quarter of an inch thick, which could easily be peeled off. The left lung and pleura appeared healthy. There was no free fluid in the pleural cavity. The heart appeared normal and there was no pericardial effusion. The vessels about the base of the heart and the structures in the mediastinum were normal. Abdomen: All the organs, including the kidneys and liver and pancreas, appeared healthy. Brain: On removing the outer coverings of the brain pus was found mapping out the sulci on the anterior, superior, posterior, lateral, and internal surfaces, and also following the course of the blood-vessels. When the brain was removed the base was found free from pus except for a small area on each side where the middle meningeal artery takes its origin. The upper surface of the cerebellum was thickly covered with pus, particularly in the region adjoining the mid-cerebrum. The other surfaces of the cerebellum were free from pus. No pus could be found about the medulla. The brain when cut into showed no sign of abscess or excess of fluid in the ventricles.

Smears taken direct from the pus in the sulci, stained by Gram's method, showed long chains of streptococci as well as staphylococci. Streptococci were also found in smears from the cerebro-spinal fluid and in the blood drawn direct from the seared heart. Media of agar, broth, and trypsin legumin agar inoculated from the blood and the pus from the brain and pleura, incubated at 37° C. for 24 hours and examined, showed growths of streptococci, *Staphylococcus aureus*, and a large Gram-positive sporing organism, the last being, no doubt, due to post-mortem invasion of the tissues.

Although the nature of this case was plainly revealed by the post-mortem examination, the recognition of the cause of the sudden coma formed a difficult and perplexing problem during life. It was in the attempt to settle this question that the urine was sent for examination, and when it was found that it contained a large quantity of sugar the diagnosis of diabetic coma was raised. The absence of polyuria and of acetone and diacetic acid in the urine, as well as the absence of the characteristic sweetish acetone-like odour of the breath, rendered this diagnosis a very questionable one.

The question of uræmic coma was next raised, seeing that albumin, though in small amount, as well as a few hyaline and granular casts were found in the urine. The determination of the percentage of urea was then suggested, and for this purpose a further specimen of urine was obtained on the following day. The percentage of urea was found to be normal (i.e., 2 per cent.), and this fact was taken as disposing of the diagnosis of uræmic coma. A quantitative determination of the sugar was then decided upon, but in this specimen no trace of sugar or other Fehling reducing substance could be obtained, showing that the glycosuria was

Temperature chart showing rise of temperature to 107° before death.