VACCINATION OF TUBERCULOUS PATIENTS AGAINST SMALLPOX.

RESULTS IN TWO HUNDRED CASES

LOUIS MARK, M.D.
MOUNT VERNON, OHIO

A review of the small amount of literature on the subject of vaccination against smallpox of tuberculous patients and personal consultation with several men of authority reveal the fact that it is apparently a matter of common agreement that vaccination should not be done unless it is extremely necessary for the safety and welfare of the patient.

The results obtained in compulsory vaccination of about 200 patients with pulmonary tuberculosis seem strongly to contradict this idea.

The reason for the compulsory vaccination was that a virulent case of smallpox broke out in the institution and the patient had been mingling with all the other patients for about ten days prior to the time of the eruption and the making of the diagnosis.

REPORT OF CASE

W. Mc., man, aged 34, white, admitted, Aug. 2, 1918, with the ordinary history of a tuberculous breakdown, was well nourished and was only 5 pounds underweight, weighing 170 pounds. He had no cough and very little expectoration, but what there was gave a positive reaction for tubercle bacilli. Chest examination revealed very little. The patient was classed as incipient (Turban Stage I).

August 12—ten days after admission—the patient's temperature rose to 102.6. He complained of headache, general malaise, nausea, vomiting and constipation. Chest examination disclosed no cause for the rise of temperature. On the 16th his temperature dropped to subnormal and he developed an eruption. A diagnosis of smallpox was made, which was confirmed by the state dermatologist.

VACCINATION OF THE OTHER PATIENTS

As the patient had been mingling with the other patients and employees in the main dining hall and cottages, compulsory vaccination was decided on, and all patients, regardless of stage or activity, were vaccinated, August 17. The number vaccinated was 200, made up of seventy in the first stage, eighty-three in the second, and forty-seven in the third. The majority had some activity and some elevation of temperature. Among these patients there were forty who either had never been vaccinated before, or had been vaccinated with negative results.

Three weeks after vaccination all cases were closely observed, and it was found that there were about 50 per cent. positive vaccinations. Fifteen per cent. of the patients had sore arms and 8 per cent. were "on tray" for one or more meals. On the ninth day after vaccination, one patient developed a typical varioloid eruption about the neck and also had a positive vaccination on the arm. Several patients who had to be kept on tray had a rise in temperature which lasted for a day or two and was present at the time the redness, swelling and soreness of the arm were the greatest. There were no instances in which patients complained of any conditions pertaining to the chest that could in any way be traced to the vaccination. The majority of the patients were examined three or four weeks after the vaccination, and in no case were there any changes incompatible with the usual course of tuberculosis.

In reviewing the results obtained from this series of vaccinations, it would appear that when a case of smallpox breaks out among tuberculous patients and there is a possibility of a spread of the disease among the patients, compulsory vaccination is advisable.

The results obtained in far advanced cases could not be differentiated from the results obtained in the first and second stage cases, and it would seem, therefore, that these cases should not be exempted when considering vaccination. Although it would not be advisable to vaccinate all patients against smallpox when first admitted to an institution, yet it should be done at once when the necessity arises.

The smallpox patient made a normal recovery from his smallpox and was discharged from the institution within three months as an apparently arrested case of pulmonary tuberculosis. His sputum was negative for tubercle bacilli on discharge. No other cases of smallpox developed at the institution.

STREPTOCOCCUS PNEUMONIA

JAMES G. CUMMING, M.D. (ANN ARBOR, MICH.)
Lieutenant-Colonel, M. C., U. S. Army

CHARLES B. SPRUIT, M.D. (BOSTON)
Captain, M. C., U. S. Army

ERNEST J. ATEN, M.D. (CORAOPOLIS, PA.)
First Lieutenant, M. C., U. S. Army
NEWPORT NEWS, VA.

At Fort Sam Houston, Texas, during the winter of 1917-1918, it was noted that the majority of pneumonia cases presented a clinical type not often seen in civil practice. The patients were highly septic; the pneumonia, for the most part, followed measles, and when not fatal, it often terminated in empyema. It was further observed that the usual type of lobar pneumonia was often complicated by relapses, which were attributed to delayed resolution. The frequency of what we shall call, for the present, septic pneumonia, and its occurrence as a complication of measles, led to an investigation as to the causative organism in this type of the disease, as well as in the complicated, so-called lobar pneumonia. It is well known that the hemolytic streptococcus is almost exclusively the causative organism of complications in scarlet fever cases; and since pneumonia is not an uncommon complication of this disease, we were led to map out a bacteriologic investigation of measles and pneumonia cases; this included the identification of this organism.

As previously reported, it was readily determined that a large percentage of measles patients harbored the hemolytic streptococcus in their throats on entering the hospital, and further, that this organism was found in practically all specimens of sputum from septic pneumonia patients, also in the pus from empyema patients, and finally in the blood stream shortly before death.¹ Furthermore, this organism was found in the throat, sputum, empyema pus, and blood stream of many patients suffering from so-called lobar pneumonia.

It would seem that this organism is of sufficient importance to warrant a brief consideration of its epidemiology. Epidemiologic studies of the occurrence of

1. Cumming, J. G.; Spruit, C. B., and Lynch, Charles: The Pneumonias: Streptococcus and Pneumococcus Groups, J. A. M. A. 70: 1066 (April 13) 1918.

more or less extensive milk-borne epidemics of tonsillitis or septic sore throat in England, as well as in this country, chiefly in the vicinity of Boston, Chicago and Baltimore, make plain the epidemiology of hemolytic streptococcus infection. It is the opinion of Smith and Brown that the organism is of human origin, that it gains access to the cow's udder from infected milkers, and that milk is the chief, if not the only, disseminator of this disease on a large scale, in these milk-borne epidemics. Smillie reports its frequent occurrence in the throats of scarlet fever patients and in the blood stream in the fatal cases. It has been isolated from the blood stream of septicemia patients following trivial wounds, and is now believed to have been the chief cause of sepsis following childbirth.

In pursuance of our plan of investigation of throat swab cultures from measles patients, it was at first believed that we should determine the prevalence of type pneumococci as well as hemolytic streptococci. But owing to the early discovery that the hemolytic streptococcus was the predominant, if not the sole, cause of pneumonia following measles, we later limited our investigation to the identification of hemolytic streptococci.

HEMOLYTIC STREPTOCOCCUS CARRIERS

With the view of determining the percentage of hemolytic streptococcus carriers among measles patients on entrance to the hospital, throat swab cultures were taken in 452 cases. These fell into two groups, one showing Types I and II,² and the second showing Type III.³ Of these, 294, or 64 per cent., showed Types I and II hemolysis, while 158, or 35 per cent., showed Type III hemolysis. Among the 294 patients showing Types I and II hemolytic organisms, thirty, or 10 per cent., developed these complications: otitis media, four; bronchopneumonia, five; tonsillitis, seven; scarlet fever, three; lobar pneumonia, two; acute bronchitis, seven; erysipelas, one; cellulitis, one.

In contrast to the foregoing group, which shows 10 per cent. of complications, we find that among 158 patients showing Type III hemolytic streptococci, sixty-five, or 41 per cent., developed complications. These were: bronchopneumonia, fifty-two; empyema, nineteen; tonsillitis, nine; otitis media, eight; arthritis, four; acute bronchitis, five; mumps, one; lobar pneumonia, one; cellulitis, six. These eighty-four complications occurred among the sixty-five patients harboring streptococci.

If only the pneumonia cases are considered in these two groups of findings, it will be noted that among those measles patients from whom organisms showing Types I and II hemolysis were isolated, only 2 per cent. developed pneumonia; while among those harboring Type III hemolytic streptococcus, 32 per cent. developed pneumonia. The incidence in percentages was sixteen times greater among streptococcus carriers.

Considering the total pneumonia cases in both groups of throat findings among measles patients, it is shown that of the 452 cases, 12 per cent. developed pneumonia, and over 94 per cent. of this 12 per cent. occurred among streptococcus carriers.

In this series of cases it has been established that 35 per cent. of the measles patients were hemolytic streptococcus carriers on entering the hospital, and

that 33 per cent. of these developed streptococcus pneumonia; while among the 65 per cent. of patients who were noncarriers, only 6 per cent. developed this complication.

In contrast to the high percentage of hemolytic streptococcus carriers among measles patients, it is pointed out that among 168 swabs from average throats, there was found only 6 per cent. streptococcus carriers. This percentage corresponds to that obtained by Smillie in his investigation of hemolytic streptococci in Boston. It is a significant fact that there are six times more streptococcus carriers among measles patients than among average well persons.

It might seem that the streptococcus in the throat of a carrier prepares the way for measles virus infections, and that the latter infection, in turn, produces a condition—lowered resistance—that favors streptococcus pneumonia—an extension of this infection to the lung, the pleural and pericardial cavities, and the blood streams, joints, etc.; or it is possible that both infections occur simultaneously through the same avenue of transmission. In order to determine whether or not measles patients are streptococcus carriers prior to measles infection, it is suggested that at least 1,000 recruits at some camp be examined for this organism, dividing these into carriers and noncarriers, and that they be kept under observation for the occurrence of measles in the two groups. Are streptococcus carriers more susceptible to virus infections?

BACTERIOLOGIC FINDINGS IN ANTEMORTEM PLEURAL EXUDATES AND BLOOD OF PATIENTS WITH PNEUMONIA FOLLOWING MEASLES

Of ninety-two examinations of pleural exudates from patients in the pneumonia wards, organisms were found in seventy-four specimens. Of forty-six specimens from patients diagnosed as having lobar pneumonia, 61 per cent. proved to be pure hemolytic streptococcus, and 39 per cent. pneumococcus. Of twenty-eight bronchopneumonia specimens, 86 per cent. were streptococci and 14 per cent. pneumococci. By consolidating these two groups we find that 72 per cent. were hemolytic streptococci and 28 per cent. pneumococci. From these figures it would appear that clinically there is either a considerable error in the differential diagnosis of the pneumonias, or that a large percentage of the lobar types are complicated by streptococcus infection.

From patients with pneumonia following measles, thirty-eight blood specimens were examined. Of these, twenty-nine were negative, while nine, or 24 per cent., were positive for hemolytic streptococci. With few exceptions these positive specimens were obtained within twenty-four hours before death, showing that the blood stream invasion with this causative organism was a late event. In the series of necropsies to be reported, 52 per cent. of heart's blood cultures were positive.

BACTERIOLOGIC FINDINGS AT NECROPSY

In thirty-four necropsies on nine lobar pneumonia and twenty-five bronchopneumonia patients, streptococci—Type III hemolysis—were isolated from twenty-one lung specimens, twenty-three pleural exudates, fourteen pericardial exudates, and sixteen heart's blood specimens.

In twenty-seven, or 80 per cent., of the specimens from serous exudates or heart's blood, the strepto-

2. Small area hemolysis representing pneumococcus and *Streptococcus viridans*.

3. Large area hemolysis representing *Streptococcus hemolyticus* colonies.

coccus was isolated from pure culture, and in four, or 12 per cent., of the specimens it was coexistent with the pneumococcus; furthermore, it was found in one or more of these specimens in thirty, or 88 per cent., of the necropsies, lobar pneumonia cases included.

The streptococcus was isolated from each of the twenty-five necropsies on bronchopneumonia patients. Of these, 75 per cent. were known to have followed measles.

Of the nine lobar pneumonia postmortem specimens from clinically diagnosed lobar pneumonia cases, in only four instances was the pneumococcus unaccompanied by the streptococcus. In two others only streptococci were found, and in the remaining three there were both pneumococci and streptococci.

In only one necropsy on a patient that had had pneumonia following measles was a type epidemic pneumococcus found, and this was coexistent with the hemolytic streptococcus; while in the nine lobar pneumonia cases, hemolytic streptococci were found, either in pure culture or coexistent with pneumococci, in five, or 55 per cent., of the necropsies.

It would appear that the hemolytic streptococcus was the immediate cause of death in thirty, or 88 per cent., of the thirty-four cases, lobar pneumonia cases included; and since this organism was found in fairly pure cultures in all lung specimens as well as in pure cultures, in either serous exudates or the heart's blood or both, in cases of pneumonia following measles, it may be concluded that the hemolytic streptococcus is the causative organism in this complication following measles, and moreover, that it was the immediate cause of death in a large percentage of so-called lobar pneumonia cases.

SWAB AND SPUTUM SPECIMENS

Since, in this series of necropsies, the hemolytic streptococcus was identified as the causative organism of bronchopneumonia following measles, a series of sputum specimens was examined in conjunction with throat swabs. Of the swabs, 76 per cent. were positive, as were also 87 per cent. of the sputum specimens. In 55 per cent. of the cases the swabs as well as the sputum specimens were positive; while either swab or sputum was positive in 100 per cent. of the cases. These findings show the importance of taking both swab and sputum specimens.

HEMOLYTIC STREPTOCOCCI IN REMOVED TONSILS

Of 142 removed tonsils, 82 per cent. showed hemolytic streptococci. The tonsil acts as a reservoir for the organism. On removal of the tonsil, the throat rapidly clears of this, the most common of all secondary invading organisms.

IDENTIFICATION AND CLASSIFICATION

On account of the unquestioned prevalence of streptococcus pneumonia, both as a complication of measles and as a coexistent or a terminal infection in lobar pneumonia, it was decided to examine for streptococci sputum specimens submitted to the pneumococcus division for type identification. A total of ninety-two specimens was examined, and forty-nine, or 53 per cent., of these showed Type III hemolytic streptococci. Of these ninety-two sputum specimens, pneumococci Types I, II or III were identified in only 9 per cent. of the cases. Owing to the prevalence of streptococcus infection as a secondary and fatal involvement in lobar pneumonia, as well as in the virus infections, there is

need for the instruction of all laboratory workers in the use of the hemolytic plate in the diagnosis of pneumonia. The clinician should insist on a report of the presence or absence of streptococci, as well as of type pneumococci. The importance of these findings in the sputum is substantiated in nine so-called lobar pneumonia cases, in which, at necropsy, hemolytic streptococci were isolated from the heart's blood or serous cavities in 55 per cent. of the specimens.

We are led to believe from our investigations that a diagnosis of Type IV pneumococcus should be made guardedly; furthermore, since massive lobular pneumonia may be incorrectly diagnosed as lobar pneumonia, and as lobar pneumonia involvements may be associated with streptococcus infection, which at Fort Sam Houston gave a mortality of 82 per cent. during January and February, we believe that all specimens of sputum should be examined for hemolytic streptococci as well as for pneumococci. Since the pneumonias are of two main groups—pneumococcus and streptococcus—and as these may be coexistent, the isolation of the causative organism or organisms in pure culture is essential to group as well as to type identification.

The value of routine streptococcus identification will become of particular importance in the event of the successful production of antistreptococcus serum.

PREVALENCE OF HEMOLYTIC STREPTOCOCCI

	No.	Positive for Hemolytic Streptococci, Per Cent.
Swabs from measles cases	857	35
Swabs from lobar pneumonia	165	69
Sputum specimens from lobar pneumonia	92	41
Swabs from scarlet fever	192	87
Swabs from otitis media	204	74
Swabs from mumps	147	37
Swabs from tonsillitis cases	79	54
Cultures from removed tonsils	142	82
Nose swabs from carriers	69	55
Average percentage of streptococcus carriers among foregoing cases		59
Throat swabs from average healthy individuals	168	6
Total	2,135	

The clinical classification of the pneumonias is important, though less so than the classification from the bacteriologic point of view. From the findings here presented it would seem that the disease may be classified according to the causative organisms into two main groups, secondary streptococcus or pneumococcus pneumonia following virus infections and primary pneumococcus which may be followed by secondary streptococcus pneumonia.

We confirm the contention of Smith and Brown that the hemolytic streptococcus Type III hemolysis is a definite type, in that the colonies produced by us in 10 per cent. horse blood agar were uniform in character, and that 150 strains secured from necropsies, throat swabs, and sputum specimens gave the characteristic sugar reactions. Since the hemolytic streptococcus has proved to be such a common cause of death, in the accompanying tabulation we present in brief the percentages of prevalence of this organism in the 1,967 specimen examinations.

It is interesting to note that nose swabs were taken from sixty-nine known carriers, and that 55 per cent. of these proved positive for streptococci. This finding, in conjunction with our experiments showing that streptococci in droplets are forced into the air during the act of coughing, might lead one to believe that

droplets as well as indirect contact infection would be a serious problem in measles and scarlet fever wards. To gain some information on this point, the dust from seven wards was examined, with negative results. Although ward dust may be quite free from streptococci, yet our observations in the surgical wards would indicate that bed linen and garments soiled with streptococci may act as a transmitting agent in the case of wound infection. Among clean operative cases the wounds in twelve became infected with hemolytic streptococci. Throat swab examiners showed that all these patients were streptococcus carriers. Auto-infection, either through soiled linen as a medium or through the blood stream, would in these cases appear to be the only two avenues of infection; and of the two, the former would seem to be the more possible.

From the results of throat swab examinations of twenty-eight attendants in the streptococcus wards, it was found that 25 per cent. of these were hemolytic streptococcus carriers. If this finding alone is considered, one would conclude that streptococcus pneumonia may be called a hospital infection. But this is not so, because it was shown that 94 per cent. of all streptococcus pneumonia patients were streptococcus carriers on entrance to the hospital. In addition, it was found by the examination of convalescent measles patients that streptococcus infection rarely occurred in the hospital wards. Infection of ward attendants is naturally the result of prolonged exposure and close association, and from the results of our investigation it would seem that cross-infection among patients seldom occurs. In order to prevent the possibility of cross-infection in the wards, which we do not believe takes place to any great extent, patients harboring hemolytic streptococci should be isolated as soon as possible after admission.

In the large cantonment hospitals where a large number of patients are cared for, it is preferable to have three or more wards—one of these used exclusively for a detention ward, where regulations providing for the sheet method of separation, the individual face mask, and the frequent washing of hands of attendants are rigidly enforced; other wards may be designated as streptococcus and clean wards.

In the streptococcus wards, the usual methods of disinfection and the practice of scrupulous cleanliness of hands should be insisted on in order to protect attendants against infection. In our opinion, the sheet and individual mask method of protection is of value only in the detention ward, or when it is impossible on account of lack of laboratory facilities to classify the patients into carriers and noncarriers.

CONCLUSIONS

From the evidence here presented it will be seen that:

1. The hemolytic streptococcus was present in the throats of 35 per cent. of all measles patients on entrance to the hospital, and that 94 per cent. of measles patients who developed bronchopneumonia belonged to this group of carriers, while only 6 per cent. of noncarriers developed pneumonia complications.

2. This streptococcus was found in pure cultures at necropsy in the serous cavities and heart's blood of all measles-pneumonia patients, and in 55 per cent. of lobar pneumonia patients.

3. This streptococcus is the same as the beta streptococcus of Smith and Brown.

4. Type III hemolytic streptococci isolated from the serous cavities and heart's blood were always large zone hemolyzers.

5. The presence of hemolytic streptococcus in the throat and sputum of so-called lobar pneumonia patients and in the serous cavities and lungs at necropsy shows that in the classification and treatment of the pneumonias, the streptococcus must be as diligently sought for as the type pneumococci.

6. Laboratory men should acquire a technic for the recognition and isolation of the hemolytic streptococcus from throat swabs and sputums of measles and all pneumonia patients.

7. In practically all cases (94 per cent.), the hemolytic streptococcus is acquired *before* entrance to the hospital.

Prevention consists in: (a) the pasteurization of all milk supplies, thus eliminating the possibility of septic sore throat followed by the carrier state; (b) the removal of tonsils of those individuals who are identified as streptococcus carriers; (c) the control of measles epidemics by the early recognition and isolation of patients; (d) the segregation of streptococcus carriers in hospital wards; (e) scrupulous cleanliness of the hands of attendants and patients, and (f) the development of a reliable vaccine and serum treatment.

TWO NEW DIAGNOSTIC METHODS FOR PATIENTS WITH CAR- DIAC DISEASE

JULIEN E. BENJAMIN, M.D. (CINCINNATI)

AND

ERNEST R. BROOKS, M.D. (CLEVELAND)

Captains, M. C., U. S. Army; Members, Cardiovascular Board
CAMP FUNSTON, FORT RILEY, KAN.

1. TACHYCARDIA

In the recent articles on the subject of so-called soldiers' heart, latterly termed "neurocirculatory asthenia," it is noted that this syndrome is encountered in the civilian as well as in the soldier. This fact is borne out forcibly if one is on the alert during the entrance examinations of recruits. Since the beginning of the very large draft increments, every man entering the service has been examined for evidence of cardiovascular disease. This work, formerly done parenthetically by the tuberculosis board at Camp Funston, is now conducted by medical officers of the cardiovascular board, who have had a rare experience in examining many subjects and in the detection of suggestive signs of cardiovascular disorders. At an average, during a large draft, 1,500 men pass through their hands each day. Two medical officers, working in shifts of two hours each, are responsible for this work. From 5 to 7 per cent. of the total number (1,500) each day are referred to the cardiac room for suspected cardiovascular disease. This room is manned by two officers. Here the subjects have a searching examination, the results of which are recorded, if necessary.

The most difficult and trying task in the preliminary examinations is the proper decision in cases showing various grades of tachycardia. They represent an enormous number of cases. When one considers that at this receiving station, which represents a