hypothenar muscles placing the thumb and little finger in the position described. The index and ring finger respectively are passively drawn over by the displacement of the thumb and little fingers.

Austregesilo, A. SEXUALITY AS A FACTOR IN THE NEUROSES AND PSYCHOSES. [Arch. Bras. d. Med., Vol. 9, 1919, No. 2.]

It is the experience of many years of observation as a neurologist and psychiatrist that the factors of sexuality play a predominant rôle in the neurotic and psychotic. He states that in every form of dementia, toxic psychoses, psychoneuroses, as well as in all types of imbecility and idiocy anomalies in the genital sphere are to be found in almost every case, if not in all. Erotic elements are evident everywhere in the delirious. This is inevitable he declares since the history of mankind is a record of war and religious mysticism, both of which are expressions of the sexual instinct. Heretofore the problems of sexuality have been recorded chiefly by poets, novelists and artists of all ages, but scientists have been very reserved until Freud's definite stand.

Brown, Wm. HYPNOSIS, SUGGESTION AND DISSOCIATION. [British Medical Journal, June 14, 1919.]

The author says that there is an overwhelming consensus of opinion that the one satisfactory method of treating the various forms of functional nervous disorder is that of mental analysis and reëducation which can be summed up in the term "autognosis," or self knowledge. He believes that hypnosis can be used to great advantage in this plan of treatment, but not, as formerly, for the purpose of reinforcing suggestion. Its field of usefulness lies in bringing into the patient's consciousness all of the circumstances through which he passed at the time of the development of his nervous disorder. By inducing light hypnosis and causing the patient to go through all his initial experiences these are thus brought into his consciousness and many of his disabilities are at once removed, those most readily removed being mutism, paralyses, contractures, and tremors. Various disorders of the vegetative nervous system are also much benefited, even more in fact than paralyses of the voluntary muscles. The extent to which the conditions are relieved depends upon the completeness with which the patient's original experiences are revived. The method has proved most satisfactory in early cases, but is not available in chronic cases in which there is marked fixation of symptoms.

Frank, A. EMOTIONAL DISTURBANCES IN CHILDREN. [Corresp. f. Schw. Aerzte., May 10, 1919.]

This article gives some illuminating examples of the unconscious sexual motivation which lies behind incomprehensible behavior in children. One case history is as follows: W. had been a bright boy and

prize winner in the school. Without warning, he suddenly turned to a most abandoned and perverse character. He lied and stole, obtained money by most precocious artifices so that his father had to warn the tradesmen that he was quite irresponsible. The boy was flogged and talked to but to no purpose. He was subjected to all kinds of penalties, and even the aid of the police was enlisted. The author was some time in gaining the boy's confidence. He catered to his weaknesses, such as love of chocolate, and finally learned the secret which the boy let out involuntarily and then bitterly regretted. Either the mother had ignored its full bearing or had chosen to be silent about it. While the boy was still well behaved a beautiful servant girl had been hired and had struck up an innocent friendship with the boy. At first he simply preferred the girl to his mother as a companion, but the affection at last ended in the boy's seduction by the girl. The mother then promptly discharged the servant as soon as she learned of her misconduct, although perhaps not fully aware of the degree of intimacy. The boy blamed the author for extorting his secret, cursed and raged at him and would not see him again. The moment the girl was discharged the boy "went bad" and showed hatred, first to the mother and only later and to a less degree to the father. The patient was finally sent to a small private institution, where he showed a marked irritability and hatred toward his mates. After two months of careful psychoanalytic and general treatment according to the author's modified form of psychoanalysis, a mixture of Forel's and Jung's techniques the boy became tractable and behaved as if nothing had happened. He had acquired such insight into his case that he took the physician back into favor. He finally made a healthy readjustment to his home and school.

Etienne, G., and Richard, G. BLOOD PRESSURE AS INFLUENCED BY WAR EMOTIONS. [Paris médical, August 9, 1919.]

These authors report the results of several years' observation of the effects of shelling in the town of Nancy on blood pressure. All violent shocks were found to affect the blood pressure, generally bringing about a brief stage of hypertension, lasting five or ten minutes, followed usually by one of hypotension lasting from a few minutes to two or three hours. The average rise of systolic pressure was eight to twelve millimeters of mercury, and of diastolic pressure, three or four millimeters. Where the emotional shocks are repeated, *e.g.*, if the town is shelled throughout the day, each successive shock finds the blood pressure a little higher than it had been before, and subsequently the return of the pressure to normal may be slow, especially in old persons, in whom several weeks may elapse before it reaches normal. Where the shocks are repeated for weeks or months, the systolic pressure shows a constant elevation of from ten to forty millimeters, and of the diastolic,