til very recently, almost entirely superseded all other local applications. I refer to pyrogallic acid.

I do not think that pyrogallic acid is by any means so well known as an application for the relief of psoriasis as is chrysarobin. If I may judge by the infrequency with which its virtues are mentioned in the journals (although I believe all recent text-books speak of it), it is not in general use. But it is, in my opinion, one of the very best remedies we have for the cure of cases of psoriasis of average severity. Employed in the form of ointment, of the strength of one-half to one drachm of the pyrogallic acid to one ounce of simple ointment, the effect produced by it is almost as rapid and decided as that brought about by chrysarobin, without the accompanying discoloration. A slight blackish staining is all that is produced, and the ointment can even be employed in the scalp without markedly discolorizing the hair, if applied cautiously. It is desirable, however, not to apply soap or alkalies at the same time, because this causes a more permanent and deeper stain.

Pyrogallic acid cannot be used in extensively generalized psoriasis, when large surfaces are affected by the disease, without a certain amount of danger from absorption, as indicated by strangury and olive-green or tar-colored urinary sections. With care, however, and the occasional suspension of the remedy for short periods, I believe this remedy could be employed even in universal psoriasis with good effect.

One more external application in psoriasis remains to be spoken of—namely, naphthol. This drug, a derivative of coal tar, was introduced into use several years ago by Kaposi, of Vienna, as a sort of substitute for carbolic acid. He recommends it very highly in psoriasis, in the form of ointment, about eighty grains to the ounce, and I have used it in this and other strengths, and also in alcohol and oil, with fairly good effect.¹

Naphthol is not so active in its effect as chrysarobin or pyrogallic acid, but it is much more agreeable and is, I think, peculiarly well adapted for employment upon such parts as are exposed to the view, as the face and hands. Like pyrogallic acid, it must be used with caution over large surfaces, as absorption with toxic effects may be produced.

It remains to mention briefly two or three methods of application of these remedies which have recently been brought forward. The first of these is the treatment by medicated gelatine, which was introduced by Prof. Pick, the well-known dermatologist of Prague. My attention was first drawn to this by a pamphlet which Prof. Pick kindly sent me, in which he gave an account of his earlier experiments with medicated gelatines, but I have not as yet had an opportunity of testing this method of medication as I should desire. I may say, however, that the method does not seem to me calculated to prove convenient and popular in private practice. I had for some time been making some experiments in my service at the Polyclinic in the preparation of gelatines impreg-

nated with chrysarobin and pyrogallic acid, but without much satisfaction, when Dr. Chas. L. Mitchell, the well-known pharmacist of this city, sent me some excellent preparations of his own, which seem to be perfectly adapted to the purpose for which they are intended. A bit of one of these gelatine sticks is cut off and placed in a water-bath, where it soon melts into a clear homogeneous fluid, which may then be applied to the lesions of the skin by means of a paint-brush. The advantages claimed are cleanliness and transparency. The coating of gelatine does not rub off on the clothes, and is therefore not so dirty as an unctuous application. A fresh coating can be painted on every day or two as the old layer wears off. The chief disadvantage of this method of treatment is that it requires apparatus which is not convenient to carry about, nor can the patient be trusted to employ it at his discretion. My own experience is that in psoriasis, at least, the gelatine applications are not active enough. I have not, however, used them with sufficient frequency to pronounce a positive opinion.

Recently a solution of chrysarobin in collodion has been recommended in the treatment of psoriasis by Dr. George H. Fox, of New York, and several dermatologists have confirmed his statements with regard to its efficacy. I have employed this preparation in one or two instances, but it has seemed to me so much less active than the chrysarobin ointment that I have not been encouraged to continue its use. It has one great advantage over the gelatine applications, however, and that is that it can be applied extemporaneously, and without the paraphernalia which must accompany the use of the gelatine.

A few weeks since a pamphlet by Prof. Auspitz, of Vienna, reached me, in which that distinguished dermatologist recommended liquor gutta-percha as a vehicle for the application of chrysarobin. I at once obtained a ten per cent. solution, or rather emulsion, of chrysarobin in liquor gutta-percha, and happening to have a case of psoriasis of the face and scalp under treatment, I gave some to the patient to apply once daily. The effect was so happy as to encourage me very much to hope that we have in this preparation the most convenient method of applying chrysarobin yet devised; and as chrysarobin is, after all, the most efficient local agent in the treatment of psoriasis yet brought forward, I have no hesitation in urging the trial of this preparation on any one who may have a case of psoriasis under treatment. It is to be noted, however, that the same watch must be kept upon the skin for fear of exciting dermatitis as when the ointment is used. Only when the liquor gutta-percha dries, which it does very quickly, there is little or no danger of rubbing the chrysarobin over the good skin, nor is there much danger, if any, of staining the clothing.

BLOOD-LETTING.

BY GEORGE HAMILTON, PHILADELPHIA.

In the JOURNAL OF THE AMERICAN MEDICAL ASSOCIATION for September 22d, is an article entitled "Good Remedies out of Fashion," by Dr. C. T.

¹ In a communication read before the American Dermatological Association, last month, and published in the *American Journal of the Medical Sciences* for October, I have given the results of my experience in the use of naphthol in various diseases of the skin, psoriasis among the number. I may refer to that paper for further details as to the action of the drug in this disease.

Hare, President of the Metropolitan Counties Branch of the British Medical Association, in relation to emetics and blood-letting.

In regard to the utility of emetics in the cases adduced there are probably very many who coincide with the opinions of Dr. Hare, while, in reference to the propriety of bleeding in the congestive conditions alluded to, and others of a similar nature, a vast majority of practitioners, including nearly all the younger, would, perhaps, withhold their assent. The signs of the times, however, seem to indicate that a beneficial change of opinion upon this subject is about to occur; and the writer cannot but express his gratification that another distinguished authority has added his testimony in favor of a practice that, abused as it may have been in past years, is likely to assume its true position as an not only useful, but indispensable therapeutic measure in the cases to which Dr. Hare has referred. The earnest, impressive injunction of Dr. H. is that the physician, in the dangerous conditions he refers to, will not lose time in the employ of means wholly inadequate to the emergency, but at once resort to that one measure which, in the practice of the most experienced physicians, has been proven to be more successful than all other agencies combined.

Following the example of Dr. Hare, a case in illustration will now be given. A man of 22 years, unusually muscular and strong, after a few hours of chilliness was seized with a most violent and protracted chill, and in a very short time great oppression and stricture of the chest, with extreme difficulty of breathing, ensued. Early the next morning the patient came under my care. He lay upon his back, gasping for breath to such a degree that, if the motion of a fan was suspended for a moment, he would, with difficulty, exclaim, "Fan, or I will suffocate!" Face pallid, tips of the ears and nose cold, the hands, feet and legs in the same condition, the nails livid, the pulse very feeble and frequent. Inexperienced and alarmed, Dr. Thomas F. Hewson, my preceptor, was called to the patient, and at once directed me to bleed; but, for several minutes, a few drops only of dark blood could be had; yet by placing the hand in hot water the blood began to flow, and soon changed to a more natural color; the pulse increased in strength and became slower, the pallor and anxious expression of the face were less marked, and the patient said he felt easier. The usual warm applications and a stimulating liniment were applied to the feet, and moderate portions of carb. ammonia and wine were prescribed. The abstraction, however, of the carbonized blood had already produced reaction, and no matter how violent this reaction, or subsequent inflammation might be, this condition would be infinitely preferable to the profound passive congestion that oppressed the heart and lungs, paralyzed the action of the brain and entire nervous system, and thus, momentarily, endangered the life of the patient. Dr. Hewson, promising to return in the evening, said that this was by far the most strongly marked case of approaching pneumonia that he had ever seen; and now, strangely enough, after more than a fifty years experience, no such case has occur-

red in the practice of the writer. At the evening visit the whole aspect was changed; reaction was fully established, the pulse excited, gasping greatly diminished, expression of face less anxious. As the oppression of the chest, and now, for the first time, some degree of pain was complained of, Dr. Hewson, anticipating a severe attack of inflammation, directed the vein to be reopened, from which florid blood bounded out as if from the vein of a man in robust health. The subsequent march of the disease, contrary to what had been feared, was not as severe as some others coming under my notice. To bleed under the condition just stated would perhaps not be thought of by an immense majority of physicians, but would probably be derided or ridiculed; but neither derision nor ridicule can successfully confront irrefutable facts. But some may say in objection: Is it not a fact that the case adduced is, by your own admission and that of Dr. Hewson, exceptionally rare, and therefore of little practical importance? In answer, it must be admitted that cases of such violence are seldom met with, but many others, endangering life in a lesser degree, do occur, not only in pneumonia, but in several other diseases, sometimes ushering in the attack, oftener in the latter stage. In the destructive epidemics that ravaged Europe several centuries ago, the deaths would often occur so quickly that it was impossible to refer many of them to the violence of fever or inflammation; whilst the opposite condition, profound venous congestion, stifling, as it were, vitality at the fountain head, would readily explain the fearfully sudden mortality. Again, in the cholera of 1832, many of the slaves in Louisiana suddenly fell victims to the malignant nature of the disease, without the occurrence of vomiting or purging. In malignant intermittent, not very often seen in this country, death may occur so soon after invasion that no rational cause can be given other than that of profound congestion.

Physicians in this section of the country are, fortunately, not often called to cases of this character, but when they do occur the course to pursue is that so emphatically commended by Dr. Hare. In this connection the writer may state that nearly all who, for a few years past, have adopted and written upon this mode of practice, are found to occupy distinguished positions as practitioners, authors or lecturers. In a physiological and pathological point of view the cases in illustration, by Dr. Hare, perhaps more decidedly the case of the young man attacked with pneumonia, are of extreme importance. In disease, the greater the divergence from the normal physiological state, the more distinct, impressive and instructive is the pathological condition; and when, as in an ordinary case of pneumonia, where inflammation and fever may be violent, the contrast between this and the congestive abnormal is still more marked and instructive. The normal action of the brain, spinal marrow and nervous system depends, entirely, upon the stimulus of oxygenated, vitalized blood; and the physician who abstains from the abstraction of carbonized, in fact poisonous blood, must do so in ignorance of the real nature of the case, or from dread of the wide-spread sentiment in opposition to bleeding,

both in and out of the profession. Another class of cases of congestion is frequently seen, of an entirely different character, and occur under very different conditions. They are generally noticed in the latter stages of disease, in fever, inflammation, or in chronic affections; and whilst they may exhibit no startling signs of immediate danger, they are, doubtless, the final cause of more deaths than occur in the former class. As a rule, and by common consent, bleeding is totally inadmissible in these cases, unless done in the first stage of the disease, and the strength of the patient yet is unbroken.

It was not the purpose, nor is it within the scope of this paper to enter upon the subject of bloodletting in ordinary inflammation, and especially in pneumonia. The difference, or rather the antagonism, of opinion in regard to the treatment of this disease, and the conflicting statistical reports of success, based upon the use or neglect of the lancet, may not be regarded as complimentary to the profession. Inflammation and its attendant fever does not occur without a cause, a noxious agent; and hence, such inflammation and fever must be regarded, in tendency, at least, as conservative; and so long as inflammation does not become so violent as to threaten the integrity of tissue, and organs essential to life, there would appear to be no necessity for the abstraction of blood; whilst in cases of great intensity, unless reasons of special, imperative character forbid, blood should be taken at the earliest moment, and to such extent as the constitution and existing conditions will admit. The danger so often spoken of in opposition to bleeding sinks into insignificance compared with that resulting from intense, unrestrained inflammatory action. The ability of the constitution to bear great loss of blood is exemplified in the fearful condition to which women in abortion, or post-partum, are at times reduced, and from which they often recover with marvelous rapidity.

CATHETER BROKEN OFF IN PROSTATIC PORTION OF URETHRA--EXPEDITIOUS REMOVAL.

BY ARTHUR L. WORDEN, M.D., DES MOINES, IOWA.

I notice, in your issue for Sept. 15, the paper of Dr. Hard, in which he recommends a sharp, screw-pointed instrument for removing broken catheters from the urethra. This cut represents an instrument which once helped me out of such a difficulty without the patient even knowing what had happened.

I had occasion to pass a soft rubber catheter, No. 11, into the bladder of a colored man. After drawing his urine and attempting to withdraw the catheter, imagine my dismay at having it break off in the prostatic region. Without mentioning the fact, I immediately passed the urethral forceps, and easily securing a grasp upon the broken end, at once withdrew the fragment entire.

On another occasion, I was unfortunate enough to have a pledget of absorbent cotton slip off my applicator and remain in the cavity of the uterus. Again my alligator urethral forceps came to my rescue, and I removed the foreign body without difficulty. The blunt end renders it as easy to introduce as an ordinary silver catheter. The cut represents the mechanism by which the greater part of the instrument works by a sliding motion, which in turn opens and closes the short alligator blades. With this instrument a firmer hold may be secured, and the fragment is not so liable to break. The compound silver catheters sometimes break off, and may be easily removed by the forceps—whereas the screw would be entirely useless. The greater safety of the former is obvious.

MEDICAL PROGRESS.

THE ACTION OF THE CONSTANT CURRENT IN ELECTRICITY APPLIED TO THE BRAIN ESPECIALLY AS CONCERNS THE EYE.—Dr. Gillet de Grandmont in the *Recueil d' Ophthalmologie* (August) after a series of experimental researches, reviews this subject at considerable length. In using this means for the relief of eye affections, he places the negative pole on the forehead near the termination of one of the branches of the tri-facial nerve, while the positive pole is placed near the superior conical ganglion of the sympathetic. The impression produced shows itself in various ways, viz., by flashes of light which are indicative of an irritation of the optic nerve and a corresponding disturbance of the retinal elements, and serves as a means of diagnosis to determine the degree of alteration of the retina. When, for example, an amaurotic does not see the brilliant blue light, but a feeble yellow glimmer, the prognosis would be far from favorable. The metallic taste in the mouth is not so easy of explanation, it has been supposed to be the result of a dialytic action upon the saliva; the iron taste has been supposed to be due to the decomposition of the elements of the blood containing iron, but this does not explain the copper taste. It is evidently due to a direct excitation of the lingual and glosso-pharyngeal nerves. The buzzing is due to an excitation of the auditory nerve. The dizziness is the result of a congestion of the encephalic centers. This appears only when the current is very strong, or when it is prolonged, and is very objectionable. Another objection to these strong and prolonged currents is the cicatrices which result on the forehead and neck from the destruction of tissue. The lowering of temperature which is shown on the use of a moderately strong current of electricity of short duration seems to negative the foregoing statement of congestion of the encephalic centers by a strong and long continued current; but this is explained by the action on the sympathetic and consequently upon the vaso-motor nerves. A current of feeble intensity stimulates the vaso-motors to the contraction of the vessels and lowers the temperature. A current of greater intensity or long continued exhausts and overcomes the excitability of the vaso-motors, and there is a vascular relaxation which admits of turgid-