the practical work that we do every day, until we are certain of what is shown by the X-ray. It is true that the X-ray is an aid to diagnosis, but while it may be an aid I believe that up to the present time, if admitted to the courts as evidence, it will do more injustice than justice. In a case I have at present, the man was shot a year ago, at close range, with a thirty-two caliber bullet, which entered immediately above the pubes, missing the pubic bone. He did not feel any harm from it for some time. Some weeks ago it appears that an abscess had formed and had broken into the bladder and there was a profuse discharge of pus from it. In sounding the bladder neither myself nor colleagues were enabled to detect the bullet. There was no escape of urine at the time of the primary injury, and it would have been hardly possible for it to have remained in the bladder without causing any irritation. I suspect that this bullet is located somewhere exterior to the bladder, and that sepsis has taken place, followed by suppuration and rupture into the bladder. I am exceedingly anxious to know what the radiographist will find in this case.

From a picture, such as I have seen taken, it is impossible to tell whether there was a separation of the epiphysis from the body of the bone or not, when the facts were that there was no separation whatever. Suppose in a case of this kind a party wishes to defraud us, and should fall off a car and injure the limb slightly (probably the knee is swollen a little), and should go to a radiographist and have a skiagraph taken and exhibit it in court (although we know the man is not injured in the slightest), and there should be the appearance of separation between the epiphysis and bone, there is only one thing that we could expect the jury to say, and that is, we were mistaken and the radiographist was correct. I feel that while X-ray skiagraphs are of value in our practice, as aids to diagnosis, it is necessary for us to develop the facts better before they should be admitted to the courts as evidence, either by the prosecution or the defense.

CLINICAL SIGNIFICANCE OF THE DIFFERENT FORMS OF THE KLEBS-LOEFFLER BACILLUS.

Read before the North Chicago Medical Society Dec. 23, 1897.

BY W. J. CLASS, M.D.

ASSISTANT TO THE CHAIR OF PATHOLOGY, CHICAGO POLICLINIC, MEDICAL INSPECTOR CHICAGO HEALTH DEPARTMENT (ANTITOXIN STAFF).
CHICAGO, ILL.

It is not necessary for me to enter into a detailed description of the bacillus of diphtheria, history of its discovery, growth in different culture media etc.; a few remarks however especially in regard to its morphology will be required in order to present this paper properly. Fraenkel in his work on bacteriology describes the Klebs-Loeffler bacillus as a bacterium, resembling in many respects Fraenkel's diplococcus of pneumonia. According to him, their habitat is in the oldest part of the pseudo-membrane. They are little rods of moderate size, usually slightly bent, about as long as the tubercle bacillus but twice as broad, hence of a rather coarse appearance and usually with rounded ends. But the form of this micro-organism is exceedingly variable, and the differences are striking in appearance. The bacteria are sometimes seen enveloped in a more or less capacious glassy membrane; sometimes the contents separate into several pieces divided by a broad transverse wall; one end of the rod is very frequently thickened like a club; sometimes this change appears on both ends so that there will be dumb-bell shaped structures, usually pointed out as involution forms. The bacilli are immobile, have no spores, perish rather rapidly when dried and succumb at 45 to 50 degrees C. They stain with difficulty, the terminals staining better and therefore standing out as dark spots from the pale center. In recent cases they are usually found associated with streptococci. Martin, in 1892, describes three forms of the Klebs-Loeffler bacillus, as being of comparatively constant occurrence in different cases of diph-

theria, namely, long entangled forms, medium sized bacilli arranged parallel and short parallel bacilli. Other authors have made different classifications, which, however, amount to about the same.

The significance of these different morphologic varieties, and the import of the almost constant occurrence of the streptococcus, especially in severe cases, are the chief points to be considered in this paper. E. A. Peters of London, in an article in the *Journal of Pathology and Bacteriology*, claims the long bacillus is the more constantly present and gives rise to a more severe type of the disease than the shorter form, which is of rare occurrence. He does not seem to attach much importance to the streptococci. His examinations, from which he draws these conclusions, were made from bacteria grown on different culture media instead of from slides made directly from the throat. As the size of the bacilli grown on the different culture media varies considerably, those grown on blood serum being larger than those grown on agar-agar. I do not consider his statistics quite reliable. On the other hand, Cornil and Babes in their text-book describe a long pseudo-bacillus, which occurs in mild cases and is considered, by them, to be an involution form of the true Klebs-Loeffler bacillus. Jacques of Chicago, in an article read before the Chicago Medical Society, makes the following statement; "In the early autumn the bacilli are small, producing a small amount of membrane and involving the cervical glands; little or no temperature and great malignancy causing the patient to be stupid as if under the influence of a strong narcotic. As the winter progressed this type changed. The bacilli became larger, more membrane was formed and the type was pharyngeal, with occasional involvement of the larynx." This statement, in general, agrees with my experience except that I have not noticed any change in the size of the organism as the season advanced, but have found the short bacillus, usually associated with the streptococcus, almost invariably in the severer forms of the disease.

In the cases of which I have kept a record and which I quote, the bacteria I mention were found in the material derived directly from the throat of the patient, a culture being made later in each case to verify the findings.

Case 1.—F. S., aged 3, third day,¹ had extensive membrane in pharynx. No antitoxin was used, death from toxemia fifth day of disease. Short Klebs-Loeffler bacillus and streptococci were found.

Case 2.—C. B., aged 20, fifth day, small amount of membrane in pharynx, larynx slightly affected, severe general intoxication, 5500 units antitoxin used, recovery. Short Klebs-Loeffler bacillus and streptococci were found.

Case 3.—A. M., aged 2, third day, small amount of membrane in pharynx, larynx, trachea and probably also in bronchi. Intubation was done without affording relief, 2000 units antitoxin used, death same day. Short Klebs-Loeffler bacilli and streptococci were found.

Case 4.—G. F., aged 8, second day, membrane in pharynx and tonsils, constitutional symptoms moderate, 1500 units antitoxin used, recovery. Typical Klebs-Loeffler bacilli and staphylococci were found.

Case 5.—E. C., aged 15, second day, small amount of membrane on tonsil and uvula, 2700 units antitoxin used, constitutional symptoms mild, recovery. Strepto- and staphylococci were found.

Case 6.—C. P., aged 9, third day, small amount of membrane in pharynx, constitutional symptoms moderate, 1700 units antitoxin used, recovery. Typical Klebs-Loeffler bacilli were found.

Case 7.—R. P., aged 8, fifth day, extensive membrane on

¹ In these cases the day referred to means the one on which first examination was made. The symptoms are those observed during entire course of the disease.

tonsils and uvula, constitutional symptoms of moderate severity, 2000 units antitoxin used, recovery. Typical Klebs-Loeffler bacilli and streptococci were found.

Case 8.—C. S., aged 3 years, six months, fifth day, involvement of larynx and pharynx, constitutional symptoms severe, 1500 units antitoxin used, recovery. Short Klebs-Loeffler bacilli and streptococci were found.

Case 9.—A. C., aged 3, fifth day, laryngeal type with severe general symptoms, 3000 units antitoxin used. Short Klebs-Loeffler bacilli and streptococci were found.

Case 10.—F. N., aged 8, second day, laryngeal type, pharynx only reddened, constitutional symptoms of moderate severity, 1000 units antitoxin used, recovery. Typical Klebs-Loeffler bacilli and streptococci were found.

Case 11.—H. B., aged 13, second day, small amount of membrane on tonsils, general symptoms mild. Long Klebs-Loeffler bacilli were found.

Case 12.—A. H., aged 6, third day, small amount of membrane in pharynx, general symptoms of moderate severity, 2000 units antitoxin used, recovery. Long Klebs-Loeffler bacilli were found.

Case 13.—K. L., aged 6, second day, small amount of membrane in tonsils, general symptoms mild, 1000 units antitoxin used, recovery. Streptococci and micrococcus tetragenus were found.

Case 14.—J. M., aged 8, second day, extensive membrane in pharynx and tonsils, constitutional symptoms moderate, 2500 units antitoxin used, recovery. Typical Klebs-Loeffler bacilli were found.

Case 15.—L. M., aged 12, second day, slight amount of membrane on tonsils, general symptoms mild, 1500 units antitoxin used, recovery. Long Klebs-Loeffler bacilli were found.

Case 16.—A. N., aged 5, second day, small amount of membrane in pharynx, general symptoms mild, 1500 units antitoxin used, recovery. Long Klebs-Loeffler bacilli were found.

Case 17.—G. L., aged 4, fourth day, extensive membrane in pharynx, constitutional symptoms of moderate severity, 1000 units antitoxin used, recovery. Typical Klebs-Loeffler bacilli were found.

Case 18.—P. L., aged 5, second day, extensive membrane in pharynx, constitutional symptoms of moderate severity, 1000 units antitoxin used, recovery. Typical Klebs-Loeffler bacilli were found.

Case 19.—A. L., aged 1, first day, redness and white spots on tonsils, constitutional symptoms of moderate severity, 500 units antitoxin used, recovery. Typical Klebs-Loeffler bacilli were found.

Case 20.—E. D., aged 12, second day, fairly extensive membrane in pharynx, constitutional symptoms of moderate severity, 1500 units antitoxin used, recovery. Typical Klebs-Loeffler bacilli and streptococci were found.

Case 21.—S. D., aged 8, first day, redness and few white spots in pharynx, general symptoms of moderate severity, 750 units antitoxin used, recovery. Typical Klebs-Loeffler bacilli and streptococci were found.

Case 22.—G. P., aged 8, fifth day, extensive membrane in pharynx, general symptoms severe. 2000 units antitoxin used, recovery. Short Klebs-Loeffler bacilli and streptococci were found.

Case 23.—H. B., aged 4, first day, redness of tonsils and pharynx, general symptoms mild, 1000 units antitoxin used, recovery. Long Klebs-Loeffler bacilli were found.

Case 24.—M. K., aged 17, second day, moderate amount of membrane on tonsils, constitutional symptoms moderate in severity, 1500 units antitoxin used, recovery. Long Klebs-Loeffler bacilli were found.

Case 25.—R. L., aged 5, third day, moderate amount of membrane in pharynx, constitutional symptoms severe, 2000 units antitoxin used, recovery. Short Klebs-Loeffler bacilli and streptococci were found.

Case 26.—M. R., aged 4, second day, small white patches on tonsils and uvula, general symptoms mild, 500 units antitoxin used, recovery. Pseudo Klebs-Loeffler bacilli were found.

Case 27.—G. C., aged 10, third day, extensive membrane in pharynx, constitutional symptoms of moderate severity, 1500 units antitoxin used, recovery. Typical Klebs-Loeffler bacilli and streptococci were found.

To sum up: We have seven cases with severe constitutional symptoms, with two deaths; in all of these the short Klebs-Loeffler bacillus and streptococci were found; thirteen cases in which the constitutional symptoms were not so marked, but the membrane was, as a rule, thicker and more extensive; in six of these typical Klebs-Loeffler bacilli were found alone, in five

they were associated with streptococci, and in one each are found typical Klebs-Loeffler bacilli with staphylococci and long Klebs-Loeffler bacilli, respectively. In the remaining seven cases the constitutional symptoms were mild or entirely absent; of these, four showed the presence of the long Klebs-Loeffler bacillus alone, and, in one each, streptococci and staphylococci; streptococci and micrococci tetrageni, and pseudo Klebs-Loeffler bacilli were found.

As the result of these examinations I have arrived at the following conclusions:

1. That the short Klebs-Loeffler bacillus apparently produces a toxin of greater virulency than the longer forms, although the local manifestations may not be so extensive.

2. That the long Klebs-Loeffler bacillus and the streptococci when found alone give rise to a mild type of the disease.

3. That the streptococcus is found associated with the Klebs-Loeffler bacillus in most of the severe cases. Its special significance is not so clear, but it is possible that by causing a more intense inflammatory reaction it opens avenues by which the toxins of the Klebs-Loeffler bacillus may find more ready entrance into the circulation, plus its own toxin.

4. That the apparent beneficial action of the antitoxin of the Klebs-Loeffler bacillus in cases where this bacillus is not present, may be due to the fact that though the local action of the different microbes varies to a considerable extent, the action of their toxins, as is shown by the similarity of the constitutional symptoms produced by them, present many kindred features. The thought, therefore, arises that the antitoxin of one infection may have an inhibitory effect on the toxin of another, as is shown by the fact that whooping cough and some other infectious diseases have been shown to occur less frequently in vaccinated persons and some cases have, apparently, been cured by vaccination.

NOTE.—By the term typical Klebs-Loeffler bacillus is meant the medium sized bacillus as described by Martin.

REFERENCES.

- Annual of the Universal Medical Sciences, 1896, Vol. 1 and 2.
General Jahrbuch fuer Kinderhik, 1894, Bd. 38, p. 254.
Karl Guenther: Bacteriologie, 1895, p. 298.
Cornil-Babes: Les Bacteries, 1896, Vol. 2.
Centralblatt fuer Bacteriologie und Parasitenkunde, 1896-1897.
E. A. Peters: Journal of Pathology and Bacteriology, 1896, Vol. 4, No. 2, p. 181.
Fraenkel: Bacteriologie, 1892.
W. K. Jaques: Chicago Medical Recorder, 1896, Vol. 11.
1301 Belmont Avenue.

TETANUS; ANTI-TETANIC SERUM; REPORT OF CASE.

Read before the Winona County Medical Society, July 6, 1897.

BY HUGH F. MCGAUGHEY, B.S., M.D.

WINONA, MINN.

Tetanus is a toxic infection dependent on the products of a specific germ. It is spoken of as traumatic when the infection takes place through a recognized wound. In idiopathic tetanus the infection atrium is some unnoted lesion. It is possible that the germs may sometimes find entrance through mucous membranes. Infection after labor gives rise to puerperal tetanus. Trismus neonatorum occurs as a rule through inoculation of the umbilicus. Tetanus cephalicus sometimes results from wounds of the head, the symptoms being limited to this region.

The bacillus of tetanus was first described by Nicolaier, in 1884. He obtained it from wounds and from the soil. Five years later Kitasato secured it in pure