improvement in the paralysis of the right external rectus, while there is none in the paralysis of the left.

The sinuses have been irrigated daily with a boric acid solution, resulting in a decided diminution in the quantity of purulent secretion. He was discharged from the hospital on May 2, greatly improved. The opening in the left sinus shows a tendency to close, and in consequence the occipital and vertex pains have returned, but they are not so intense. This cavity will probably have to be opened again. This case is especially interesting from the fact that the other sinuses are not involved, and the pressure, instead of being made on the optic nerve, as is usually the case, is directed upward and outward and involves the abducens nerves just before passing through the sphenoidal fissure.

DISEASES OF THE MAXILLARY SINUS IN RELATION TO THE DISEASES OF THE EYE.

The maxillary sinus is subject to the same diseased conditions that are met with in the other accessory cavities, and probably is the most frequently affected, as it has both the teeth and the nose from which pathologic processes may extend. It probably is, however, less frequently complicated with diseases of the eye than the other accessory sinuses.

Many of the eye symptoms that are met with in abscess of the maxillary sinus are reflex in character, such as narrowing of the field of vision, photophobia, dilatation of the pupil, blepharospasm, and ptosis; while others are the result of an inflammatory extension; such as osteoperiostitis, abscess of the orbit, dacryocystitis, and iridochoroiditis.

It will be observed that there are many symptoms of diseases of the eye, when associated with accessory sinus inflammations, which are common to one or more of the affected cavities. There are, however, certain symptoms which are peculiar to certain sinuses, and it is only by the closest observation of these cases, both from the rhinologic and ophthalmologic side, that we can hope to make an early diagnosis. While pain is a symptom met with in these affections, its location will frequently aid one in determining the cavity which is the seat of the inflammation. This, however, must not be depended on alone, for occasionally in ethmoidal abscess the pain is referred to the supraorbital ridge as well as to the eye and infraorbital region.

Orbital abscess may occur as readily from an extension of an inflammation of one cavity as from another. I believe, however, basing my opinion on my own experience, it more frequently results from frontal sinus disease. A narrowing of the field of vision for white and for colors may follow an abscess in either the frontal, ethmoidal, sphenoidal, or maxillary cavities. It is, however, more frequently met with when the frontal and ethmoidal sinuses are jointly diseased. Refractive disturbances are more frequently noticed in inflammations of the frontal and ethmoidal cavities, and generally in those cases where the pressure is exerted either at or posterior to the equator of the eyeball.

Exophthalmus may be present in any well-advanced abscess of the accessory sinuses; the extent and direction in which the eye is pressed from its normal plane indicates the cavity which is affected; e. g., in frontal disease the eye is pushed forward and downward, in ethmoidal, forward and outward, in sphenoidal, directly forward and slightly outward, while in maxillary disease it is directed upward and outward.

Sudden blindness, together with ptosis and strabismus, when associated with abundant discharge into the post-nasal space, is indicative of sphenoidal disease.

EYE TROUBLES ATTRIBUTABLE TO NASOPHARYNGEAL AND AURAL DISTURBANCES.*

BY JOSEPH A. WHITE, M.D.

Professor of Ophthalmology, and Associate Professor of Otology and Laryngology, University College of Medicine.
RICHMOND, VA.

The direct anatomic connection between the eye and nose makes it easy to understand how disease can spread from the latter to the former by way of the nasal duct. Moreover, the intimate nervous and vascular connection between the two organs also plays an important part in the propagation of disease from the nose to the eye. Both these modes of transmission have been recognized since the last century, and referred to by different authors, but Bresgen—1881—was the first to lay any special stress on the fact that conjunctival catarrh was often dependent on nasal catarrh. Since his article appeared, medical literature has had numerous contributions on this and other eye troubles of nasal origin, whether by direct or reflex communication. Appended to my article on "Neurosis of the Nose and Nasopharynx," in Burnett's "System of Diseases of the Ear, Throat and Nose," and to Dr. Gould's article in the same treatise, on "Disease of the Eye Dependent on Diseases of the Nose," can be found an extensive bibliography of the subject from the earliest times to 1892. Since then, other contributions on the subject have appeared from time to time, adding to the accumulation of testimony to the importance of considering the nose as an etiologic factor in eye disease.

From these references it will be seen that, according to a large number of observers, we have not only troubles of the lachrymal apparatus by direct transmission of pathogenic organisms from the nose to the eye, but a large number of other eye diseases traceable to direct or reflex extension from nasal and nasopharyngeal causes, such as asthenopia, photophobia, retinal hyperesthesia, *muscae volitantes*, ocular neuralgia, contraction of the visual field, blepharospasm, phlyctenular ophthalmia, conjunctivitis, keratitis, mydriasis—Goris and Trouseau—amaurosis, amblyopia, strabismus and even glaucoma, myopia—congestion of uveal tract (Landolt)—and astigmatism (spasmodic).

The supposed nasal and nasopharyngeal causes were as numerous as the resulting eye troubles, among them being acute rhinitis, chronic hypertrophic rhinitis, synechia, nasal polyps, growths, diseases of the accessory sinuses, adenoid tissue, enlarged tonsils, simple atrophic rhinitis, *ozena* and nasal hydrorrhea.

Coryza gives rise to lachrymation, conjunctival injection, dacryocystitis and ocular pains. Chronic rhinitis produces rebellious conjunctivitis and blepharitis, phlyctenular conjunctivitis, epiphora—asthenopia—myopia (?)—congestion of uveal tract (Landolt)—and astigmatism (spasmodic). *Ozena* produces ulcers of the cornea, dacryocystitis, neuritis, iritis, chorioretinitis, amblyopia, amaurosis and orbital phlegmon. Purulent rhinitis causes iritis and uveitis, phlebitis of ophthalmic veins, and contraction of the visual field. Sinusitis—frontal, maxillary, ethmoidal, etc.—produces orbital neuralgia, iritis, iridochoroiditis, orbital cellulitis, neuritis, exophthalmus, diplopia, etc. To nasal hydrorrhea is due neuritis and atrophy of the optic nerve. Nasal synechia cause blepharospasm, conjunctivitis, amblyopia, and strabismus, and nasal diphtheria and croup, similar af-

*Presented to the Section on Ophthalmology, at the Fiftieth Annual Meeting of the American Medical Association, held at Columbus, Ohio, June 6-9, 1899.

fections of the conjunctiva. Adenoids cause phlyctenular keratitis and conjunctivitis, blepharitis and strabismus, etc., while pharyngeal diphtheria gives rise to paralysis of accommodation, and tonsillar affection to lachrymation, blepharospasm, conjunctivitis and keratitis, and orbital thrombosis.

Eliminating the accessory sinuses, and the ordinary causes of nasal stenosis, which are treated of in other papers, I will briefly refer to the eye troubles, known or supposed to be etiologically connected with nasal catarrh without nasal obstruction, with hydrorrhea, with atrophic rhinitis and ozena, and with lymphoid tissue in the nasopharynx, such as adenoids and enlarged tonsils.

It is almost a waste of time to speak of the well-established connection between affections of the lachrymal apparatus and nasal disease, probably 80 per cent. of the former being caused by the latter, either by the direct extension of germs from the nose to the lachrymal sac, or by the obstruction of the nasal orifice of the nasal duct, thus preventing the out-flow of the lachrymal fluid, with its consequent disintegration and the injurious effects therefrom.

Although epiphora can be the result of any lid troubles causing malposition of the punctum, or of that rarer cause, closure of the canaliculi, I very seriously doubt whether the usual departure of troubles of the lachrymal sac, viz., obstruction of the nasal duct, is often due to other causes than nasal disease. I think most writers and observers are of accord on this point, although all do not lay equal stress on the necessity of the nasal treatment, if we would cure the lachrymal trouble.

It is interesting to note some of the results of slight operations on the nose and throat as indicating another line of connection with the eye. In 1892 Ziem reported having observed hyperemia of the disc with venous pulsation, following the application of the galvanic cautery to the nasal mucosa, which he explained by the anastomosis between the vessels of the nasal fossæ and the orbit, probably by way of the ethmoidal vessels—an interesting coincidence if not a proven sequence. Akin to this is the report of amaurosis—optic neuritis—following on a traumatic hemorrhage from the nose, by Schmidt-Rimpler, which he supposed due to resulting ischemia of the retina. Somewhat similar was Blok's case—reported to the Ophthalmological Society of the Netherlands, in 1894—of marked asthenopia in emmetropia with normal convergence, cured by cauterization of the inferior turbinate, evidently of reflex influence. More curious still is the case reported by P. Viollett,¹ of permanent strabismus following the cauterization of the nasal mucosa, and one more difficult of logical explanation. He supposed it to be due to nervous shock in an individual predisposed to strabismus by a pre-existing astigmatism and a general nervous state. This case recalls that of Goureau² who, in 1891, reported the cure of strabismus in a child, by the removal of adenoids, in which instance the shock worked in the inverse order to the first case. Laurens³ reported another case of strabismus cured by removal of adenoids. In May, 1894, in a paper entitled "Nasopharyngeal Therapy in Ocular Affections," presented to the French Ophthalmological Society, Guibert, among other cases, recorded one of recurring keratitis cured by removal of adenoids after local and constitutional treatment had signally failed. Ziem has reported a similar case.

I have often seen pain in the eye produced by the operation for adenoids, which is in the same line of reflex effects. Other cases of like character have been recorded, and they seem to point to a physiologic and pathologic

relationship between the nose and eye along lines not yet thoroughly explored, and in consequence the explanation of the production of such effects is not always satisfactory.

These results of operations on adenoids are probably open to criticism, and especially the supposed cure of keratitis by this means. We are all familiar with ulcerative keratitis as an accompaniment of lachrymal disease, which may have been caused by direct infection from the nose, but it is difficult to account for a keratitis secondary to nasal disease without accompanying affection of the lachrymal apparatus, even in ozena—Trousseau, Nieden and Massini—as its so-called characteristic microbe, the bacillus of Loewenberg, Fraenkel and de Vedova has been found by Terson and others in the lachrymal fluid, without any alteration of the conjunctiva or cornea. Still, Fortunati⁴, in February, 1896, reports two cases of ulceration of the cornea in conjunction with nasal trouble, in which the ocular treatment was of no avail, and which were only benefited by the improvement of the nasal condition. In these cases the lachrymal passages were in a normal condition, and moreover the nasal trouble was not of microbial origin. He explained it as a neuroparalytic trouble, the cornea being somewhat anesthetic without phagedenic infiltration, without hypopyon and almost without pain, the trophic disturbance starting probably in the peripheral filaments of the second branch of the fifth pair in the nose, extending to the Gasserian ganglion and thence to the ciliary ganglion and ciliary nerves.

Dr. Haight⁵, in the discussion of his paper on "Diseases of the Eye, Caused by Diseases of the Nose," reports a case of ulceration of the cornea resistant to all treatment, until the accompanying adenoids and enlarged tonsils were removed.

I do not think there is any question as to the apparent influence manifested by the presence of lymphoid tissue in the post-pharyngeal space and fauces on the progress of so-called phlyctenular keratitis or keratoconjunctivitis, which often develops into or is accompanied by deep ulceration of the cornea. Still, the explanations offered differ, some claiming it to be the effect of a generous scrofulous dyscrasia, and others that the condition of the eye is reflexly produced by the presence of the throat trouble, proven in many instances by the rapidity with which the eyes get well when the adenoids and enlarged tonsils are removed, an argument met by the rejoinder that the removal of the latter, with the improvement to breathing, brings about an improvement in the general health and reacts favorably on the eyes.

That iritis and uveitis have been supposed to be etiologically connected with nasal troubles is well known since Ziem first reported a case of iritis which he claimed to be caused by purulent rhinitis.

In May, 1895, before the French Ophthalmological Society, Fage reported a case of iritis associated with ozena which he supposed to be due to the latter by infection from the coccobacillus of Loewenberg, but as in Ziem's case the demonstration is not conclusive. E. Berger also observed a case associated with purulent rhinitis.

W. C. Posey⁶ reports a case of *metastatic uveitis* with loss of sight, consecutive to a purulent rhinitis. In the discussion, Randall said he had seen a similar case. There was no symptom of trouble with accessory sinuses. Panas⁷ supports this contention with the explanation that the eye trouble is due either to the passage of microbes by way of the vessels and lymphatics from the nose, an endo-infectious process, or to toxin poisoning.

So also may be explained E. Berger's case of central scotoma in a woman with ozena, in whom the nasal treatment ameliorated the visual defect.

Sulzer's⁸ two cases of optic neuritis with scotoma, in connection with ozena, cured by treatment of the nasal affection, are in the same line, due either to the propagation of an infectious process by way of the vessels and lymphatics, or to toxin action, as in post-diphtheric paralysis or in tetanus. In like manner might be explained cases of amblyopia and amaurosis in ozena by an infectious process traversing the ethmoid plate, and causing subacute basilar meningitis, with secondary involvement of the optic nerve.

Panas⁹ reports a case of double orbital thrombosis due to gangrene of the tonsil—infection by way of the anastomotic veins pointed out by Trolard, connecting the tonsil with the cavernous sinus through the foramen ovale.

Asthenopia, lachrymation and contraction of the visual fields were long ago reported by Hamilton¹⁰ in cases of atrophic rhinitis and ozena, and since observed by others—these conditions being improved by the nasal treatment. I have seen cases of asthenopia and pain in the eye and ear in connection with atrophic rhinitis without appreciable lesion of the eye or ear which appeared to be due to the nasal trouble.

The connection of optic neuritis and nerve atrophy with nasal hydrorrhea or serous rhinorrhea is a subject of interest, owing to the obscurity of the etiology. Bosworth collected 18 cases, of which 5 had optic nerve atrophy. These were reported by Baxter, Leber and Nettleship, each 1, and 2 by Priestly Smith. Professor Körner¹¹ has added four more to these, 1 each by Gutsche, MacKenzie Wallace, Wood and Hardy, and 1 by himself. He overlooked 1 reported by A. Emrys-Jones, making 10 in all, of which 4 died, 1 recovered and the result in the others is unknown. Dr. Gould reports 2 cases, 1 by Elliotson and 1 by Paget, supposed to be of the same character, but, on examination, I find no mention of the condition of the optic nerve in either case. An autopsy was performed in 2 of them—in Baxter and Gutsche's; in the former the result was negative, in the latter, death was found to be due to meningitis, but a tumor of the hypophysis was found which had extended into the sphenoidal sinus. The optic nerves, especially the right, were found compressed by the tumor and there was probable neuritis. Körner thinks the discovery of a growth in this locality is a step toward the elucidation of these enigmatic cases.

Since Bosworth made his collection of cases of nasal hydrorrhea, in 1889, many such have been put on record—the list is a long one, but these are the only ones with optic nerve complications, as far as I have been able to discover. It is possible more of them developed optic neuritis, and recovered, but, in the absence of the ophthalmoscopic examination, this is a mere surmise. In all probability, both the hydrorrhea and the neuritis are from the same cause, instead of the latter being caused by the former.

I have treated two such cases in the last year, one a typical one of two years' duration, with constant discharge of a clear limpid fluid, and a resulting chronic eczema of the upper lip, and the other with intermittent attacks. While both have shown minor eye troubles, such as asthenopia, with congestion of the conjunctiva and blepharitis, neither has had any optic neuritis. In neither have I been able to discover the cause of the discharge.

From this résumé of the relations between naso-

pharyngeal and eye troubles, it will be observed that disease is transmitted from the nasopharynx to the eye by more than one way: it may be by direct microbial contamination along the lachrymal passages; by mechanical obstruction of the nasal duct at its lower end or in its canal, with resulting pathologic changes; by propagation through the blood-vessels and lymphatics; by toxin poisoning; or it may be by reflex nervous influences. It is in any or all of these ways that pathologic alterations of the mucosa of the nose and nasopharynx can bring about secondary changes in the organ of vision and its adnexa.

Enough, however, has been brought forward to show at least the importance of examining the nose and nasopharynx when dealing with eye troubles of doubtful causation or intractable to local treatment. It has often been my experience to find nasal trouble present in cases of intractable asthenopia, and also of recurring conjunctivitis, which were benefited by the cure of the nasal complication, but I have never seen a case when I was satisfied this was the sole cause of the ocular trouble. For example, I have seen reflex trouble, such as blepharospasm, temporarily benefited by nasal treatment, which recurred later, and was found to have an ocular cause underlying it. This, however, does not apply to disease of the lachrymal apparatus in which, as previously stated, I feel assured that the nasal treatment is of as much avail, and often as necessary, as the treatment of the eye, to prevent frequent relapses.

EYE TROUBLES SUPPOSED TO BE DUE TO AURAL AFFECTIONS.

The connection between the eye and ear is one long known and often discussed. Medical literature is filled with references to this relationship in its anatomic, physiologic and pathologic aspects.

Anatomically they are connected, 1, by way of the brain envelopes and the proximity of their nerve nuclei; 2, by the anastomosis of the superior cervical plexus and the auriculotemporal with the facial at the mastoid apex, and by the branches of the tympanic plexus, which innervate the lining of the drum and the mastoid cells; and 3, by the circulation.

Physiologically they are connected by the functional action of the trigeminus; by the supposed physiologic influence of the semicircular canals; and by the peculiar phenomena observed in the reaction of sight and hearing on one another, in the conjoined action of one sense on stimulation of the other.

It has been shown by Duval, Laborde and others, that lesions of the trigeminus can bring about trophic disturbances in the eye at the same time—keratitis—and in the ear, and that certain pathologic conditions of the ear can by extension to the trigeminus at the level of the Gasserian ganglion give rise to ocular lesions, chiefly of a reflex character. The investigations of Cyon, Baginsky, Hogenes, etc., show that lesions of the semicircular canals can produce various ocular disturbances.

As for the peculiar phenomena referred to, such as light and color sensations caused by perception of sound, they have so far not been satisfactorily explained; but we know that in some people, especially those of high-strung nervous organization, or so-called neurasthenics, the perception of a certain sound or tone causes at the same time the perception of a particular color—the eye functions being undisturbed—which, I suppose, under other conditions could be reversed, and although it probably has its origin in the nerve-centers, and not in peripheral excitation of these special organs, it shows an intimate connection between them, the nature of which has yet to be elucidated. The physiologic relationship

is also shown by d'Arsonval's experience in becoming deaf for one to one and a half hours after steadily gazing at an arc light for some minutes. Waldeyer's morphologic parallel between these two organs is familiar to you all.

Retina analogue of organ of Corti.

Choroid analogue of connective tissue wall of cochlear duct.

Sclerotic analogue of bony labyrinth capsule.

Conjunct. sac analogue of tympanum.

Cornea analogue of membrana tympani.

Nasal duct analogue of Eustachian tube.

Orbit analogue of bony auditory meatus.

From these anatomic and physiologic relations we can see that it is not unnatural to look for some pathologic connection between troubles of the eye and ear.

Blepharospasm apparently results from cerumen impaction, foreign bodies in the external ear and syringing the ear—Rampoldi, Ziem, Buzzard, etc.; also from the presence of pus or polyps in the drum—Deleau, Duterné, Gottstein.

Blurred and double vision have also been reported as having resulted from examinations, traumatisms and diseases of the external ear.

Lachrymation, ciliary neuralgia and muscular disturbances have appeared as a sequence of traumatisms, causing lesions of the drum.

Amblyopia and contracted field for both light and color have seemingly been caused by obstruction of the Eustachian tube, and improved by catheterization—Kisselbach and Wolfberg. Contraction of the field was also observed by Lucae, from the use of the air douche in the external canal in perforation of the drum, probably due to irritation of the tympanic plexus.

Gervais reported strabismus with myosis from mastoid inflammation, which disappeared when the abscess was opened.

Troubles of the accommodation have been observed from irritation of the branches of the trigeminus in the organ of hearing, and cured by the improvement of the aural condition—Poltzer, Wolfberg, Urbantschitsch.

Nystagmus is probably one of the most commonly reported eye reflexes from the ear. Baginski's experiments on rabbits in this line are well known, and show the muscular disturbance to arise from water thrown into the ear, and from disease of the labyrinth and semicircular canals—Jansen, Cohn, Kipp, Urbantschitsch.

Nystagmus should be watched for carefully, as it is sometimes so slight that it is not perceived by the patient. It is more readily noticeable when it occurs from syringing the ear in suppuration of the drum, and is nearly always from labyrinthine irritation—occasionally it accompanies meningitis of aural origin, or thrombosis about the petrous bone. Geronzi—October, 1897—reported a case from an acute non-suppurative middle ear inflammation, following influenza. There was also diplopia in outer parts of the entire field. Singer¹² reported a case following a Stacke operation, whenever pressure was made on the drum, or the air in the external meatus was rarified. The pupils dilated at the same time. Urbantschitsch also reported a case of aural polyps in which nystagmus was produced each time the polyp was pressed on. Phluger¹³ had several years previously reported a similar case. Gruber has also seen such cases.

Abducens paralysis has been observed in one or two cases of intracranial involvement, and recently—German Otological Society, May, 1898—Habermann reported a case of middle-ear suppuration cured in ten days, but

which relapsed in three weeks, with headache and sudden paralysis of the abducens. The dura was exposed and found hyperemic only. The paralysis, which was permanent, was supposed to be due to extension of the inflammation into the petrous and sphenoid bones. In the discussion that followed Katz stated he had seen the same thing due to meningitis caused by ear trouble. Brüeger had also seen it in what was supposed to be latent meningitis of aural origin. Mann had seen it in perisinous (?) abscess. Jansen thought it due to serous meningitis, dependent on sinus thrombosis or epidural abscess. Habermann, however, concluded that sinus trouble could be excluded, as he had seen abducens paralysis on the opposite side from the ear trouble.

Paralysis of the orbicularis has been observed as resulting from foreign bodies in the external ear—non-perforating catarrh of the drum, purulent otitis media, etc.

Hippus—alternating contraction and dilatation of the pupils—has been observed by Pisenti¹⁴ in a person with chronic middle ear catarrh, each time the Eustachian catheter was used, and each time air was driven into the drum. Reflex might be by three ways:

1. Irritation of the trigeminal branches in the floor of the nose, transmitted to the sensory root of the ophthalmic ganglion. 2. Irritation of the posterior and inferior ethmoidal nerves and transmitted by the ophthalmic ganglion to filaments of the oculomotor. 3. Irritation of the internal ear by dilatation of the drum and transmitted from the ampulla to Deiter's nucleus and thence to the nuclei of the sixth and third pairs.

Purulent choroido-iritis from purulent otitis is reported by Kipp and Pomeroy, either of metastatic origin from emboli, or possibly by extension to brain envelopes and along optic nerve sheaths to the eye. Panas¹⁵ ("L'auto-infection dans les maladies oculaire") reports a similar case in the service of Dr. Quénu, except that the metastatic panophthalmitis was on the opposite side from the diseased ear, the eye on the same side remaining intact.

Orbital cellulitis, with secondary neuritis and amblyopia, has also been observed by the same author, as the result of purulent otitis media, which was cured by injection of antistreptococcus serum when ordinary measures had failed.

Obscuration of visual field and keratitis have been observed by Knapp—mydriasis and hemiopia by Moos, in Menière's disease; whether as a sequence or coincidence is doubtful.

Schwartz has reported a case of progressive amblyopia in a subject of an acute affection of the acoustic nerve, which was arrested and cured by leeching.

Statistics of deaf mutes show a larger number of cases of amblyopia—G. Mayer, Schermer, Gellé, Schmolz and others.

Wyatt Wingrave—October, 1897—reported a case of tuberculosis of the ear with optic neuritis. There was a discharge from the ear, but the general health was perfect. He thought it might be due to latent meningitis, or a tuberculous gumma. The good health, however, was a contraindication of meningitis, and the age, only 25, of gumma.

Optic neuritis is also frequently found in connection with mastoid abscess, suppuration of the drum, and their intracranial complications. Exophthalmus and strabismus are also found in the same connection—Zaufal, Kipp, Schwartze and others. Paralysis of the oculomotor nerves and homonymous hemianopsia have also been observed from the same causation.

These ocular manifestations, resulting from suppur-

ative otitis, are of especial diagnostic significance in determining the necessity of operative intervention. While optic neuritis is not a constant symptom of meningitis, or cerebral abscess, it usually indicates some participation of the brain or its envelopes with the ear trouble. It, like all other so-called diagnostic signs, might be misleading under unusual conditions. I have seen a case of middle ear suppuration, intractable to treatment, cured by opening the mastoid, consent of the patient being obtained only by my assurance of probable intracranial complications, because of the advent of optic neuritis. The operation showed there was no intracranial trouble due to the ear disease, the neuritis being only a coincidence, and not a sequence. This, however, does not materially detract from the value of neuritis as a symptom of brain involvement. When exophthalmus and swelling of the eyelid accompany the neuritis, we can be reasonably sure of thrombosis.

When we analyze the recorded cases of ocular phenomena in aural troubles, we find that there is occasionally some uncertainty about the connection of the two. It is easy to understand why photophobia, eye pain, etc., can be a sequence of ear trouble, because of the intimate nervous connection by the trigeminus branches, but it is hard to see how amblyopia can result from uncomplicated inflammation of the middle ear, even in deaf-mutism, although in such cases the visual function may be prevented from attaining its complete development, just as there may be arrest of development of the cerebral centers. Badal also calls attention to the frequency of serious trouble of the fundus oculi in these cases. In regard to nystagmus, I can not see how it may be physiologically explained by the reflex irritation, without some complication of the internal ear.

The question of ocular troubles in vertigo and Ménière's disease is involved in obscurity, although, according to Mendel (Berlin), vertigo is a collection of symptoms determined by an alteration of the sense of equilibrium, combined with defective function of the eye muscles. Probably, however, the eye examination may help to make a diagnosis between Ménière's disease and a cerebellar lesion.

Still, while much is yet to be explained in regard to pathologic influence of the ear on the eye, we may conclude from what we have before us that the influence really exists, and is exerted by both direct and indirect propagation, by way of the nerve tract and by the blood current.

BIBLIOGRAPHY.

1. *Annales d'oculistique*, June, 1897.
2. *Recueil d'ophtalmologie*.
3. *Med. Presse*, Feb., 1896.
4. *Archiv. Ital. di Otol., Rhin., etc.*
5. *JOURNAL*, May 21, 1898.
6. *Oph. Record*, Jan., 1898.
7. *L'auto-infection dans les Maladies Oculaires*.
8. *Annales d'oculistique*, Jan., 1895.
9. *Loc. cit.*
10. *Trans. Intercollegiate Congress of Australasia*, 1889.
11. *Archives of Otolaryngology*, Oct., 1898.
12. *Munch. Med. Woch.*, June, 1898.
13. *Berliner Klin. Woch.*, 1891.
14. *Bul. de l'Acad. de Med., Perugia*, 1897.
15. *Archives d'Ophtal.*, May, 1897.

DISCUSSION.

DR. HOLMES—I can only make a very few remarks on this subject, because of lack of time. Dr. Bull deserves the thanks of the ASSOCIATION for such a terse and instructive paper. It is my experience that the frontal sinus is not so often attacked as we assume. I had opened the sinus a number of times—a matter of exploration where there was every evidence of disease, but found that it was limited to the ethmoidal or sphenoidal cells, and at times to the antrum of Highmore. I agree with the Doctor that the lamp test for the frontal sinus is of little or no value. The danger of treating the frontal sinus through the nose is such that I think it should not be under-

taken except in the rarest cases. I attempted it a number of years ago, but after dissecting a great many specimens, I demonstrated how impossible it is to do this without danger, especially when the parts are softened by disease. The liability of invading the cerebral cavity is greater and quite a number of deaths have resulted. If I wish to know anything about the sinus, I prefer to make an exploratory opening, which can be made small, and at the end of the week nothing is left but a small white line, and in the course of a month or two nothing at all to show that any operation has been done. If you find disease you can easily enlarge the opening.

With regard to the maxillary antrum, I do not believe that the teeth play a very important part in disease of the antrum of Highmore; I believe it comes from the nose in the vast majority of cases, and I base that opinion on the extensive examination of cadavers. It is true that once in a while it may occur from the teeth, for bad dentistry, fracture of the process, etc., may cause it. I have seen quite a number of antrum affections with little or no complications relating to the eye, and it is my limited experience that we do not get eye symptoms so often in diseases of the antrum of Highmore as we do in diseases of the frontal, ethmoidal or sphenoidal sinuses. The illumination test for the antrum of Highmore is very important, but I should not place any value on any test excepting that which gives the red reflex from the pupil; I am not speaking of the new test, the retromaxillary illumination, but of the ordinary tests, and with those I feel safe in assuming the presence of infection if I do not get a pupil reflex. I have often injected water into one sinus and then made the test, getting the pupil dark; then evacuating the sinus I would again get the pupillary reflex, showing that even a small quantity of fluid can be detected.

Now I wish to say one word about iodoform. I have not used it for five years, for it is an abominable drug. I am satisfied that we get quite as good results without that stinking drug anywhere about our patients and I do not see how we can justify ourselves in continuing to use it, especially as it has been associated with diseases that patients would not like to be charged with.

As to the majority of cases being cured in six weeks, I am sorry to say that mine are not. I have some that have been going on for five years, with large openings through the canine fossa, and there are cases no one could cure. Of course the acute cases may be cured quickly.

SOME OF THE WAYS IN WHICH X-RAYS ASSIST IN MEDICAL DIAGNOSIS.*

BY FRANCIS H. WILLIAMS, M.D.

BOSTON.

For satisfactory medical work with the X-rays, the first requisite is an efficient apparatus, the second, experience in examining a large number of cases.

It is impracticable in so short a paper to describe and compare the advantages and disadvantages of the various forms of static machines and coils that have been used to excite vacuum tubes, but I present photographs of the static machine I have been using at the Boston City Hospital for some time past, which give in themselves a sufficient idea of the machine and of my method of examining patients. The patients are brought to the X-ray room on a stretcher, that fits wooden horses placed in front of the machine, so that they are spared all possible fatigue. This machine has four revolving plates, six feet in diameter, and four fixed plates, six feet four inches in diameter. In Europe they are using coils more than static machines.

The vacuum tube is the part of the X-ray outfit that demands the most care and study by the physician. Those designed by Dr. William H. Rollins of Boston I have found excellent for my purpose. They are provided with his regenerators for lowering the vacuum, and the anode in one form is so constructed as to be readily rotated if the energy used is sufficient to melt a hole through the platinum; in another form the anode is hollow and can

*Presented to the Section on Practice of Medicine, at the Fiftieth Annual Meeting of the American Medical Association, held at Columbus, Ohio, June 6-9, 1899.