scopic study showed that the floor of the mouth was infiltrated but the glands were free.

As the removal of the lower jaw, especially in the region of the symphysis is mutilating, the author has attempted to accomplish the same results in a different way.

The glands of the neck are first removed and, after the operation, their connection with the floor of the mouth below the lesion is thoroughly burned with the cauter, and the wound closed. Then the lesion in the tongue or floor of the mouth is attacked with the electric cauter. The application of this is usually repeated two or more times, until everything is destroyed down to the area of the first cauterization from below. The healed skin flap of the first operation forms the floor of the mouth and prevents an oral fistula.

The first operation after this method was performed in April, 1912,—two years ago. The lesion was a cancer occupying the floor of the mouth between the tongue and the symphysis of the jaw. It was about the size of a silver dollar. Permanent cures have been accomplished in similar cases by an *en-bloc* dissection of tongue, floor of the mouth, jaw and glands. The oldest case lived fifteen years, but this is a very mutilating operation, and the recent patient refused to submit to it. This led the author to attempt what he had had in mind for some years. At the present writing, two years since operation—there is no evidence of recurrence and no mutilation.

Since then four cases have received this treatment with, so far, apparent success.

The majority of cases of cancer of the tongue seek surgical aid at an unnecessarily late period. In every case the patient is warned. There is always something to be seen and felt in the tongue or floor of the mouth. If such a lesion is investigated at once, a local operation with the electric cauter should be sufficient; in a little later stage removal of the glands and repeated cauterizations in the mouth; in still later stages resection of the jaw must be done. The author's recent experience seems to show that this operation should be done in stages. First, thorough removal of the glands with cauterization of the floor of the mouth from the neck wound; second, cauterization of the lesion within the mouth; third, removal of the lower jaw and cauterized area. These points will be discussed in detail with illustrative cases in the complete paper.

When the cases observed until 1908,—a period of 18 years,—are compared with those observed during the past five years, the influence of education is well shown. The very early pre-cancerous lesions have increased from 8 to 30%. The late and inoperable cases have decreased from 18 to 10%. The cures have increased from 21 to 50%.

When the author considers the cases personally operated on by him by these newer methods in the past five years (14 in all) he finds that

there has been no postoperative mortality, and so far but one patient is dead of recurrent carcinoma. In this case the lesion of the tongue had previously been subjected to operation, the recurrent tumor was extensive, and the glands of the neck involved.

In this group every type of operation according to the newer methods just described is represented. At the present time there is evidence of recurrence in only one case; here the lesion was most extensive and the operation most radical.

The experience with these 14 cases proves the point as far as the immediate mortality is concerned, because considering all cases studied, the postoperative mortality has been about 22%. Since recurrences as a rule take place within one year of the operation, the results in these 14 cases also demonstrate that the improved methods promise a much larger per cent. of permanent cures, and certainly a longer freedom from recurrence.

We have, therefore, apparently conquered the technic of operations for cancer of the tongue. Now, if we can educate men to come earlier, we shall probably conquer the disease.

THE USE OF IODOFORM OIL IN JOINTS.

By E. G. BRACKETT, M.D., BOSTON,

Surgeon to Orthopedic Department, Massachusetts General Hospital.

[From the Orthopedic Department, of the Massachusetts General Hospital.]

THE use of medicinal agents in joint cavities is by no means new. That it has not always met with success is also well known, yet the principle is sound, and tallies with other surgical methods of today. Many of the results have been bad, because of some disaster, and others negative as to the final results; either because of some of the difficulties of technic and of the great risks under the older methods of surgery, or because of the improper selection of cases. The report of the results in the series here recorded is in the line of similar procedures, but may differ in some matters of technic and in the selection of cases.

There are two general types of cases for which the procedure may be used:—

1. Those in which there is no active disease at the time, and in which the object of the injection is largely mechanical, as, for instance, cases of old arthritis with adhesions, in which it is necessary to keep the surfaces apart after freeing the adhesions.

2. Those in which there is present active disease, and in which the object of the injection is largely for the effect of the agent employed

on the immediate process, as, for instance, in synovitis, of tubercular, Neisser, or other origin.

In all cases, the two things to be considered are:—

I. The special features of the technic of the operation.

II. The type of case to which the procedure is applicable.

I. *The Technic.* Several points in this procedure demand emphasis.

(a) The injection is always made by open incision, and not by puncture.

(b) Sufficient fluid is always left in the joint to keep the capsule under tension, therefore some form of stitch is necessary by which the tension can be maintained at the time of the withdrawal of the syringe.

(c) The adhesions, in old cases, are broken up before the joint is filled with oil and the capsule closed; other than this no violence is done to the cavity of the joint, and no handling is permitted, particularly of the tubercular joints.

(d) Fixation by plaster, or splints, is seldom used.

The Incision. The open incision is always used, and a word is necessary in reference to the choice of this method of inserting the oil, instead of by trocar injection. As a matter of fact the two methods have hardly enough in common to make a comparison. It is often not easy to find the cavity of these joints, not distended with fluid, even by a clear open incision, and, in the acute cases, with the very great edematous swelling of the synovial capsule, it is often difficult to find the cavity of the capsule. The synovial membrane is redundant, the pannus may cover a large part of the articular surfaces, these two surfaces are in contact and roll out into the incision, so as to completely block it, and, in the acute cases, as in the active stages of tuberculosis, the thick, velvety, synovial membrane protrudes into the wound, almost like a hernia. In the older cases with adhesions, the cavity is more or less obliterated, and a trocar will make a small area about its point. In the early cases, with a thick redundant membrane, one has only to experience the difficulty of finding the capsular cavity, even by an open and easy method, in order to realize the difficulty of finding it by a blind trocar puncture. The direct entrance by a clean cut is far more surgical than the blind hunt by a trocar, and likely to do far less damage. Another reason for the open method is the opportunity of obtaining a specimen for examination, to determine the nature of the infection. I believe it justifiable to open in some cases, to insure diagnosis, when the procedure is not dangerous, and may be advantageous. It is only too easy to call a joint case tubercular or otherwise, but in very many cases to which this method is applicable, it is impossible to positively make a diagnosis without further evidence than the clinical signs. The

method is practically safe with extraordinary precaution.

The incision in the skin (preferably on the inner side) (Fig. I.) is a blunt ellipse and the flap dissected back the width of the ellipse. An ellipse with the flap in the opposite direction, shorter than the skin incision, is made in the fascia, and also turned back. It is not always possible to successfully complete this flap when the fascia is very thin and adherent. A straight incision, shorter than the fascia flap, is then made in the capsule, about one-half inch from the border of the patella, in the vertical direction of the leg. The patella lies in the middle or slightly below the middle of the incision.

The Stitch. In order that the capsule may be kept under tension while the fluid is left in the joint cavity, it is necessary to have some form of stitch which not only will make an oil-tight cavity, but also will hold the capsule firmly about the nozzle of the syringe while filling the joint, and will immediately close the opening on its withdrawal. It would seem a very simple method to determine on such a stitch, but several methods were tried before it was found that the most effectual was a double mattress, one enclosing the other. (Fig. 2.) The capsule is closed, beginning at the two ends, and carrying the needle through the fibrous portion of the capsule, but not quite through the synovial portion. By so doing, the soft, spongy synovial tissue is pushed up against the stitch by the pressure of the oil, and acts as a kind of valve. The capsule is closed from either end, with the exception of a space of three-eighths to one-half inch in the centre. Two mattress sutures are then placed in the same manner as the first sutures, one with the opening pointing above, and the other below, the second smaller than the first and enclosed by it. The syringe is then inserted between the threads of the inner mattress, and both drawn firmly about the nozzle. (Fig. III.) When the cavity is fully distended the syringe is withdrawn, the inner mattress is first pulled, then the other, and both tied. (Fig. 4.) The double ellipse flaps give two coverings to the capsular incision, and an additional protection against direct entrance to the joint. Silk is used in all of the sutures.

Preparation Used. Olive oil has been used in all of the cases of the series. Glycerine has been used by many operators and has had much favor, but does not seem to be so well adapted to use in these cases, on account of its hygroscopic properties. It is necessary to have a medium which will resist absorption, and remain in situ to protect the surfaces as long as possible. Very great care should be used in the selection of the oil; it must be of pure quality and of neutral reaction. Very many of the olive oils contain acids, which tend to disorganization and the freeing of some irritative products, as well as to the liberation of the iodine. Some of the irritative results, following the use of oil for surgical purposes, as for instance in the abdominal cav-

ity, have probably been due to this cause. The oil in all of these cases has been the French oil of Antoine Chiris. It is sterilized for one-half hour in boiling water, in a cotton plugged bottle, the top being also covered by rubber tissue, which is kept in place until the oil is used. Enough of the solution is used to fill the joint under tension of its capsule. For an adult joint three and one-half to four ounces is a sufficient amount. The special feature which must be borne in mind in all cases is the necessity of keeping the joint cavity sufficiently distended, to prevent the surface of the synovial capsule from coming in contact either with itself, or with the other articular surfaces, therefore it is necessary that the oil be left with the joint capsule under tension. This is Nature's way of protecting an acutely inflamed joint. In the usual synovitis, an excess of fluid is poured into the joint, separating the synovial surfaces, and this acts in two ways: first, to keep the inflamed surfaces from contact, thus diminishing the pain; and, second, to thus prevent adhesions in the acute stage, when adhesions are formed so easily. When the fluid is early absorbed, or removed, and the surfaces allowed to come in contact, while still acutely inflamed, the pain and sensitiveness, and the tendency of adhesions is increased. By imitating Nature's method,—filling the cavity with fluid,—the freshened surfaces are kept from adhering in the cases in which adhesions have been broken up, and, in cases of active disease, a decided amount of rest is given to the inflamed synovia, through the avoidance of friction.

In old cases with adhesions it is necessary to freely break up all bands in order to obtain as large a capsular cavity as possible. Many old bands are so strong as to give risk of fracturing a portion of the articular surfaces, and these may be cut with much greater safety. Also, experience of many operators has shown that the ordinary forcible straightening of these knees, held firmly by strong adhesions, is attended with a large element of danger. Many times the adhesions are stronger than the bone, which is much atrophied by long restricted use, and force results in a tearing off of a fragment of the articular surface. Puncture by trocar, at the best, will find only a small isolated cavity, and cannot meet the conditions which demand thorough freeing. It is particularly essential, in this class of cases, that sufficient oil be left in the joint to insure the actual distention of the whole capsule. Naturally, fixation in plaster is not indicated.

Selection of Cases. This method is applicable to:—

1. Cases of old infectious arthritis, with adhesions and with partial obliteration of the capsular cavity, but with no true ankylosis, and without destruction of cartilage.

2. Cases of infectious arthritis, in the acute stage, particularly those of Neisser origin, and more especially the type where there is one re-

maining persistent joint, which usually results in ankylosis.

3. Cases of primary synovial tuberculosis before involvement of cartilage or bone.

4. Cases of osteo-arthritis, with marked thickening of capsule and with pannus, in the sensitive stage, but before there has been loss of cartilage.

In other words, the method may be used in the affections of the synovial membrane, either when it is desired to keep the surfaces apart, or when it is desired to apply a remedial agent to the joint surfaces.

For the purpose of illustration, one case (a favorable one) of each of the types to which the method has been found applicable will be cited.

CASE 1. An infectious arthritis of two or three years' standing, which had had several times brisément forcé. At the time of operation the case showed increasing sensitiveness, disability and pain; knee held in flexion of 20°; motion 5 to 10°, all painful; knee swelled; capsule evidently very markedly thickened; patella not movable. X-ray showed fragment of bone on the inner side of the joint, apparently from a fracture of a portion of the articular surface of the inner condyle, and a corresponding space on the under and anterior surface of the inner condyle. Incision was made along the inner side, the capsular cavity was found practically obliterated, with only a few small areas or pockets remaining to indicate the capsular cavity. The patella was so strongly adherent that it could not be freed by either blunt dissection or with the finger, but required the knife or scissors. There were no surfaces of bone exposed, but all articular areas were covered by thick, tough membrane, apparently simply inflammatory in character. On the other side, a piece of bone with cartilage upon one surface was found lodged between the tibia and condyle. This was removed, and the areas of cartilage which were exposed by this were clear and white. About one-half of the capsular cavity was restored. The adhesions on the outer side were so firm as to resist all attempts at freeing, except by actual section, and complete restoration of the joint cavity was not attempted. The joint at this time allowed flexion to 80° and nearly complete extension. The joint cavity was then filled with oil with the capsule under tension.

The case showed practically no pain during convalescence, which was uneventful. Knee kept in fixation for about two months, but slight amount of motion was given each day without force. Patient began walking with crutches in a month and during the following months used leg increasingly.

Twelve months later the knee allowed extension to within 5° of normal; 90° flexion; thickening had markedly disappeared; practically no pain; patella was somewhat movable from side to side. Patient walked without protection about the house and with a splint allowing motion on the street. Eighteen months after operation patient walked several miles daily without cane.

Microscopically the specimen showed an old chronic inflammatory process, and although every attempt was made to find evidence of some tubercular process the examination was negative in this respect.

CASE 2. Neisser infection, polyarticular. All joints cleared in the usual time, excepting one knee, which remained persistently painful, swollen and sensitive; flexed at 20°; no motion on account of marked spasm and pain; no fluid; very marked thickening of capsule. Case apparently one of the types in which in one joint the Neisser infection remained, and there being no fluid to keep the surfaces apart, it gave every indication of a stiff joint from adhesions. No progress had been made for two months. The joint was opened by incision on the inner side, a few adhesions found, but these were easily broken up, the joint moved to a position of 90° flexion, and the capsule, filled with oil, was left under tension. Little pain in convalescence. Motion allowed of a few degrees in a week. After three weeks gentle passive motion was given, which did not cause particular pain. Recovery with full extension; flexion to 110°. Patient has been able to work freely since (2½ years).

CASE 3. Tubercular synovitis of six to eight months' duration. Unable to work for six months; knee flexed at 45°; motion practically nil on account of spasm; whole knee swollen; tender; marked capsular thickening; no fluid; complete disability. X-ray showed no involvement of bone; articular surfaces clean; normal cartilage space between bones. T. B. von Pirquet reaction, positive. Incision on inner side opened to markedly thickened synovial membrane, fragile, vascular, of typical tubercular appearance, pannus extending over to the edges of the articular surface of bone; remaining cartilage clear, smooth; no sign of erosion of any part of joint that could be inspected through the incision. No manipulation was given to the inside of the joint except to remove a small bit of the synovial capsule sufficient for microscopic examination. Oil injection, under tension. Recovery from operation negative. Leg in plaster for several days. Split plaster allowing passive motion daily. At end of eight weeks a second injection was given, and at this time free oil was found in the joint. No weight-bearing was allowed for six months, but patient wore a split plaster and used crutches. At end of eight or nine months patient returned to work with plaster. Two years later leg showed practically normal motion except that of extreme flexion; no swelling; no tenderness. Has worked for 2½ years at his occupation as a plasterer with no return of symptoms. Microscopic examination of specimen showed the process to be tubercular.

CASE 4. Infectious arthritis with hypertrophic areas, with increasing disability for two years. Permanent flexion of knee of 15° to 20°; motion allowed in 25°; marked capsular swelling and villous condition, evident to palpation; pain and sensitiveness with use, to such a degree that disability was nearly complete. X-ray showed hypertrophic process, without involvement of cartilage. This case had a somewhat multiple arthritis, but the knee had been the longest involved and its disability was the greatest. The process was not at the time active. Incision on the inner side showed thickening of whole synovial portion of capsule, pale in color, with pannus extending over onto the edges of the articular surfaces but not covering them, and several large lipomata, which were removed. Oil injection under tension. Motion allowed at end of two weeks. No pain during convalescence. Three months later swelling and capsular thickening had

disappeared; knee could be extended freely and flexed to 100°; no pain on weight-bearing. Eight months later knee had remained quiet; patient was using joint freely; there was no swelling of capsule; crepitus had disappeared.

The list in this series includes cases of the four classes above mentioned operated on by this method. The post-operative period of observation varies from six months to four years. From the study of these cases the following summary has been made:—

The first group comprises the cases of old infectious arthritis with adhesions—object of operation may be called mechanical, viz., to keep apart the freshened surfaces after freeing the adhesions—10 cases. Seven have been successful. Two were unsuccessful and they showed by x-ray destruction of the articular surfaces. These two illustrate the type of case which should not be attempted by this method, viz., the cases in which the arthritis has resulted in destruction of cartilage. In both, this destruction was present, and the operation gained nothing. In the others, motion resulted varying from 45° to 100°. In cases sufficiently severe to warrant an operation it is doubtful if complete motion is ever to be expected.

The second group comprises the cases in the earlier stages of disease. Object of operation is also in a large measure mechanical. This group includes the three types, viz., (a) acute infections (Neisser, etc.), (b) early tubercular synovitis, (c) chronic infectious (osteo-arthritis, etc.).

Type A—Neisser and Acute Infectious. Three cases have been treated, none in the very early stage. With the strong tendency of this form to a spontaneous recovery, it is doubtful if this operative procedure is to be generally advocated in the early period of the disease, for it is recognized that this infection is the most amenable to treatment, and in the usual case, even in the persistent joints, radical measures are rarely necessary. But these joints do occasionally go on to destructive changes, and it is in these persistent ones, before this stage of articular invasion has been reached, that the use of oil is to be advocated. It is in this later stage, where the capsular thickening persists, where adhesions are forming and becoming firm, that the indications for this operative treatment are evident, and has resulted in a rapid diminution of the capsular swelling, disappearance of pain and sensitiveness, with increase of motion.

Type B—Tubercular Synovitis. This type forms the most interesting of the group. It can be definitely stated that the operative procedure is not applicable to cases in which there is involvement of the articular surfaces. The old records of cases show that these cases of tubercular disease of the knee practically all came to ankylosis or to excision; but the records also show that the average case did not present itself for treatment until about two years after

FIG. 1.

FIG. 2.

FIG. 3.

FIG. 4.

the onset of the disease, and at this time the disease was well advanced. At present many cases are seen in which there is no evidence, either clinical or by the x-ray, to indicate involvement of bone, either because diagnosis of these conditions is made in much earlier stages, or because the cases are presenting themselves earlier in the course of the disease. Moreover, the further study of these cases, especially by early operation, shows that a large number of these begin as a primary tuberculosis of the synovial capsule. The excision of these knees, in this stage of purely synovial disease, is a step most radical, yet experience has shown that these cases progress more or less steadily to distinct bone disease. It is in this stage that the attempt has been made to cut short the process by the internal medication of the joint. The usual ultimate result is such an unfortunate one, that attempts to this end are to be desired, provided the method is a safe surgical procedure, and a few favorable early cases gave encouragement. Ely has shown that the cure of the tuberculosis is in many instances by the formation of fibrous tissue, and may serve as the means through which the cure is effected in these cases, but so far there is no proof in this series. One feature of this method, by no means an unimportant one, is the establishment of a positive early diagnosis, a thing not always easy or even possible by clinical signs alone, in the early stages of the process, and the evidence thus obtained in these cases has very materially aided in the general diagnosis of the early tubercular joint lesions.

Thirty cases are available for report. From the standpoint of the results they may be divided into three groups.

1. Cured Cases (8). It is doubtful if the final result of a disease so insidious can be definitely given until a very long time has elapsed. Freedom from symptoms, even for a year, is not sufficient to warrant placing the case among those cured, as has been shown by one of the best early results. An apparent cure has occurred in eight cases, which have stood the test for one and one-half and two and one-half years. The cases were placed as cured by the following tests: Freedom from pain or sensitiveness, maintenance of motion, absence of added deformity, absence of capsular thickening,—all with continuous use.

2. Arrested Cases (14). To this second group belong the less successful and more usual cases, which have not shown definite improvement. There has, however, been a decided checking, as shown by the diminution of (1) sensitiveness, (2) deformity, (3) swelling of the capsule, by the maintenance or increase of motion. In these cases, the x-rays have given no evidence of extension of the disease to the erosion of cartilage or involvement of bone. In other words, the condition has changed from one in which the evidence was pointing toward a progressive process, to a condition in which the

evidence was directed toward a retrogressive process of the disease. The question will arise whether this was not the result of the usual rest and fixation of the treatment, but the fixation was no greater than before, and usually much less, for these joints are not kept in perfect fixation, and the change in the clinical course was too synchronous with the operation to be regarded as a coincidence.

3. The unsuccessful cases (8) are those in which there has been no decided improvement, or the present state cannot be learned. Three were Italians who have quite dropped out of sight. Experience has shown that these are not a favorable class. A few were operated upon in which the x-ray showed a slight involvement of the articular surfaces and the result in these was uniformly bad. The disease was in no way checked, and the cases came later to an excision.

At the present, the most practical value of this operative procedure, in these tubercular cases, depends upon the value which is put on the "checking of the disease." To estimate this, one must have in mind for comparison, the usual course of this disease under fixation and rest. Without exception, the non-operated cases which have been kept under observation, have progressed steadily, by a remitting course, to increasing disease and to increasing disability, and ultimately to excision or to amputation. This applies also to the old cases in the past, whose records have been looked up. Besides the three cured, in seven of these cases there has been decidedly an arrest in the condition as shown by the symptoms, the clinical signs, the x-rays, and the clinical usefulness of the joint. This seems decidedly a gain, particularly when it is considered that this is the stage when excision would not be resorted to, at least as a routine. It must be also borne in mind that the after-care in an out-patient clinic cannot be properly carried out, and many times patients were found walking on the joint three or four months after operation, when fixation had been carried out up to the time of operation.

Type C. Chronic Arthritides. The cases of chronic arthritides in which benefit has been obtained are those of the osteoarthritic type, of the chronic form of the infections, with enlarged villi but without the destruction of cartilage. The procedure is not of use in the atrophic type, or in the atrophic stage of the infectious type. Three cases of this type have been operated upon and without benefit. The cases which have been benefited have shown a decided diminution of the capsular swelling and thickening, diminished sensitiveness and increased use of the joint, and of this type three cases have been operated upon with distinct benefit to each.

CONCLUSIONS.

This method is applicable to the cases of capsular involvement, of various types of infection, and in the stages in which there is no involve-

ment of the articular surfaces. These cases may be divided into two groups:—

1. Cases of old infection in which adhesions have been freed, and it is desirable to keep the surfaces apart. (The use in these cases is largely mechanical.)

2. Cases of infection, of

(a) Acute infection (Neisser, etc.).

(b) Tubercular synovitis (early stage).

(c) Chronic arthritis (selected cases).

The procedure is in no way a substitute for arthroplasty; is not applicable to cases of disease of any origin, in which the x-ray shows involvement of the articular surfaces; but is essentially for those cases, only, in which the abnormal condition is confined to the synovial membrane. In tubercular cases, the early cases naturally show the most marked definite improvement, and in these cases repeated injections are advocated, preferably at intervals of eight to twelve weeks. This procedure should be considered as an incident in the treatment of the tubercular synovitis, not a substitute for the fixation and rest. Neither is permanent fixation desirable. The absolute immobilization is, perhaps, quite as disastrous as the other extreme. A daily gentle passive motion, without forcing the arc of motion allowed, is decidedly a benefit, does not increase the activity of the disease, and does prevent the contraction of the capsule and formation of adhesions.

In Group I, in which the object is largely mechanical, the motion is begun early, within the first two or three days, but never with violence or even to the point of causing pain. In the other cases (a) and (c) of Type A, the acute infectious and the chronic arthritides, motion is also begun early, on the third or fourth day, and given once or twice daily. After two weeks the leg is put in a split plaster, and locomotion allowed without weight-bearing.

Medical Progress.

REPORT OF PROGRESS IN SURGERY.

BY JOHN BAPST BLAKE, M.D., BOSTON.

It becomes yearly more and more impossible to give, within moderate limits, an adequate outline of the progress of general surgery as indicated by current literature. The number of monographs steadily increases, and the duty of the reviewer becomes correspondingly complicated, even with the assistance given by abstracts and indexes. America is certainly contributing, at present, a generous share to the sum-total of investigation and treatment of disease. This is true both in the output of the laboratories and of the surgical wards of the hospital.

In a way, it may be said that conspicuous advances have been made in the broad sub-division of,—anesthesia; surgery of the head; standardizing treatment of fractures; surgical conditions of the thorax; surgery of the stomach; sterilizing the skin both in minor injuries and in preparing for operations.

1. Anesthesia:

There have been at least one hundred important monographs on this subject in the past twelve months. These papers have ranged from the Anoci-association of Crile, through local anesthesia pure and simple, to the new rectal anesthesia of Gwathmey.

RECTAL ANESTHESIA.

Technic for the introduction of oil ether—colonic anesthesia devised by J. T. Gwathmey, M.D., abstracted by Charles Gordon Heyd, M.D.

Contraindications: 1. Whenever ether is contraindicated *except* when the patient has been ill from a previous administration. Here it can be given with impunity. It can also be given in bronchitis, asthma. 2. Pathological conditions of the lower bowel, colitis, hemorrhoids, fistula, etc. 3. When the patient complains very much upon its introduction.

Preparation of the Patient: 1. A laxative, mild, avoid all purging. 2. Irrigation of the colon until the return is clear, three to four hours before operation.

Apparatus: 1. A small catheter and a funnel. 2. Two small rectal tubes inserted side by side to flush and withdraw fluid.

Administration: 1. *a.* One hour before operation: Morphine, $\frac{1}{8}$ to $\frac{1}{4}$; atropin, $\frac{1}{200}$ to $\frac{1}{100}$ by hypo.; *b.* one-half hour before operation per rectum: Chlorotone, gr. 5-10; ether, drams 2-4; olive oil, drams 2-4; 2. *a.* Mixture for adults: Olive oil, oz. 2; ether, oz. 6. Mixture for weak anemic patients: Ether, 55, 65%; oil, 45, 35%. Mixture for children: Ether oil, 50%. *b.* Patient in Sims' position. The catheter is well lubricated and inserted, four inches within the rectum. The mixture poured slowly into the funnel, at least 5 minutes being consumed. Introduce oz. one, for every *twenty* pounds of body weight except for obese. From 10 to 30 minutes is necessary before patient can be moved.

Danger Signals: 1. Loss of lid reflex. 2. Stertor or embarrassed respiration. 3. Approaching cyanosis. If any of the above signs are present withdraw two or three ozs. of the mixture from the rectum immediately. If the breathing is regular with reflexes active, the patient will be relaxed and in surgical narcosis.

Post-Operative: Immediate irrigation of the colon with cold-water soap suds. Then withdraw one of the tubes from rectum and introduce two to four ozs. of olive oil and one pint to one quart of cold water. Withdraw tube from the rectum.