been advocated because of its definite composition, but the reaction with it is often disappointing and tardy.

Because of its permanent strength and prompt action, sodium perborate stands us in good stead. From healthy urines, containing about 1 to 5 mgs. (.001 to .005) of indoxyl salts per liter, I obtained a distinct indigo-blue reaction, uniformly, by the following procedure:

METHOD.

To a test-tube filled one-fourth with urine an equal quantity of concentrated hydrochloric acid is added to liberate the indoxyl-sulphuric acid or urinary indican and then as oxidizer a piece of sodium perborate as large as a full-sized pea. The mixture, which immediately effervesces briskly, is gently agitated to dissolve the perborate. The urine promptly deepens in color and if the amount of indican be large, turns faintly blue. To concentrate the color, one cubic centimeter (16 minims) of chloroform is added, the tube closed with the thumb and the contents *gently shaken for at least two minutes*. The chloroform separates at the bottom as a layer varying in depth of blueness with the proportion of indican. Some pathologic urines yield a layer almost black in color. If the shaking is too vigorous the chloroform is emulsified and remains milky, though with a decided blue tinge.

If the patient is taking potassium iodid, Jaffe's test is always liable to a fallacy from the violet blue color of free iodine in chloroformic solution. If this complication is suspected, the acid fluid is poured off and to the blue chloroform potassium hydroxid is added. Color due to iodine disappears while indigo-blue remains.

RECOVERY FROM BROKEN NECK.*

ARTHUR AYER LAW, M.D.

Instructor in Operative Surgery, Medical Department of the University of Minnesota.
MINNEAPOLIS, MINN.

This case is of interest, not because a broken neck is especially rare, nor is recovery with partial restoration of function very unusual, but complete recovery following operative interference from fracture through the cervical vertebræ is almost unprecedented. I use the term broken neck advisedly—the conclusions of practically all observers have been that a dislocation of the cervical vertebræ without some fracture is impossible. If we are familiar with the bony contour of the vertebræ, with their articular processes placed nearly vertical, we readily understand how a dislocation without a fracture of greater or less magnitude is impossible.

Medical literature contains report of a number of cases of fracture of the cervical vertebræ showing partial recovery with persistent deformity of the neck, greater or less pressure on spinal cord, possible paralysis, bed sores, etc., but I have been able to find but three cases where there was complete and absolute recovery following surgical interference in these cases. The case is as follows:

History.—Roger C., aged 16, weight 110 pounds. On March 6, 1907, he was looking over the gate of a freight elevator, had his chin over the same, the elevator descended, striking the back of his head, and at the same time the gate ascended, catching him at about the region of the larynx, the two lines of force exerting a peculiar shearing pressure on the boy's neck. He was hurried to the City Hospital, where I

saw him twenty minutes later, at which time he presented the clinical picture of pressure on the cord.

Examination.—The boy was unconscious. He was rigid, with violent incoordinate movements, the limbs markedly flexed, all the flexors predominating, complete anesthesia, greatly exaggerated knee jerk, ankle clonus, paralysis of sphincters of rectum and bladder. Breathing was diaphragmatic, shallow and slow. He was cyanotic, the pulse being 120 and thready. His head was slightly crowded back out of position; there was a dimple in the back of the neck over the spine of the third cervicle vertebra. The palpating finger in the pharynx could distinctly make out the prominence of the body of the second vertebra dislocated forward. The boy was *in extremis*, and I advised his parents that he would probably die on the operating table, but he was doomed in any event, and should be given the chance of surgical interference. Dr. J. E. Moore, whom I asked to see him, concurred in this opinion, and the lad's parents consented to the operation.

Operation.—He was placed on his back on the operating table, without anesthesia, his head and shoulders extending over the edge of the table. While the hospital internes held him by his shoulders and body, with one hand under his occiput and the other under his chin (my feet on the edges of the table for a leverage) I made powerful traction on his

head with no other result than twice to dislocate his lower jaw, which had to be reduced. Then with a folded towel over the teeth of his upper jaw to protect my hand, I made traction from here and the occiput. Dr. Moore meantime, with his fingers in the pharynx, crowded back on the vertebræ; this also was unsuccessful, and finally I had an interne grasp my wrists and pull with me; we used all our strength, first extended and then flexed the head on the neck sharply, when the vertebræ slipped into place. The relief to the patient was instantaneous, showing pressure on the cord was relieved; his pulse and respiration quieted down, the incoordinate movements ceased, and sensation returned. He was put to bed with the head of the bed raised, a ten-pound extension on his head pulling from the vertex through the medium of a Barton's bandage, pulleys and weights.

Further Treatment.—Bromids were crowded with the expectation of meningitis developing. He was catheterized for the first twenty-four hours, then voided urine involuntarily, while the bowels responded to enemas. His temperature for the first ten days ranged from 100 F. to 102 F., while his pulse continued high, averaging 120. He had no symptoms of effusion or pressure other than that he was unconscious for twelve days. He was fed by rectum for four days. At the end

* Reported before the Minnesota Academy of Medicine, May 3, 1907.

of the seventh day, still making extension on his head, his body was slipped over the edge of the bed and held by orderlies and internes while I applied an extremely heavy cast from his hips to the top of his head, enveloping him entirely save his face from his mouth to his brows. In the photograph (Figure 1), the cast has been cut away from his chin to dress a small area of pressure necrosis due to the extension. This cast absolutely immobilized his head, neck and shoulders. After the plaster had set the extension was removed. He was left in this cast four weeks. After his return to consciousness on the twelfth day all medication was stopped, and save for a burn on his left arm and leg, resulting from leaking hot water bags applied by a green orderly when first received in the patrol room, he was very comfortable. His temperature had dropped to normal, although his pulse still persisted between 100 and 110. At the end of the fifth week after his injury his cast was removed to find that the boy had made a perfect recovery. He was allowed to get up and go about with a simple collar of plaster, more to remind him to be careful than anything else. At the end of the sixth week he was discharged, cured, with perfect function.

A skiagraph was kindly made for me by Dr. J. Clark Stewart seven weeks after the injury. In comparing it with skiagraphs of a normal neck it could be seen that the second intervertebral space where the dislocation occurred was a little more pronounced, but the vertebrae were in perfect position.

TREATMENT OF HAY FEVER.

W. H. FITZ GERALD, M.D.
HARTFORD, CONN.

As all rhinologists are aware, it is often found that the only areas of contact in the nostrils of the hay-fever patient are formed by congested sections of the anterior third of the inferior turbinate and the resisting nasal septum. Such areas are frequently limited to the very tips of the inferior turbinates, however, as is well known, contact areas may be found in various other sections of the nostrils.

A contraction should be brought about through the application to these areas of a weak solution of cocaine and adrenalin by means of pencil or pledget. If these areas of contact are not extensive, the contraction may be made permanent by the cauterization of their irritable surfaces with chromic or trichlor-acetic acid or both, providing the patient speedily becomes a normal breather and remains such. Atropin sulphate, administered in doses of gr. 1/200 immediately preceding cauterization, will aid contraction and prevent copious secretion. Applications of this drug may be repeated occasionally. It may be found that doses of gr. 1/400 will prove as efficient as the larger dose.

If the obstructive areas are extensive and it is difficult or quite impossible for the patient to get air through the nostril I remove sections of the areas involved wherever they exist, even at the height of the attack, with scissors and snare or punch-forceps, whether congestion or hypertrophy exists; and when it is necessary to prevent contact areas I do not hesitate to remove even extensive areas entire. I have been doing this for the past five years with most excellent results.

If the affection is bilateral it is usually well to leave one nostril for a day or two.

If septal obstructions are present their removal and subsequent nasal breathing may be all that is necessary to bring about a normal condition. As it is usual after nasal operations, the patient should be instructed to sit erect with basin directly beneath the chin until all bleed-

ing has ceased. The nostril will require little or no packing if the patient is mindful of the instruction to breathe through the nose, to keep the nostril operated on as free from clot, etc., as possible, blowing the nose when necessary (never both sides together) and following the same instruction regarding bleeding if there is a recurrence after leaving the operating room, substituting gauze for basin if such a recurrence happens before reaching home.

If the operator decides wedging is necessary he should use just enough sterile wedging to prevent adhesion and only between those sections absolutely requiring it, for it is essential that the patient breathe through that nostril, if not freely, at least as much as is possible. Occasionally it may be necessary to keep a wedge between certain surfaces for a few days, but it is remarkable how speedily healing takes place if the patient is persistent in nasal breathing. A slight coating of chromic and trichlor-acetic acids over small areas may sometimes prove useful following operations to favor contraction and to prevent secondary hemorrhage. It may also prevent the necessity of wedging, as it promotes a crusting which often makes adhesion impossible.

It is well to remember that atropin sulphate, in the above mentioned doses, may be helpful during the healing process in these cases. Also in subsequent years if there is the slightest nasal disturbance due to small recurring contact areas, when the patient has not been entirely faithful to his instruction in breathing, atropin sulphate and absolute nasal breathing will overcome what otherwise might prove a severe recurrence of hay fever.

As I stated in my first letter to *THE JOURNAL*,¹ I do not consider a nostril capable of establishing a contact area, especially below its median line, normal. If, for the purpose of diagnosis, I can establish such an area in the nostrils of a hay-fever patient at any period other than his hay-fever attack, in other words, if I have not seen the patient during his hay-fever period or during an attack of "cold in his head," I irritate the nostrils, and if I am able to bring about a contact area I treat it appropriately, being governed in this by the extent of the area involved.

The nostrils having been relieved of all obstruction, it must be insisted that the patient wear the plaster mouth guard, during sleep at least, as soon as possible, and one may be certain that when his patient informs him that he has been able to wear the plaster the entire night without discomfort he is going to have little further trouble, and this he should be able to do within the first four days of treatment unless the obstructions are more than ordinarily extensive.

When the contact areas are very small it is not at all unusual to find that the patient has worn the plaster without discomfort the entire night following the first treatment. The plaster mouth guard must be worn until the patient has overcome his faulty breathing.

A nasal spray or douche should not be ordered under any circumstances. Air breathed normally through unobstructed nostrils, and nasal secretion, providing the patient is following Nature's general requirements necessary to good health, will bring about a more speedy recovery than any production of the laboratory possibly could.

The patient must be impressed with the necessity of nasal breathing to resist asthmatic attacks during these periods, and he who has had such attacks must be assured that he may expect a recurrence yearly if he does not possess nostrils as nearly normal as possible and use them as Nature intends he should. After the physician

1. *THE JOURNAL*, Sept. 8, 1906, p. 793.