DISCUSSION

This department has been established by the editors in order to afford to those interested in questions relating to economic geology an opportunity for informal discussion. Contributions are cordially invited either in the form of discussion of more formal papers appearing in earlier numbers or bearing upon matters not previously treated. Letters should be directed to the Editor, Sheffield Scientific School of Yale University, New Haven, Conn. The full name of the author should be attached to all communications.

ORIGIN OF THE BENDIGO QUARTZ VEINS.

Sir:—The paper by Dr. F. L. Stillwell on "Replacement in the Bendigo Quartz Veins and its Relation to Gold Deposition" in *ECONOMIC GEOLOGY*, Vol. XIII., pp. 100-111, and the discussion of the paper by Prof. Alan M. Bateman on pages 222-223 of the same volume bring up anew the problem of the origin of those interesting deposits. The "saddle reefs" of the Bendigo Gold-field have made that district famous. This type of ore deposit is also found in Nova Scotia and several other places, but nowhere do the "saddle reefs" seem to be so well developed as in Victoria where they were first recognized. "Saddle reefs" are merely bedded veins which have formed in folded rock strata along the axes of the anticlines. Other types of bedded veins also occur in the Bendigo district, and much of the gold ore has been derived from "spurs"—veins that intersect the beds of sandstone and shale at all angles. These different types of veins are, however, similar in origin, and were evidently formed during a single period of ore formation, for they are everywhere closely related and in places grade into one another.

The "saddle reefs" of Bendigo have usually been attributed to the filling of cavities produced by the arching of the more rigid strata, and some geologists have stated that there is no re-

placement of the country rock by quartz.¹ Dr. Stillwell, in his interesting researches, has now demonstrated that at least some replacement of country rock by quartz has occurred. He does not definitely exclude cavity filling, but strongly emphasizes the part that replacement has played in the formation of the veins. Professor Bateman, in discussing the paper by Stillwell, argues against the exclusive application of the replacement theory, and suggests "that the saddles may represent a combination of both filling and replacement such as takes place in most veins." Neither of these gentlemen refers to the alternative hypothesis adopted by Mr. E. J. Dunn in his beautifully illustrated "Reports on the Bendigo Gold-Field."

In discussing the origin of the "saddle reefs," Dunn states (page 22) that they "have been formed by the deposition of silica from solutions along the axial lines, either in cavities formed by the bending or arching over of beds differing in texture, composition, rigidity, and thickness from one another, . . . or else the saddle reefs filled cavities, more or less occupied by fragments broken off by disruption of the beds along the axes, the quartz filling the interstices first, and then exerting enormous pressure to increase its bulk in a manner common to all substances when crystallizing." In regard to the "leg reefs," which represent the downward extension of the "saddle reefs," he says (page 23): "First the finest film of quartz would be deposited from the solutions that certainly traversed the strata along all the lines of fracture; then these thickened, and wedged open spaces for an extension of the quartz. The solutions of heated water charged with silica were under great pressure, and would consequently penetrate the minutest fissure or crack. Once the silica began to deposit, the crystallizing force would be exerted to widen and enlarge the joints, etc., it occupied, the rift extending as the quartz accumulated."

The views expressed by Dunn are, with certain exceptions, similar to my own, and, since I have recently been making a

¹ Dunn, E. J., "Reports on the Bendigo Gold-Field," Nos. I. and II., *Special Reports, Victoria Department of Mines*, p. 16, 1896.

Lindgren, Waldemar, "Occurrence of Albite in the Bendigo Veins," *ECONOMIC GEOLOGY*, Vol. I., p. 164, 1906.

special study of the mechanics of vein formation, it may be of value to summarize here the evidence in favor of the hypothesis that the Bendigo veins have largely made room for themselves by mechanically displacing the inclosing country rock.

Although, as described by Stillwell, some of the inclusions of country rock have unquestionably been replaced by quartz, there is much evidence that the veins as a whole could not have been formed in this way. According to Dunn (page 16), "the opposite edges of the fissures in the country rock now filled by quartz are all quite angular and correspond in outline frequently on opposite sides of quartz veins"; and (page 27) "the frequent fitting of the country rock on opposite sides of the quartz veins shows how the quartz has gradually wedged the opposite sides apart." This observation was confirmed by Lindgren (page 164) who states that the "opposite sides of the jagged walls of the 'spurs' in many places would fit closely if joined again."

Stillwell notes (page 105) that "in many instances, however, in the larger reefs the included blocks of country rock are not replaced in the slightest degree"; and in his illustration of "preserved brecciation" (Plate V.), as well as in the photographs published by Dunn, the inclusions are angular and have sharply defined boundaries that separate them from the massive white vein quartz.

Specimens of ore from some of the gold-quartz veins of the Swauk district in central Washington contain similar inclusions of black shale, which microscopic examination shows have been largely replaced by silica; but the inclosing white quartz was evidently deposited in a very different manner.² The resemblance is increased by the fact that the silicified shale fragments contain segregated masses of graphitic material frequently arranged in broken lines.

Dr. Stillwell thinks that the laminated quartz, which "is very common and frequently carries the most gold," has originated through replacement of slate, and that this explains the peculiar structure of the ore. This theory is certainly not applicable to

² Taber, Stephen, "The Mechanics of Vein Formation." To appear in an early number of the *Bulletin, Am. Inst. Min. Eng.*

all laminated vein quartz; and there are certain facts concerning the laminated quartz of the Bendigo veins which seem to be irreconcilable with Dr. Stillwell's hypothesis.

Laminated quartz is found in many veins in the crystalline belt of the Southern Appalachians, and, while the gold production from these veins is to-day of comparatively little importance, the belief persists among the older miners that laminated quartz is a favorable indication of the presence of gold in payable quantities. Some of these veins possess a banded structure resulting from the alternation of narrow veinlets of quartz with thin parallel sheets of slate or schist. The veinlets are more or less lenticular in form, pinching out in places to be replaced by others a little to one side or farther along the strike, while frequently the layers of schist are wholly included within the quartz. The wall rock, which is commonly a quartz-sericite schist, shows little if any evidence of replacement by quartz, but often contains pyrite in single crystals and larger masses that partly replace and partly displace the older minerals. The same is true of the layers of schist occurring as inclusions within the vein except that from the thinner layers the quartz, in some cases, has been largely removed, leaving the sericite with a little chlorite and other accessory minerals. In a few rare instances garnets, cyanite and other metamorphic minerals have developed. The veins show every gradation from this banded structure to the typical laminated structure consisting of plates or layers of quartz separated by thin films of sericite.

Now while it is unquestionably true that replacement has usually been a factor in the formation of these veins, there is conclusive evidence that it has not been the most important factor. The veins frequently swell and pinch, and, in extreme cases, consist of a series of lenses disconnected or connected by almost imperceptible stringers. In the immediate vicinity of the veins the folia of the schists are curved to conform with the curvature of the lenses in such a way as to prove that the walls have been gradually pushed apart. Some of the large quartz lenses are subdivided by thin partitions of schist into smaller lenses and

layers.³ In these instances it is clear that structural features, antedating the formation of the veins, have been displaced rather than replaced by the vein quartz.

The illustrations and detailed descriptions given in the reports of Dunn, Rickard and others indicate that the Bendigo veins have been formed by processes that are essentially similar to those which have resulted in the formation of other gold veins, such as those in the Southern Appalachians. The quotations given below are taken from Dunn's reports and require no comment. On page 23 he states: "Some legs consist of a series of lenticular bodies of quartz lying between the same bedding planes, and either connected by a thin quartz film or simply by a joint. Lying as they do between the bedding planes, the quartz forming the legs is generally well laminated." On page 24: "The whole thickness of the quartz in a leg is in some cases laminated (see Plate 15, B), . . . or it is laminated on the footwall, as in Plate 16, B, on the hanging wall (Plate 15, A), or it may consist entirely of white quartz, free from laminations (Plate 12, B)." Again on page 27 he says: "Laminated quartz abounds in all the mines, the quartz alternates with thin black slate films that have been stripped from the walls of the reef during the accretion of the quartz. In thin veins, or close to the walls, the quartz is most laminated. Such quartz is usually considered favorable for the occurrence of gold.

"The quartz along the walls in saddle reefs is generally laminated. As a rule, where quartz reefs cut across the beds there is no lamination, but an exception is met with in Collmann's fault where the walls of the fault were rubbed together so violently that films were formed along the walls. These were again detached as the quartz accumulated.

"Where the reefs are widest there the laminations are further apart, where narrowest, they are closest together. Sulphides appear more commonly in laminated stone, and perhaps gold also; it is certainly very frequent in such stone, the gold generally lies

³ Taber, Stephen, "Geology of the Gold Belt in the James River Basin, Virginia," *Va. Geol. Survey Bulletin* No. VII., pp. 165-168 and 224-229, 1913.

along the black films. . . . Instead of thin laminæ, in some cases thick substantial flakes of slate are included in the quartz."

The "spurs" that cut across the bedding planes in the slate show no evidence of laminations either parallel to the walls or parallel to the structural planes of the adjacent country rock, and yet the "spurs" in places branch off from the "leg reefs" and "saddle reefs." Why should replacement of slate be in any way limited to the veins that run parallel to the bedding planes?

The hypothesis that the Bendigo veins represent the filling of yawning cavities has been widely held in the past, but it is likewise open to serious objections. In several other districts veins have been found occurring in the crests and troughs of folded sedimentary rock strata, but open spaces, similar in form to "saddle reefs" have never been recorded in anticlinal folds, although folded rocks in all stages of dissection are exposed in many parts of the earth.

It does not seem possible that large cavities could have been formed and maintained under the conditions of deep burial and great pressure that existed during the intense folding to which the Lower Ordovician strata of the Bendigo gold-field were subjected. The beds are contorted into a series of folds, the anticlinal axes being only 300 to 1,300 feet apart, and superimposed on these major folds there are in places, especially in the slates, numerous complex minor crenulations along the lines of stratification. The great lateral pressure to which the beds have been subjected is also shown by the development of a slaty cleavage which frequently cuts across the bedding.

Where sharply bent the sandstone has been fractured along parallel or slightly radiating planes so as to give the rock a platy structure, and displacement has occurred along some of these fractures; but, under the same stresses, the more plastic slate has accommodated itself to the displacement by flowing. This is illustrated in Dunn's photographs especially in Plates 4, B and 11, A. Additional evidence of rock flowage is furnished by the way in which the slate beds are usually much thickened in the axes of the folds and correspondingly compressed in the limbs. Under

such conditions of rock flowage is it possible for large cavities to form and remain open?

While lateral pressure might conceivably, under favorable conditions, cause the formation of open cavities in the crests of anticlines, it would certainly tend to close all openings along the limbs of the folds, and yet, according to Dunn (page 23), "cases occur in which the quartz forms over an anticline down a syncline and over another anticline without any interruption," and "in some cases in this field the leg reefs are very persistent . . . and yet there is no development of a body of quartz or saddle reef where the plane they occur in is traced over the anticlinal axis." Lateral pressure can not account for the formation of a series of quartz lenses along the limbs of a fold between the same bedding planes.

It is inconceivable that veins as tortuous as those shown in some of Dunn's photographs (Plate 14, A and B) could ever have been formed in preëxisting open spaces, or that the numerous angular inclusions of country rock (see Plates 20, B and 22, A) could have reached their present position, if the veins were deposited in widely open cavities.

The phenomena discussed in the preceding paragraphs are all difficult or impossible of explanation under any theory of vein formation by replacement or by deposition in open spaces; but they all furnish evidence tending to confirm the hypothesis that the veins have largely made room for themselves by mechanically displacing the country rock. According to the latter concept the veins were deposited from ascending solutions, probably magmatic in origin, which were under great pressure and at high temperature. These solutions penetrated slowly along the lines of least resistance, such as fault fractures and planes of bedding or schistosity. Appreciable openings were filled, and, under favorable conditions, some of the country rock was replaced, but both of these processes were of minor importance. As the vein minerals separated from solution, pressure was developed tending to force apart the inclosing rock and thus make room for the growing veins. The mechanics of the process by which growing

crystals exert pressure has been discussed elsewhere,⁴ and therefore need not be repeated here.

Where the solutions penetrated along closely spaced planes of bedding or schistosity, the thin layers and films of slate were more or less silicified, and at the same time they were gradually separated as the veins increased in size. The laminated quartz originated chiefly in this way. Solutions, entering the zones of complex fracturing, commonly present where beds of the more rigid sandstone had been sharply bent, deposited their minerals along the intersecting fractures, thus forming narrow veinlets which on growing gradually separated the rock fragments and formed the breccia ore. Mineral deposition probably began about favorable nuclei in the larger open spaces and where the least force was required in enlarging the openings. Circulation of solutions through the thin films that separated the growing crystals from one another and from foreign bodies was necessarily slow, and therefore under these conditions diffusion through the solution became an important factor in supplying additional material.

The forms assumed by the growing deposits were largely determined by the nature of the passage ways along with the solutions entered and by the forces resisting enlargement, the tendency always being to produce that shape which required the least expenditure of energy. This explains the growth of the largest deposits—the “saddle reefs”—in the crests of anticlines where the rocks were under least pressure, and also the swelling of “spurs” where they pass from sandstone into slates. Rickard states that “frequently a big ‘blow’ or irregular body of quartz is formed in the slate separating two beds of sandstone.”⁵

The carbonaceous matter in the slates has probably acted as a precipitating agent for the sulphides and possibly for the gold. The sulphides seem to be more abundant near the walls and in vicinity of the slate inclusions. In places the pyrite is found

⁴ Taber, Stephen, “Pressure Phenomena Accompanying the Growth of Crystals,” *Proc. Nat. Acad. Sci.*, Vol. 3, pp. 297–302, 1917.

⁵ Rickard, T. A., “The Bendigo Gold-Field (Second Paper): Ore Deposits Other Than Saddles,” *Trans. Am. Inst. Min. Eng.*, Vol. XXI., p. 688 and Fig. 5, p. 693, 1892.

along the vein walls replacing both slate and vein quartz, thus furnishing additional evidence that the ore-bearing solutions, during the deposition of the vein minerals, were not confined to the widely open portion of partly filled fissures. Lindgren, in describing pyrite crystals found along the contact between the vein quartz and slate, states that they are in the form of cubes "of which one half is embedded in quartz and the other half in the black slate."⁶

STEPHEN TABER.

Sir: In your issue of May, 1918, Mr. Jacob W. Young comments on my paper on the "Halogen Salts of Silver at Wonder, Nevada," and suggests the hypothesis that the iodine and bromine had their origin as emanations from the interior of the earth rather than as solutions from the surface. This explanation of the origin of the silver haloids at Tonopah and Wonder was fully considered before either of my papers on this subject was written, but it was abandoned in favor of the theory of the derivation of the halogens from surface waters because of the controlling weight of field evidence to the contrary. A consideration of chemical facts and laws brought out no unexplainable objections to the theory that I presented, and only served to emphasize the fact that laboratory reactions very incompletely cover the field of possible chemical phenomena. Ostwald writes:

What we call the chemical character of an element and employ for the estimation of its similarities and differences with respect to other elements, is made up of the sum total of the chemical processes, and physical properties known to us which this element and its compounds exhibit. It must, however, be at once admitted that our knowledge of these processes and properties is incomplete and highly onesided. It is incomplete even by reason of the limitation of all human knowledge; it is onesided in the highest degree because of all possible chemical processes, those occurring at medium and slightly higher temperatures are known with incomparably greater thoroughness than, for example, the processes occurring at much higher temperatures. . . . To employ a picture, we know only a certain, and, in part, somewhat arbitrarily cut section of the great organism of chemical facts while the rest of the body remains almost a secret to us.

⁶ Lindgren, Waldemar, *op. cit.*, p. 164.

The questions Mr. Young raises are: (1) Why should the natural order of the precipitation of the halogen silver salts be inverted? (2) How did such an unusual concentration of bromine and iodine in proportion to silver come about? (3) Why should the solutions from which the silver was precipitated be assumed to be alkaline when the literature on the secondary enrichment of ore deposits emphasizes the acidic character of mine waters, especially those in the weathering section of the veins?

The answers to questions (1) and (2) can best be given together. In regard to question (1) I assume that by "natural order" Mr. Young means customary order, and refers to the order of precipitation of the halogen salts of silver customary under laboratory conditions. It must indeed be granted that the ore deposits under discussion were deposited in a natural order. In regard to question (2) Mr. Young appears to be under a misapprehension as to the relative amounts of the halogens in the solutions at Wonder required under my hypothesis. His inference is "that bromine is in amount equal to chlorine, and iodine is in big percentage." This is quite contrary to my conception of the case, which is that the circulating water contained a comparatively large proportion of chlorine, a very small proportion of bromine, and much less iodine. A consideration of the law of mass action, which I need not go fully into here, will suggest that although in the light of laboratory experience we might expect the salts to be precipitated in the order of iodide, bromide, chloride; a sufficiently great preponderance of chlorine over bromine, and bromine over iodine, would cause the order of precipitation to be precisely opposite. It is not necessary that all the chlorine be precipitated from the solution before the precipitation of bromine begins, but only that the concentration of chlorine be reduced to a point where mass action will favor the precipitation of bromine. In like manner, the precipitation of iodine may begin before either the chlorine or bromine are eliminated from solution, and consequently the quantities of the halogens found in the orebody have no recognizable relation to their relative proportions in the solution. To quote Ostwald again: "Not only the nature and, say, the temperature are the

determining factors for the occurrence of a chemical process, but also the *ratio of the substances present to the given volume or the concentration as well.*" The concentration of solutions worked with in the laboratories are so immensely greater, and the time of reaction so infinitesimally less than those involved in ore deposition, that the results of the laboratory cannot always be compared with those of nature, especially when facts and natural laws indicate the possibility of different results.

The occurrence of these salts in Nevada in the form of crystals, suggests a set of conditions of deposition very different from those of test tube reactions. To me it indicates concentrations so low as to be within the range of solubility of the halogen silver salts. Obviously we are here dealing with conditions about which we have no experimental evidence.

With regard to question (3), the difference should be emphasized between this class of ore deposits, which contain minute proportions of sulphide minerals, as compared with heavy deposits of base sulphides. In the Wonder vein descending alkaline surface waters would remain alkaline to a much greater depth than in a deposit heavy with sulphides. However, this would make no difference in regard to the formation of the silver haloids, as they are readily precipitated in acid solution. In this connection it should be remembered that there was practically no transportation of silver and the secondary minerals were found either on, or within a few inches of, the ore from which the silver was derived. My favorite conception of the process of formation of these minerals is that the alteration took place above the level of the ground water, as follows: Consider a piece of rich silver sulphide, cracked so that water can percolate through it, and lying in such a position that a slow seepage of oxidizing surface water passes through it. As a result of oxidation silver from the ore goes into solution as a sulphate, but it is immediately combined with one of the halogens and its precipitation begins. Its concentration may be so low that the silver salt will remain in solution long enough to be carried a short distance before the deposition is complete, and as a consequence we find crystals instead of amorphous precipitate. While I have pictured this as

taking place above water level, there is no serious objection to considering it as occurring in slowly moving solutions below water level.

The occurrence of the alkaline salts of chlorine, bromine and iodine at the locus of the orebody may be due either to wind blown salt-dust from nearby saline lakebeds, or from similar beds or salt lakes existing at the place before erosion occurred to its present extent.

Although I have here given first the theoretical considerations upholding my theory, their acceptance was, as I have already indicated, forced on me by the weight of field evidence. Where silver chloride is found, its invariable occurrence on the upper edges of silver-bearing veins, extending from the surface downward but limited to the depth of the oxidized zone, has never been questioned as being the result of action by surface water. I do not think it is necessary to review the arguments on this point. In the deposits that I have described the interrelations between the chloride, bromide, and iodide of silver were such that they must have had a common origin. In Tonopah they were found only in outcropping veins although there were numerous veins that did not have outcrops. At both Tonopah and Wonder there had been a large but undetermined amount of surface erosion, and it is not improbable that at Wonder a zone of silver chloride had been eroded away. Possibly this is the explanation of the absence of the chloride at Wonder.

In concluding this argument I would say it was inevitable that the theory proposed by Mr. Young should arise, and in my paper on the Tonopah deposits I anticipated it and gave reasons to the contrary; but, since the discussion has resulted in a more detailed exposition of the facts as well as of my theory, I thank Mr. Young for his interest in the subject.

Mr. Waldemar Lindgren's contribution telling of a description written fifty years ago of a similar deposit in Mexico is of interest. It is remarkable that there have been so few contributions to literature of descriptions of the "horn silvers," although they occur commonly enough.

J. A. BURGESS.

Sir: The paper by Victor Ziegler on "The Movements of Oil and Gas through Rocks," although brief, is the clearest and most complete review of the workable theories of oil migration. I agree with all its major features and conclusions, and will make only a few remarks on minor points.

Professor Ziegler calls attention to the possible effects of adhesion and adsorption of oil on various rock substances. Both adhesion and adsorption are phenomena that retard migration. They are interesting in their selective action on various parts of crude oil, as shown by filtration through fullers earth. From these experiments it is evident that different oils have different powers of adherence to a given substance; also that different substances hold an oil fraction with different powers of adsorption or adherence.

It appears that unsaturated hydrocarbons are absorbed on rock surfaces more strongly than saturated hydrocarbons. It appears that heavy oil fractions adhere more strongly than light fractions. It appears that clay surfaces are more efficient than quartz surfaces in producing these effects; also that they are increased by increase of area of surface through comminution of the solid particles. It appears also that oils are made heavier when they are subjected to these processes of fractionation, if all the fractions are again brought together.

There is much mystery about adsorption and adhesion. We still lack a complete explanation of the oil-flotation process of concentrating metallic ores which exhibits related phenomena. No one knows just why the grease tables at Kimberley are successful in separating diamonds from pulverulent "blue-ground." The diamonds stick to the grease, and the other minerals slip over it and pass off the table.

It is generally thought that molecular forces are involved. I have crystals of calcite with inclusions of oil, in many very thin green layers that tend remarkably to follow certain buried faces of the growing crystal.

¹ Discussion of paper by Victor Ziegler, *ECON. GEOL.*, Vol. XIII., No. 5, July, 1918, pp. 335-348.

A complete explanation of these phenomena seems to require further advance in the science of physics. Their main interest lies in their mystery.

I have a crystal of rock salt containing square inclusions of oil and round inclusions of gas. The difference probably indicates some mystery of crystallization.

The ability of oil to penetrate clay-shale is proven by its commercial occurrence in crevices in clay-shale. At Florence, Colo., and other places, these crevices do not connect with any sandstone or other porous stratum. The oil seems to have been formed in the shale. Regardless of its origin it must have migrated through clay-shale to reach the crevices.

I have hand specimens of basalt in which the steam-blebs contain zeolites with needles projecting to the center of the vesicles, showing that the cavities were once full of water. They are now full of the residue of oil. This oil must have penetrated the basalt for several inches and also a few millimeters of calcite and zeolite. Doubtless the oil made its way along invisible capillary passages between the rock particles and crystals.

CHESTER W. WASHBURN.