chief to the structures of the middle or internal ear, or that the pain and discomfort incident to this process would offset the anesthetic qualities obtained, but practical experience has proven these fears unfounded.

My first experiment with this method was done on a patient suffering from a left facial paralysis of seven months' standing; the left ear was filled with exudate and almost deaf. The case is more fully reported by my friend, Dr. H. O. Reik,¹ who suggested a paracentesis of the drum membrane on the affected side for the relief of the paralysis. A jet of ethyl chlorid was allowed to play into the canal for some seconds and the incision was then made. With the exception of a slight burning sensation due to the ethyl chlorid nothing was felt by the patient, who was a very nervous woman. She almost completely recovered from her facial paralysis and her hearing returned to normal. Encouraged by this success, which demonstrated the harmlessness of the procedure, I have since used this freezing method on every available case that presented itself during my service at the above mentioned clinics and also in my private practice. In no instance was the subsequent course of the case any different from what might have been expected, had ethyl chlorid not been used.

In my earlier cases, I used an ordinary ethyl chlorid tube held twelve to fifteen inches from the ear and allowed a jet of the liquid to play into the auditory canal through the previously inserted speculum. This had some disadvantages. One had no assurance that the drug actually reached the desired spot and hence anesthesia was sometimes imperfect; much liquid was wasted and more than necessary was injected, producing more or less pain and discomfort to the patient, and it was almost impossible to avoid getting some of the liquid on the face or even into the eye of the patient, unless previously protected by a towel.

These disadvantages have been completely eliminated by a new nozzle especially designed for the purpose and made for me by the Gebauer Chemical Co. of Cleveland, Ohio. This nozzle is about two inches long, very slender and is attached to the metal container at such an angle as to permit a good view of the parts beyond when partly introduced into the speculum. The apparatus is held in the same manner as any ear instrument, the body of the container forming the handle and a very fine *spray*, not a jet, is easily directed to the desired spot (Figs. 1 and 2).

In practice, I begin with an almost imperceptible spray, which I allow to play on the part, until the patient feels a certain degree of coolness within the ear, then very gradually increase the volume of the spray by slowly opening the valve, at the same time facilitating its rapid evaporation by means of a jet of air blown into the canal from a Politzer bag. With a little practice one soon learns to know how far it is necessary to carry the procedure in order to obtain a sufficient degree of anesthesia. For my part, I usually shut off the spray when the patient begins to complain of an aching sensation, then blow in a little more air and operate; the whole procedure taking but a few minutes. A minimum amount of ethyl chlorid is used and none wasted, an item of some importance when dealing with such an expensive drug.

In searching the literature on the subject I find

that Brieger² mentions the use of ethyl chlorid for similar purposes, but up to the present time have not been able to get access to his paper and am not familiar with his method of application.

In addition to the usual minor operations about the ear canal, such as paracentesis, incising furuncles, extracting polypi, etc., I have recently begun to use ethyl chlorid in the manner above described as a preliminary to the application of Lucæ's sound in the treatment of chronic otitis media and find it an ideal way to avoid pain, the one objectionable feature of that form of treatment. The results thus far seem to indicate a far greater percentage of markedly improved cases than I was able to obtain before introducing the preliminary freezing; however, further experience is necessary to more fully confirm this statement, and I shall take pleasure to report on the same at a future date.

449 East Twenty-second Street.

THE TRANSMISSION OF DISEASE BY THE MOSQUITO.*

JOHN R. TAYLOR,

Chief of the Laboratory of Las Animas Hospital.
HAVANA, CUBA.

The study of the life and habits of mosquitoes, not only in their adult stage, but in their cycle of development from the egg to the mosquito, is one of the most fascinating subjects that a person who has the time could possibly take up. Not only is the subject fascinating, but it is also of great importance that the life history of all species of mosquitoes should be understood, as they have been proved beyond a doubt to play a prominent part in the transmission of disease to human beings and animals, and more especially is this true of malaria, yellow fever and filariasis.

Mosquitoes belong to the order of *Diptera*, of the family *Culicidæ*. They are differentiated from other insects which simulate them by, 1. The piercing mouth. 2. The scales on the veins of the wings. 3. The particular venation of the wings.

There are four stages in the life of the mosquito, the egg, larva, pupa and imago or winged insect. The early stage of the mosquito is always aquatic.

EGGS.

The eggs are deposited habitually on the surface of either natural or artificial collections of water. Occasionally, however, they may be laid on moist soil, or on grass above the water, and hatch out later when water reaches them. The eggs are laid in one of two ways, either singly, each egg lying on its side (*Stegomyia fasciata*); or in a boat-shaped mass containing from 200 to 400 eggs all glued together, each egg resting on its end (*Culex*). The eggs hatch out in from 12 hours to 2 days.

LARVÆ.

The larvæ are eruciform, and easily recognized in the water by their wriggling movements. They have a disproportionately large head. The larvæ of the subfamily *Anophelina* habitually rest in a horizontal position at the surface of the water. Other species hold themselves in an oblique position, head downward, at the surface. The larvæ remain in the larval stage about a minimum of six days, but the duration may be much longer.

1. Reik: "The Relationship Between Otitis Media (non-supuration) and Facial Paralysis of the Refrigeratory Types." Trans. Am. Otological Soc., July, 1904, p. 376.

2. Brieger: *Klinische Beiträge zur Ohrenheilkunde*, 1896.
* Read before the St. Louis Medical Society, 1904.

PUPÆ.

The pupæ, third stage of the mosquito, differ from the larvæ in the great swelling of the thoracic segments. They habitually remain at the surface of the water, unless disturbed, when they rapidly descend. They remain in the pupæ stage about 24 hours to 3 days, when they hatch out and become mosquitoes.

MALARIA.

The mosquito malaria theory was held by the ancient Roman writers nearly 2,000 years ago. The same belief prevailed among the peasants in certain parts of Italy and Tyrol, and among barbarous tribes in the malarial districts of Africa. Many other theories prevailed, those most generally accepted being that the disease was due to foul air, night mists, odors from swamps or marshy soil, stagnant water, upturned earth, etc.

In more recent years, we find the mosquito malaria theory advanced in a scientific manner by Americans. Dr. J. E. Nott, in 1848, discussed the question of the transmission of malaria by mosquitoes, being convinced that they transmitted the disease. Dr. A. F. A. King, in 1833, wrote a very thorough article on the etiologic relation of mosquitoes to malaria. Laveran, in 1891, and Koch, in 1892, believed in the theory, but without producing any facts in its support. Dr. Patrick Manson, in 1894, offered the most plausible theory, stating his belief that it must be carried by some blood sucking insect, and from the habits and distribution of the mosquito, he believed the disease was transmitted by this insect. Dr. A. Bignami, in 1894, was probably the first to definitely bring out the theory that the mosquito conveys the infection of malaria by inoculation.

Dr. Ronald Ross, in 1895, began to work on the mosquito theory. He allowed the insects to bite cases of malaria, but failed at that time to get his mosquitoes infected, due no doubt to the fact that he was experimenting with the *Culex* species. In 1897, Ross tried to infect *Anopheles* mosquitoes that he had not tested previously. On examining the insects after a day or two, he was able to find positive evidence of malarial infection in the stomach. The walls of this organ were found to contain encysted parasites (zygotes).

Dr. W. G. MacCallum, in 1897, proved that the gametocytes are sexual forms, describing the process of the micro-gametocyte flagellating and penetrating the macro-gamete. He was probably the first to recognize the true sexual process that takes place between the male and female parasites of malaria.

Dr. Ronald Ross, in 1898, followed out a series of experiments with the malaria of birds, using the *Culex* mosquitoes. He was able to demonstrate the cycle of development of the parasite, from the time that the mosquito sucked into its stomach the blood of an infected bird, until the parasite reached the salivary glands of the insect. Later he succeeded in infecting healthy birds by allowing them to be bitten with infected mosquitoes. What Ross proved in connection with sparrow, or bird malaria, was taken up by Italian scientists, especially Prof. Grassi, who worked out a similar relation between the *Anopheles* mosquito and human malaria. Grassi was able to trace the whole process, step by step, bringing together and applying to the human parasite the knowledge gained by previous workers in different parts of the world, until at the present time no single point in the cycle of development is incomplete.

Dr. Patrick Manson, in 1900, conducted an experiment in London, to test the mosquito-malaria theory. It was not necessary that this should be done to prove

the positiveness of the theory, but it served to add another link to the already long chain of evidence, proving that malaria is transmitted to the human being by the medium of the mosquito. For this experiment insects were infected in Italy, then sent to London, and allowed to feed on two men, Dr. Thurburn Manson and Mr. R. Warren of the London School of Tropical Medicine. In each case the fever developed exactly eighteen days after they were bitten, and the benign tertian parasites were found in their blood by competent observers.

Drs. Sambon and Low, with two others, in 1900, volunteered to spend the fever season in Ostia, Central Italy, the most malarial locality in the Roman Campagna, with no protection against malaria, except to avoid being bitten by mosquitoes between sunset and sunrise, the hours during which the *Anopheles* usually bite. This protection they secured by means of a carefully constructed hut, the windows and doors of which were covered with wire gauze, into which hut they retired an hour before sunset. They moved freely about the country during the day time. They lived in this manner for a period of three months, using no quinin or other prophylactic, and not one of the four contracted malarial fever.

A control experiment was inadvertently made during their residence in Ostia. After the assassination of King Humbert, which occurred about that time, fifteen or sixteen police agents were sent to Ostia to arrest suspected anarchists. Though these police spent only a part of the night in the district, every one of them developed fever about two weeks later.

Experiments have also been made by the Japanese army on the Island of Formosa. Two companies of soldiers were stationed in a very malarial locality. The men of one company were very carefully protected from the bites of mosquitoes, and did not suffer from attacks of malarial fever, while those of the other company who were not protected from mosquito bites suffered severely.

Many similar experiments and scientific investigations have been made, so that we now know that a certain species of mosquito is, so far as is known up to the present time, the only natural means of transmission of malaria. The *Anophelina* mosquitoes are the intermediary hosts necessary to complete the cycle of development in malaria. In the case of yellow fever it is a different species of mosquito (*Stegomyia fasciata*) that acts as the intermediary host. The conditions necessary for the existence of the parasites of human malaria and yellow fever are only to be found in the human blood, and in the particular species of mosquito.

Mosquitoes are never born with the disease, it is only when they are given an opportunity to suck the blood of a person having malaria or yellow fever, that they become infected. The parasites have never been found in water, earth or in the air. Experiments made in different parts of the world indicate that the infected mosquito does not transmit the infection to its offspring.

To become infected with malaria, the *Anophelina* mosquito must bite a patient whose blood contains either pigmented spheres or crescent-shaped bodies. The sporulating form of the parasite will not infect the mosquito. It requires from seven to fourteen days, after the insect has bitten a case of malaria, before it is able to transmit the infection to a human being. This period constitutes the time necessary for the parasite to go through its cycle of development, and to reach the salivary glands. It is only then that the mosquito is able to

infect. At the time of the next bite, while the insect is taking up blood with one tube, it is pouring out saliva from the salivary glands with another tube, and, with the saliva, the parasite of malaria.

A somewhat similar process takes place in connection with yellow fever, but the cycle requires at least twelve days. There is this difference, however, that to be infected with yellow fever, the *Stegomyia fasciata* must have bitten a patient during the first three days of the illness, otherwise it will not acquire the infection.

So far as we know, no other animal shares with the human being the capability to be the intermediary host of yellow fever, or of the human malarial parasite. Intracorporeal parasites akin to malaria have been found in many animals, in cattle, horses, sheep, dogs, monkeys, frogs, bats, sparrows, finches, crows, hawks, owls, pigeons, blackbirds, etc., but the parasites are of a distinct species.

Dionisi has found a hematozoon of the bat, which resembles human varieties very much. Koch specially investigated the possibility of transmitting human malaria to the lower animals in Java. He used the orang-outang and hylobates, but failed entirely to produce the disease in these higher apes, as other investigators have failed with other animals.

Mattei injected blood of malarial patients into sixteen pigeons, with negative results. The same negative result was obtained in four instances, in which he injected the blood of infected birds into the arms of healthy men.

The incubation period of malaria in the human being, from the time of the bite by an infected mosquito, varies from two days to three weeks, depending on the particular kind of parasite, the amount of blood taken in by the mosquito, and its richness in parasites.

OPPOSING THEORIES.

It may seem absurd to take up the reasons, or at least some of them, that are given to combat the mosquito-malaria theory. However, it may not be out of place to mention a few of the objections:

Malarial Localities.—It is said that certain places are malarious, in which there are no mosquitoes. I think, if careful search were made, it would be found that the malarial bearing mosquitoes were present.

Foul Air of Swamps.—Some infer that the disease is acquired from foul air. The gas given off from the mud of stagnant pools and swamps has been found to be a compound of hydrogen and carbon-methane, or marsh gas, which can be easily proved to have nothing to do with the malarial infection.

Contaminated Water.—There are still a few persons who maintain that the disease is contracted by drinking infected water. It is possible that many of the supposed cases of malaria from drinking bad water were typhoid cases. Those who have studied the malarial parasites can not understand the possibility of the germ living or propagating in water of any kind.

Night Air.—The idea that people get malaria, as a result of being out at night, is explained by the fact that the anophelina, as a rule, only bite at night.

Upturned Earth.—Epidemics of malaria are known to break out among men engaged in work requiring up-lifting of the soil. In these cases the infection is no doubt due to the fact that some of the new arrivals had malarial parasites in their blood when they came, also that conditions were more favorable for the breeding of mosquitoes, on account of the new pools and ditches formed during the digging operations.

It is important that we meet wrong ideas as to the cause of malaria, by disseminating the knowledge that we now possess, either by education or by vigorous sanitary laws if necessary.

YELLOW FEVER.

From observations made in connection with yellow fever, we have reason to believe that infection can not take place except in one of two ways, either by the bite of an infected mosquito, or by direct inoculation of infected blood. Over two hundred experiments have been made on human beings. About 128 of these experiments were mosquito inoculations; the persons being bitten by mosquitoes infected, or supposed to be infected, with the germ of yellow fever. The others were direct blood inoculations, except three. Two of the experiments consisted in making an excoriation on the skin, one centimeter square, by means of scraping and then applying a drop of virulent serum from a case of yellow fever. The result was negative. Another experiment was made, whereby a non-immune subject was allowed to drink water, in which had been triturated the bodies of four infected *Stegomyia fasciata* mosquitoes, the result being negative.

Dr. Charles P. Finlay, at the session of the Royal Academy of Sciences, in Havana, Cuba, August 14, 1881, first gave to the world his eventful theory of the transmission of yellow fever by the medium of the mosquito, reporting, at that time, a series of experiments that he had made on human beings, by which he claimed to have proved his theory. He was the pioneer in mosquito inoculations. From 1881 to 1900, Dr. Finlay, assisted by Dr. Claudio Delgado, performed over one hundred inoculations with supposed infected yellow fever mosquitoes. Time will not permit me to dwell on these experiments, I will simply read an extract of Dr. Finlay's original theory given out in August, 1881:

Three conditions will, therefore, be necessary in order that yellow fever may be propagated: 1. The existence of a yellow fever patient into whose capillaries the mosquito is able to drive its sting and to impregnate it with virulent particles, at an appropriate stage of the disease. 2. That the life of the mosquito be spared after its bite on the patient until it has a chance of biting the person in whom the disease is to be reproduced. 3. The coincidence that some of the persons whom the same mosquito happens to bite thereafter, shall be susceptible of contracting the disease.

Dr. Finlay's theory was confirmed by the American Military Yellow Fever Commission composed of Drs. Reed, Carroll, Agramonte and Lazear. The positive results obtained by this commission, experimenting in Cuba in 1900 and 1901, proved beyond any question of doubt, that yellow fever is only contracted, naturally, by the bite of the mosquito. They also proved that contact with the patient, fomites, clothing, etc., plays no part in the transmission of the disease.

At the present time, Dr. Finlay explains the process of yellow fever infection in the following manner. He assumes that the germ is a two host, microscopic, perhaps ultra-microscopic endo-parasite of the female of a particular mosquito (*Stegomyia fasciata*). In order that it may complete its life cycle, perpetuate its species, and find its way into other healthy stegomyias, it is necessary that it should penetrate from an infected insect into the body of a non-immune person.

It appears that after the germ leaves the infected insect, and enters the body of the susceptible person, it passes through a certain cycle or evolution, requiring usually three to six days, although the period of incuba-

tion in the human being may be extended to at least thirteen days in exceptional cases.

Dr. Finlay holds that the penetration of the germ into the body of the non-immune, is intended for the accomplishment of its sexual reproduction, and that the analogy between it and the malarial parasite holds good, with the difference only, that the latter accomplishes this act in one of the anophelina mosquitoes, while the sexual cycle of the yellow fever germ is supposed to take place in the human host. He suggests the following as the probable cycle of development of the germ in the body of the contaminated mosquito:

- a. Introduction of young sporozoites sucked up with the blood of the patient.
- b. Growth of the sporozoites to adult age.
- c. Formation of schizonts and merozoites.
- d. Formation of gametes.

In the body of the non-immune, counting from the time of inoculation by the bite of the infected *stegomyia*, the following stages, in his opinion, are likely to occur during the period of incubation:

- a. Localization of the gametes in appropriate sites.
- b. Fecundation of the macrogametes by the microgametes of the yellow fever germ.
- c. Formation of the ookinet, and development of the oocyst in appropriate cells (endothelial?).
- d. Discharge of free sporozoites of yellow fever into the circulating blood (perhaps ultra microscopic).
- e. Development of the toxemic symptoms characteristic of the invasion of yellow fever in the inoculated non-immune.

The following facts, in connection with yellow fever, are now very generally admitted:¹

1. The only natural method that is known by which yellow fever is transmitted from the sick to the healthy person is by the bite of mosquitoes previously infected by biting a sick person in the first three days of an attack of yellow fever.
2. On the fourth day of the disease the blood no longer contains the virus of yellow fever, even when the fever is high.
3. Up to the present time it has not been proven that yellow fever is transmitted by any other species of mosquito except the one known as *Stegomyia fasciata*.
4. The infected mosquito is only dangerous after an interval of twelve days from the time that it has ingested virulent blood.
5. Contact with a patient, personal effects, clothing, excretions, fomites, etc., are incapable of producing yellow fever.
6. One-tenth of a cubic centimeter of virulent serum injected under the skin is sufficient to cause yellow fever.
7. The virus of yellow fever rubbed on the denuded skin does not give the disease.
8. The virus of yellow fever preserves its vitality and virulence in defibrinated blood for a period of five days, when protected from the air, preserved under liquid vaselin, at a temperature of 24 C. to 30 C. At the end of eight days under similar conditions the virus is no longer active.
9. Virulent serum exposed to the air at a temperature of 24 C. to 30 C. is inactive at the end of forty-eight hours.
10. The virulent serum becomes harmless after heating for five minutes at a temperature of 55 C.
11. In the serum of the patient the virus of yellow fever

1. These conclusions are based on the experiments and investigations of the following workers: Drs. Charles J. Finlay and Claudio Delgado, in Havana, 1881 to 1900; Dr. A. Agramonte, in Havana, 1899; American Military Commission, composed of Major W. Reed, M.D., and Drs. J. Carroll, A. Agramonte and J. W. Lazear, in Havana, 1900 and 1901; Dr. J. Guiteras, in Havana, 1901; Commission Working Party No. 1 of the Yellow Fever Institute, Public Health and Marine-Hospital Service, composed of Drs. H. B. Parker, G. E. Beyer and O. L. Pothier, in Vera Cruz, 1902; Dr. A. Agramonte, in Vera Cruz and Mexico City, 1902; French Commission, composed of Drs. Marchoux, Salimbeni and Simond, in Rio Janeiro, 1902 and 1903; Brazilian Commission, appointed by Dr. Emilio Ribas, composed of Drs. L. P. Bareto, A. Gomez de Silva and Julio de Barros, in Sao Paulo, 1903.

passes through a Berkenfeld or Chamberland filter without dilution, providing the density of the filter is not too great.

12. The blood serum of a yellow fever convalescent presents preventive properties. The immunity produced by such serum is still present at the end of twenty-six days. The serum also appears to possess therapeutic properties.

13. The usual period of incubation of yellow fever is from three to six days.

14. Apart from the natural transmission by means of the bites of infected *Stegomyia fasciata* mosquitoes, the only other proved method by which it has been possible to transmit yellow fever artificially consists in injecting non-immunes subcutaneously with yellow fever blood.

15. Experimental yellow fever has been produced by the bite of an infected *Stegomyia fasciata* mosquito fifty-seven days after it acquired the infection.

16. Yellow fever can only spread in regions where the *Stegomyia fasciata* prevail.

17. It is possible to eliminate yellow fever from the world by three methods: 1. Mosquito proof isolation of all cases, including those suspicious. 2. Thorough fumigation of patients' quarters and those adjoining. 3. Attempt to destroy all mosquitoes in all their stages.

18. Neither in the mosquito nor in the blood or any other tissue has anyone succeeded in discovering the germ of yellow fever.

EXTERMINATION OF HUMAN MALARIA.

We have reason to believe that human malaria would disappear from the world, if any of the following methods could be put into effect:

1. If the parasites could be eliminated from the blood of all malarial persons by quinin or other method. This is difficult, because we could never get at all infected persons.
2. If all healthy mosquitoes could be prevented from biting infected persons; or if infected mosquitoes could be prevented from biting healthy persons. This method would be practicable if everybody were to adopt preventive measures against the mosquito.
3. If all the anophelina mosquitoes could be destroyed. This is possible if all the people in all localities of all countries begin, and keep up, a vigorous crusade against the mosquito in all its stages.

Unfortunately, not one of these methods is capable of universal application, and while we may or may not be able to entirely eliminate malaria, there is no question of doubt but that we can reduce the number of cases to a minimum.

EXTERMINATION OF YELLOW FEVER.

From the positive results obtained in Cuba in 1901, under the direction of Col. W. C. Gorgas, of the American Government of Intervention, and still maintained by the Cuban sanitary authorities, we have reason to state positively that yellow fever could be eliminated from the world by the use of measures similar to those used in Cuba, as follows:

1. Destroy as many mosquitoes as possible, so as to limit the number of insects capable of transmitting the disease. We know that if there were no mosquitoes there would be no yellow fever.
2. Prevent mosquitoes from biting yellow fever patients. If we prevent the mosquitoes from getting infected we also prevent yellow fever. This is, perhaps, the most important measure, namely, strict mosquito-proof isolation of all cases of yellow fever and all suspicious cases.
3. Destroy, by thorough fumigation, all mosquitoes that may have become infected by biting yellow fever cases.

EXTERMINATION OF FILARIASIS.

We know that both *Culex* and *Anopheles* species of mosquitoes act as intermediary hosts of the filaria; the worm going through its cycle of development in the body of the insect. After a period of days the insects

are capable of transmitting the infection to human beings. We can only expect to exterminate, or at least diminish the disease, by similar methods as against yellow fever:

1. Compulsory mosquito-proof isolation.
2. Thorough fumigation of patients' quarters, etc.
3. Destruction of all mosquitoes in all their stages.

QUARANTINE.

We should endeavor by all means possible to prevent importation of all persons having transmissible diseases. Quarantine, if strict, will exclude yellow fever and diminish malaria in any country. However, should yellow fever gain entrance to a place, it is easy to control and exterminate by the methods above mentioned.

All persons on steamers, coming from countries where yellow fever is prevalent, should not be allowed to land until a period of five days has elapsed, counting from the time that the steamer set sail from the infected port, and providing no suspicious case has occurred during the trip. If there has been a suspicious case of sickness on board, quarantine should be enforced for at least five days after arrival of steamer. (Persons who present a satisfactory certificate of immunity, may be allowed exemption from quarantine.) The temperature of all persons in quarantine should be taken morning, noon and night, and any one having the least rise of temperature should be at once placed in a mosquito-proof isolation hospital for further observation.

Even if no sickness has occurred on the steamer, quarantine should be enforced for at least one day after arrival of a steamer from a malarial infected country. Temperature of all persons should be taken, and any one having a rise of temperature, sufficient to arouse suspicion, should have his blood examined and, if found to contain malarial parasites, he should be detained in a mosquito-proof hospital, and put under medical treatment until the parasites are known to have disappeared from the peripheral blood, before he is allowed to land.

INFECTED PERSONS.

When persons are found to have malaria or yellow fever, they should be compelled to undergo isolation in mosquito-proof rooms. The bed should also be covered with mosquito gauze so as to afford double protection. In the case of malaria, the patients need only be protected, as a rule, between sunset and sunrise, the time that the anophelina usually bite. In yellow fever, the patients should be protected against mosquitoes day and night, during the early stages of the disease. All persons suspected of having yellow fever must be reported to the board of health, who will immediately arrange to have the patient conveyed in a mosquito-proof ambulance to a specially constructed mosquito-proof hospital. The patient should be covered while being carried to the ward, so as to prevent any possibility of being bitten by mosquitoes. When the patient is inside the special hospital, all danger of spread of the disease is eliminated.

After removal of the suspected patient from his quarters, all the rooms in the building, and in the adjoining houses, must be immediately fumigated, so as to kill any mosquitoes that may have bitten the patient. All persons employed to fumigate rooms should be immune to yellow fever.

METHOD OF KILLING INFECTED OR NON-INFECTED MOSQUITOES.

All rooms or buildings must be tightly sealed before commencing fumigation. This work should be done

with care so as not to disturb the mosquitoes in the rooms, to the effect that they might escape. Men employed in fumigating work can secure temporary protection against the bites of infected mosquitoes, by rubbing exposed surfaces of skin with oil of citronella, pennyroyal, or 5 per cent. menthol ointment.

Sulphur.—This is the quickest and most effective agent to kill mosquitoes in rooms, destroying even hibernating forms; but its use is objectionable because it tarnishes metals and bleaches fabrics. From one to four pounds of sulphur to one thousand cubic feet of space should be burnt. A small amount of alcohol is added to the sulphur, and, after igniting, everything should be closed tight for a period of from one and one-half to four hours. Pans of water should be placed in the room.

Datura Stramonium.—The powdered leaves and stems of the Jimson weed mixed with saltpeter and burnt, kills hibernating mosquitoes. The proportion used is one ounce of the mixture to each two hundred cubic feet of space. It is the only cheap, safe and effective material for the purpose and was worked out by Mr. George Keller of Newark, N. J., for the New Jersey Mosquito Commission.

Pyrethrum Powder.—Use from one to six pounds for each one thousand cubic feet of space, and expose from three to six hours. This has been found useful, but does not certainly kill mosquitoes, but stupifies them so that they drop to the floor. When the rooms are opened up, the insects must be swept up and burned at once. It is said to be ineffective against hibernating forms.

Formaldehyde Gas.—This is effective against active summer mosquitoes if rooms are kept closed and tightly sealed for two or three hours. Generally use one liter of formalin to one thousand cubic feet of space. It is said to be ineffective against hibernating forms.

TO DESTROY MOSQUITOES IN THEIR EARLY STAGES.

It is while the insects are in the aquatic form that successful extermination can be made, and especially while they are in the larval stage. On account of the difference in the breeding places of the yellow fever and malarial bearing mosquitoes it is best to consider them separately.

YELLOW FEVER MOSQUITO.

This insect prefers to breed in water, in rain-water barrels, cisterns, tubs, cans, bottles or any receptacle containing water. This does not necessarily have to be rain-water.

City or state laws should be in force during the breeding season making it a penalty for any one to be found harboring larvæ, commonly called wrigglers or wiggle tails, around their premises. No water should be allowed to stand in any receptacle, unless it is covered, so that no mosquitoes can enter. The sanitary inspectors should receive special instructions to see that the laws are complied with. They should be provided with petroleum distributing cans, to pour oil on all uncovered, still or slowly moving water that they may find. Generally use about two teaspoonsful to the barrel, or one ounce to each fifteen square feet of water. Rain-water barrels and cisterns should be abolished by furnishing a cheap water supply. However, in localities where receptacles for water are absolutely necessary, make it compulsory that they be mosquito-proof. A strong, tight-fitting lid can be made, having a small space in the center about six inches square. This space is to be covered with fine wire gauze to prevent mosqui-

toes from entering, and still allow enough ventilation for the water. Surface drainage should be abolished wherever possible. Ditches and open drains should be cleaned occasionally and thoroughly flushed. Pools or ponds that cannot be drained should be filled in with sand or other material, or stocked with fish, or treated once a week with petroleum.

ANOPHELINE OR MALARIAL BEARING MOSQUITO.

This insect breeds in water in puddles, in hollows made by the hoofs of cattle, in pools, ditches, sluggish streams and especially in swamps. In order to prevent such accumulations of water, make superficial drains through swamp lands, or eliminate the water by filling in with sand or earth. Remove the grass and weeds from the edges of pools, streams and ditches. See that irrigation is conducted under restrictions. Pour oil over surface of water where it is practical; this forms a thin film over the surface which kills the larvæ and pupæ by preventing them from breathing. Renew the application once a week. It is doubtful whether the use of oil is successful in the treatment of swamp lands, and it is better to have these filled in or drained.

LARVICIDES.

The New Jersey Mosquito Commission have experimented with various larvicides, with the following results:

Crude Petroleum.—This may be used to kill larvæ as well as adults that attempt to lay eggs. It is useful on salt and fresh water but does not spread rapidly.

Kerosene.—This has the same effect as crude petroleum, but spreads better.

Fuel Oil.—This is like kerosene, but cheaper, somewhat heavier, and more lasting. It spreads equally well on salt or fresh water. It is the most effective mineral oil.

Chloronaphtholeum.—This is a creosol preparation, mixes readily with fresh water and kills mosquito larvæ in proportions of 1 to 1,000. Good for gutters and lot pools. Useless in salt water.

Phinotas Oil.—This is a creosol combination. It is the most effective larvicide known. Kills everything from larvæ to fish. Especially useful in sewer basins, gutters, cesspools, etc. Does not work with salt water.

THE ELEMENTS OF DIAGNOSIS OF CUTANEOUS SYPHILIS.*

E. A. FISCHKIN, M.D.,

Adjunct Professor of Dermatology, College of Physicians and Surgeons; Attending Physician (X-Ray Department), Cook County Hospital; Dermatologist, Norwegian Lutheran Deaconess Hospital.
CHICAGO.

The difficulties in the diagnosis of syphilis have been admitted by the most skillful diagnosticians. The reason for this is that the cutaneous manifestations of syphilis may be imitated by quite a number of other less obnoxious skin diseases and that syphilis may often appear in forms which simulate the lesions of other dermatoses. As Dr. White says, the skin, by its comparatively simple structure, is incapable of displaying a large variety of pathologic manifestations, and the limited number of skin lesions are therefore reappearing in a large number of morbid conditions.

No less an authority than Kaposi says:

The syphilids do not offer any other morphologic prop-

erties than the non-syphilitic dermatoses, as they all appear in macules, papules, pustules with secondary formation of scales and crusts. Their unquestioned specific clinical features, by which they can be distinguished from all other non-syphilitic dermatoses, and which impress us as peculiar typical syphilitic symptoms, are therefore not based on their morphologic properties, neither do they depend, as it is often taught, on certain physical properties, as the sepia (copper) brown color, their predominant localization on the flexor surfaces and around the orifices of the body, their symmetrical distribution, circinate appearance, formation of groups, their polymorphism, excessive formation of crusts and scales and absence of itching. For all these external features are encountered also in the non-syphilitic eruptions.

The specific properties of the syphilids are said to be the sum total of the phenomena which compose the pathologic-anatomic course of the individual lesion of which we distinguish three cardinal anatomic features:

1. The products of cutaneous syphilis are all equally composed of a uniform cellular infiltration of the papillary body and the corium and vary only in size.

2. The cells do not show a tendency to form a lasting organization into connective tissue, but undergo retrogression and complete involution.

3. The growth and consummation of the infiltrate follow in certain order and direction. They spread and disappear centrifugally.

But the peculiar specific clinical characteristics of the lesions of syphilis, if determined by these pathologic-anatomic features, are not pathognomonic of syphilis exclusively; they are the clinical expression of cellular infiltration of the papillary body in general and are everywhere met where there is such infiltration.

1. The dense sharply defined cellular infiltration of the corium and papillary layer which is manifested clinically by a papular elevation is not the exclusive feature of secondary syphilis. It is encountered in a large number of other skin diseases, such as chronic eczema, psoriasis, lichen planus, herpes tonsurans, pityriasis rubra, lupus erythematosus, etc.

2. They all show the essential feature that they do not form lasting organizations. They are distinguished by these very same features, which permit a *restitutio ad integrum* from the granulomata, in which proliferating cells undergo further development and formation of new tissue (tuberculosis, leprosy, syphilitic gummata, etc.).

3. The centrifugal spread of the infiltrate is not the expression of the pathologic process, but is determined by the anatomic structure of the skin. It is not the morphologic expression of a specific disease, but is encountered in all morbid processes in the vascular cutis, as, for instance, the different forms of erythema, psoriasis, scleroderma, pityriasis rubra, etc. All diseases, the pathologic processes of which are confined to the papillary layer or around the blood vessels, are marked by a circular spread, due to the fact that the blood vessels perforate the tissue in all directions which permit a free, that is, a circular spread of the pathologic process.

If these three cardinal symptoms are not characteristic for the pathologic process of syphilis, neither can the clinical phenomena derived from them be regarded as peculiar to the syphilitic infiltrate only. In fact, all the characteristic symptoms which have been ascribed to the papules of syphilis can be applied as well to the papules of lichen planus.

If we analyze the characteristic clinical features of the syphilids, we see that none of them is conditioned by the specific process, and that they all have their

* Read before the Chicago Medical Society.