the family history to throw any light upon the case. The past history was negative, save for the fact that the patient had suffered for several years from a small incarcerated umbilical hernia.

The present illness began June 24th, and was ushered in by a severe pain in the mass at the umbilicus. This pain persisted until June 26th, when it suddenly ceased. During this time the patient vomited nearly twenty times, but whether the vomiting was faecal, was not known. On the evening of the 25th the vomiting ceased and there were several evacuations of the bowels following a dose of citrate of magnesia. On June 26th a redness and tenderness appeared about the umbilicus and the hernial mass disappeared. On June 27th the area of redness had extended, and was very much more tender.

The physical examination showed a well developed, and rather stout woman, about fifty-two years of age. The examination was generally negative except for the abdomen. The abdomen was somewhat obese and was not distended but lax and tympanitic. There was no protrusion at the umbilicus but the depression, thereof, was unusually large and surrounded by an area of redness and tenderness and induration about six inches in diameter.

The patient entered St. Margaret's Hospital on the evening of June 27th, and was operated on the following morning under ether. As the greatest redness was below the umbilicus a vertical incision about three inches long was made below the umbilicus, and at a depth of 1 1/2 inches in the fat about two ounces of foul smelling pus was evacuated. The pus had an odor characteristic of an appendix abscess. After the excess pus had been wiped out, there appeared in the upper end of the wound a red mass about the size of a lemon.

On enlarging the wound this appeared to be the sac of a hernia with a rent about 1-2 inch in diameter. In the left lower border protruding from this rent was a sharp, hard, foreign body. The rent in the sac was slightly enlarged upward and a finger being introduced cautiously, came in contact with a loop of small intestine. The foreign body was seized in a hemostat and withdrawn and found to be a piece of animal bone, 1 7/8 inches long and 3/8 inches wide, as per illustration. Further inspection revealed a piece of small intestine strangulated in the sac, reddened but not gangrenous and showing a perforation of 1/4 inch in diameter, with its mucous membrane everted. The perforation in the gut was closed with interrupted plain cat gut sutures, in two layers. In the presence of so much infection it was not deemed advisable or safe to attempt a radical cure of the hernia, but the abdominal ring was enlarged slightly to relieve the strangulation and no effort was made to return the bowel into the abdominal cavity. Cigarette drains were placed around the sac and a large dressing applied.

The patient made an uneventful convalescence, save for a slight faecal discharge which entirely ceased by July 9th. On July 12th the patient was discharged from the hospital, the wound being clean and granulating and at the upper angle showing the mass of the sac almost on the level with the skin. On July 19th the patient's husband reported that the wound was almost healed and that she was out of doors taking a little exercise.

A report from the patient's physician, a few weeks ago, shows her to be perfectly well.

This case is unique almost, in that there was a perforation of a strangulated hernia by a foreign body. This foreign body was probably a piece of animal bone, looking more like a piece of spare rib. The patient had no recollection of swallowing any foreign body or of having any difficulty in swallowing at any time, yet it would seem almost impossible for her to swallow a piece of bone of such size without much trouble.

Another remarkable feature of the case was that the bowel practically disappeared within the abdominal cavity and in spite of the presence of so much pus, produced no general peritonitis.

RINGWORM OF THE SCALP AND ALOPECIA AREATA APPEARING SIMULTANEOUSLY IN THE SAME LOCATION.

By JOHN E. LANE, M.D., NEW HAVEN, CONN.

THE following case is of some interest on account of its possible bearing on the explanation of the cause of some of the epidemics of alopecia areata, which are occasionally reported.

I. C., an Italian boy, five years old, was seen for the first time at the Dermatological Department of the Yale University Clinic, on March 20, 1916. At the center of the nape was a bald spot, of about five centimeters in diameter, with a slightly irregular border. The skin was smooth, white and completely depilated, with the typical appearance of alopecia areata in one of its favorite locations. A more attentive examination disclosed the fact that there was considerable scalliness of the border on the side of the spot towards the scalp. Some of the scales were removed and examined microscopically. In them were found large numbers of typical ringworm fungi.

The question then presented itself, whether this was a case of the not very infrequent "bald ringworm" (Tinea) or "tinea decalvans" (Tillbury Fox), or whether it was a spot of alopecia areata

with a concomitant ringworm infection. I was unable to settle this point to my own satisfaction at this visit, but it was decided somewhat later by the course of the disease.

For a short time the spot increased a little in size and the scaly border and fungi disappeared with very little treatment in the course of a couple of months, and the spot began to be covered with hair again. This rapid cure of the ringworm without the development of any new areas, strongly suggested that the depilation which had taken place was due to the alopecia areata, inasmuch as it was as efficient in curing the ringworm as an application of a suitable dose of the x-ray would have been. Complete confirmation of this diagnosis came about a month later, when a new spot of typical alopecia areata appeared a little to the right of the location of the first spot.

Book Reviews.

The Essentials of Chemical Physiology. For the use of students. By W. D. HALLIBURTON, M.D., LL.D., F.R.S., Fellow of the Royal College of Physicians, Professor of Physiology in King's College, London; author of "Text-Book of Chemical Physiology and Pathology. Ninth edition, with colored plate. London and New York: Longmans, Green & Co. 1916.

To those of us who received our first instruction in this little book, a new edition, including all of the advances in biochemistry, is extremely welcome. Holding a place between a quiz-compend and the larger and more extensive works on this subject, it fills a very active need of students.

The experiments to be performed are stated in very simple language, are not complex, and so complete that even a beginner without other knowledge than that of the use of chemical apparatus ought to be able to conduct them successfully. Organic chemistry, too, in so far as it is necessary for the understanding of biochemical processes, is quite fully treated. The aldehydes and ketones, the ethers, the amino-acids and aromatic compounds so commonly occurring in physiological processes are fully considered and clearly described.

The chapter devoted to fats contains all the newer information which we have acquired by investigations regarding the lipoids, galactosides and phosphatides.

The classification of proteins, which the author refers to as simply provisional, because we are daily learning more and more with reference to their structure, follows that adopted in the larger works. A very valuable chapter on foodstuffs is incorporated in the volume, which

contains a brief but accurate report on the progress of our knowledge of vitamins.

In a work, however, which is supposed to be devoted to biochemistry, the description and plates of glands producing the different secretions seem out of place and indicate that the author is primarily a physiologist.

The chapter on blood is rather brief and contains practically none of the newer investigations which have been conducted on the importance of variations of the amount of urea, uric acid, sugar, and creatin, which are revolutionizing our methods of testing for pathological changes in the body by means of the blood.

On the whole, we may regard this work as playing a very important part in the instruction of students, but rather too brief for the needs of anyone engaged in a thorough prosecution of any line of investigation.

Diagnosis and Treatment of Surgical Diseases of the Spinal Cord and Its Membranes. By CHARLES A. ELSBERG, M.D., F.A.C.S., Professor of Clinical Surgery at the New York University and Bellevue Hospital Medical College; Attending Surgeon to Mount Sinai Hospital and to the New York Neurological Institute. With 158 illustrations, three of them in colors. Philadelphia and London: W. B. Saunders Company. 1916.

An attractive book of 300 pages, profusely and most liberally illustrated; easy to read, because of large type and wide spacing; easy to hold, because it is not too bulky; and interesting from cover to cover, because, as Dr. Elsberg says in his preface, the "volume is a record of personal experiences in the surgical treatment of diseases and injuries of the spinal cord and its adnexa," and because he has limited himself rigidly to "a consideration of the diagnosis and treatment of diseases of the spinal cord that may require surgical interference."

We have referred to the illustrations: there are no less than 150 of them; many half-tones; many excellent reproductions of x-ray plates; a number of diagrams, and two admirable colored plates illustrating the difference between the spinal veins in inflammation and the spinal veins in compression. The illustrations, as a rule, take half a page, and occasionally a whole page, so that the text is not more than 200 pages, and probably less. Emphasis is laid upon this fact, because the tendency at present seems to be to make medical books too long: either the writer over-elaborates his subject, or else he does not remain in the field which he originally set for himself. Dr. Elsberg avoids both these dangers; he speaks with highest authority and from a very wide experience; his book is unlike anything heretofore published in America, and it should be in the hands of every surgeon who may be called upon to explore the spinal cord.