joints as the result of better knowledge of the nature and distribution of the disease, the treatment of appendicitis and the investigations on peritonitis and diseases of the appendix, the more complete operations for cancer following fuller study of the mode of spread of the disease, and so on.

But while pathologists are thus working out problems which affect the general well-being of mankind and the solution of which can be no personal gain to themselves, is it too much to ask mankind to furnish the means for such research? The English are looked on as a thoroughly practical people, and yet it is a very remarkable thing that England is almost the only country which does not realize the importance of scientific research and the result is that in England, with very few exceptions, men who might otherwise have thrown much light on these matters are compelled to turn their attention to practice in order to make a living. Unless work of this kind is done how can we hope to advance with any rapidity in the treatment and cure of disease? The surgeon or physician must wait till the information of which he is urgently in need has been acquired for him by the pathologist. Such apathy can surely only be the result of ignorance. A rich man affected with an obscure or incurable malady can not understand how it is that he fails to obtain the definite opinion or the relief which he so earnestly desires and for which he is prepared to pay any price. Surely if he understood the meaning and importance of pathologic research and that the practicing physician can only apply and carry out what is taught by the pathologist, he would bestir himself to aid research in order to gather information which might be of much use to him and others.

ORIGINAL ARTICLES.

EXPERIMENTAL BASIS OF THE DIETETIC AND MEDICINAL TREATMENT OF HYPERACIDITY AND GASTRITIS.

Presented to the Section on Materia Medica, Pharmacy and Therapeutics, at the Forty-eighth Annual Meeting of the American Medical Association, held at Philadelphia, Pa., June 1-4, 1897.

BY JOHN C. HEMMETER, M.B., M.D., PH.D.

CLINICAL PROFESSOR OF MEDICINE IN THE BALTIMORE MEDICAL COLLEGE; CONSULTANT TO THE MARYLAND GENERAL HOSPITAL. BALTIMORE, MD.

Boas, who first used the finding of fragments of mucosa in the vomited matter and gastric wash water for diagnostic purposes, attributed great importance to this way of finding out the real state of the mucosa. He held that, in certain conditions of suppressed secretion, the differential diagnosis between a possible neurosis and a genuine gastritis with a glandular atrophy was only possible by examination of such pieces of mucosa. Rosenheim, Boas and Julius Friedenwald, *Med. News*, June 22, 1895, emphasize the value of qualitative and quantitative testing of rennet zymogen to differentiate between chronic gastritis with glandular atrophy and carcinoma on the one hand, and nervous dyspepsia and secondary gastritis on the other. However, Ewald and also Einhorn have asserted that absolute deficiency of rennet zymogen is not pathognostic for atrophy, therefore it would indeed seem as if a certain diagnosis could only be made by a small piece of mucosa.

Is there any clue which can be derived from these

pieces regarding the state of the mucosa in the secretory disorders? This I will try to answer in the following. Hayem, to whom we are indebted for the best histologic investigations of the gastric mucosa, emphasizes that the individual elements of the mucosa, gland ducts, superficial epithelium and interstitial tissue can become diseased in a variety of ways; the various portions of the stomach fundus, pylorus and cardia may exhibit different affections; and finally the mucosa may at different parts show different phases of disease. He distinguishes a parenchymatous and an interstitial gastritis. First the parenchymatous:

1. *Gastrite parenchymateuse hyperpeptique chloro-organique*.—Under this he has two subclasses: *a*, *d'emblée*—dyspepsia coming on at once, in the first stage of digestion; *b*, *tardive*—coming on in later stages, in one and one-half to two hours. Under this hyperpeptic parenchymatous gastritis Hayem means, clinically, a hyperpepsia with hyperacidity, and anatomically, degeneration of the principal central or chief cells with proliferation of the parietal border or oxyntic cells.

2. *Gastrite parenchymateuse muqueuse*—"gastritis mucipara," by which he means a mucous degeneration, a process taking place principally in the vestibules to the gland ducts (which are lined with columnar epithelium), and corresponds to the Schleim Katarrh of most German writers. This is associated with hyperpepsia and subacidity.

3. *Gastrite parenchymateuse atrophique*, which signifies anatomically the total atrophy of the glands without interstitial processes, and clinically, anacidity or achylia. The interstitial forms he separates into two classes: those in which the round cell infiltration, and those in which the sclerosis, *i. e.*, connective tissue proliferation, predominates. These processes are described as occurring purely as such, or mixed with forms of parenchymatous gastritis, and as leading to sub- or anacidity. In order to bring my results in critical consideration with those of Einhorn, I have adopted his classification of the anatomic conditions found in these fragments. There is, however, one objection that can be urged against it, and that is the apparent fact that he has based his system on conditions of the gland tubes and interglandular tissues exclusively, and mentions the state of the cells only once in six types described. I will therefore supplement his categories by adding the state and condition of the vestibular or alveolar columnar cells (*Vorraum Zellen*), and the condition and numeric relations of the chief, central or ferment cells (*Hauptzellen*) and the parietal, border or oxyntic cells (*Belegzellen*).

1. Normal. Gland ducts and interglandular tissues exist in normal proportions. Columnar epithelium of the surface and that of vestibule normal, with scarce cells showing at their free ends slight mucoid metamorphosis. The average number of parietal or oxyntic cells in six ducts which were sectioned very nearly down the center was twenty-two to forty.

2. Connective tissue excess. Proliferation of connective tissue around the glands, glands and epithelial cells as in normal condition.

3. Proliferation of glands. Under this class I have in nineteen cases been impressed with the probability that there must be three types of this condition:

Type *a*.—In this subtype there is a proliferation of gland tubules. Under the same field of microscope there will be more than under normal conditions, since they are much closer to each other, but the number

of central and oxyntic cells are from eighteen to forty-two, or the same as under the normal condition.

Type *b*.—Increase of oxyntic or parietal cells with normal number of gland ducts. Here there seems to be no proliferation of the gland ducts, the connective tissue and the ducts bear the same relation as in class 1, but the anilin staining, oxyntic cells may be so increased that they lie in juxtaposition, giving the whole duct the appearance of a peptic duct of the dog; the number may reach seventy in one duct. The oxyntic cells are increased in size; the nuclei stain very dark.

Type *c*.—Increase of the number of ducts in which the number of oxyntic cells appear normal in size and number, and in the same fragment or section portions of mucosa in which the ducts are not augmented, but the oxyntic cells are increased in number and size; this third is then, it would seem, a combination of types *a* and *b*. When there are many oxyntic cells above the normal, the entire gland duct assumes a tortuous or elongated shape. It seldom extends down into the mucosa in the same plane, therefore it is very rare that a section will strike down the middle of a duct. Generally the counts in six ducts struck fairly along the central canaliculus are taken as an average.

4. Incipient atrophy. To the same field under the micrometer there are fewer glands than normally present. They appear shrunken and smaller, and at the same time the spaces between the glands are larger than normal, owing to an increased connective tissue formation; the latter is thickly invaded as a rule, with small round cell infiltration.

5. Atrophy. In complete atrophy there are only remnants of glands left, a few degenerated cells lying in empty circular spaces where glands had previously existed; there is also a diffuse round-celled infiltration.

6. Vacuolization. Round or ovoid vacuoles exist within the glands in large numbers, being the result of mucoid degeneration of some of the glandular cells; this is generally associated with connective tissue proliferation. Vacuoles are present in the gland cells normally and can be seen in the drawings of Kupfer and Stöhr. I have also seen them in both longitudinal and cross sections of the gland tubules, but rarely more than two to three to the entire duct. It is conceivable that they may be produced by the process of hardening and imbedding. Some of the fragments obtained from stomachs may show characteristics of two types.

DEDUCTIONS FROM THIRTY-SIX CASES.

In eight *healthy* persons the mucosa fragments were normal in six; proliferation and autodigestion marked in one, which also showed beginning small round cell infiltration between the ducts; connective tissue increase in one. In the first of these cases the examination showed proliferation in one fragment and a normal condition in a second one found in the same wash water.

In eighteen cases of *hyperacidity*, the fragments of gastric mucosa were found apparently normal in four; atrophy of gland tubules and connective tissue increase so that there were fewer glands, but in these few there were contained a larger number of oxyntic cells than normal, in two cases; proliferation of gland ducts with apparently normal oxyntic cells in six cases; proliferation of oxyntic cells, generally without marked increase in the gland tubules, in six cases.

In twelve cases of *anacidity* or *subacidity* the frag-

ment was apparently normal in two cases; proliferation of glands with marked small round cell infiltration was found once. Atrophy in some form was found in the fragments from the nine remaining cases.

In establishing the classification of *euchlorhydria* and *hyperchlorhydria* we could not be guided exclusively by the amount of free HCl found after the double test meal used at the Maryland General Hospital and described in the author's book (Hemmeter, "Diseases of the Stomach," p. 111. P. Blakiston Son & Co., Philadelphia, 1897). Thus, a young, vigorous farmer, aged 25, who had never had any disease, showed on repeated examination an amount of free HCl equal to 60 degrees, with a total acidity of 80 degrees. Ordinarily, judging simply from the analysis, such a case would be diagnosed as hyperacidity. However, these cases can be diagnosed justly and accurately when considered together with concomitant signs and symptoms only. Although this case had the large amount of free HCl, there was no starch indigestion, no erythrodextrin, no pyrosis; there were no symptoms referable to the stomach at all, the man was in perfect health.

Another case, a neurasthenic female, had intense suffering from hyperacidity and occasional gastrodynia, and the amount of free HCl was never over 30 degrees. This case showed hypermotility. The stomach as a rule was empty twenty-five minutes after an Ewald test meal; with my intragastric rubber bag in connection with the kymograph, she showed very frequent and sudden gastric peristalsis of unusual tonicity.

SUMMARY.

Eight *healthy* persons: Perfectly normal in six; *a*. glandular proliferations; *b*. normal in 1; connective tissue increase in 1.

Hyperacidity in eighteen cases: Normal in four; atrophy in two; proliferation of glands in six; proliferation or hypertrophy of oxyntic cells in six.

Anacidity or *subacidity*: Normal in two; proliferation of glands in one; atrophy in nine.

Proliferation, therefore, according to this table is present in two-thirds of these cases of hyperacidity, and atrophy in three-fourths of these cases of anacidity or subacidity. Einhorn does not give any results from examination of perfectly healthy individuals as his cases of *euchlorhydria* seem to be in patients.

Of his twelve hyperacid cases three were normal or very nearly so, six showed proliferation and three showed connective tissue proliferation. In his cases of anacidity, or rather what he calls *achylia gastrica* of which there were seven cases, there was atrophy three times, marked vacuolization once, proliferation once, and normal condition twice.

On the whole, judging from Einhorn's results, Cohnheim's, Hayem's and my own, the conclusions seem justifiable that proliferation of glandular elements is present in from one-half to two-thirds of the cases of hyperacidity, and atrophy is present in from one-half to two-thirds of the cases of anacidity.

Adolf Schmidt (*Virchow's Archiv*, Bd. cxliii, S. 478), asserts that the epithelium of the stomach is preserved better than the gland cells in inflammatory conditions of the mucosa. This he says is particularly so in chronic gastritis which forms island-like foci in stomachs otherwise not much changed. My experience and that of W. D. Booker, is not in accordance with this observation (see pathology of simple, acute and chronic gastritis in the clinical portion of Hemme-

ter's "Diseases of the Stomach"). Although I preserved the stomachs by injecting them with alcohol immediately after death (within twenty minutes), also with formalin and sublimate so that autodigestion was at once checked, my sections showed generally a more serious destruction of the surface epithelium than of the gland cells. At times both are so much altered

cardia or esophagus they are most frequently found in the lower or side opening of the tube, as it must pass through or over the growth on its way into the stomach; but even in malignant growths of other parts of the stomach, patient searching in the sediment of the wash water will sometimes reward the clinician by the discovery of tumor fragments. The first wash water

Fragment of mucosa showing normal condition of glands; very slight round-celled infiltration. From Hemmeter's "Diseases of the Stomach," P. Blakiston Son & Co., Philadelphia, Pa., October, 1897.

Atrophy and vacuolization of glandular elements; mucoid degeneration of chief or central cells; comparative preservation of border or oxyntic cells; from a case of gastritis acida. From Hemmeter's "Diseases of the Stomach," P. Blakiston Son & Co., Philadelphia, Pa.

that it is impossible to say which is most or least affected. It seems in chronic gastritis that new epithelium will be reformed quite rapidly where the old has been lost or destroyed.

In suspected cases of malignant neoplasms fragments of the growth are occasionally found and are of importance in the diagnosis. In carcinoma of the

in the morning, about 500 c.c., should be permitted to settle twelve hours in a conical glass such as is used for the settling of urinary solid constituents, and the sediment should be examined under a low power (about 50 diam.) or the sediment may be more expediently brought down in the centrifuge.

Once I made the diagnosis of carcinoma when no

tumor was evident from repeatedly finding involuntary muscle fibers when no meat had been eaten for three days after preceding lavage. It proved to be a broad, flat carcinoma of the posterior wall.

The sections were stained in a variety of ways, principally in the eosin-hematoxylin, Golgi and Bismarck brown stains. The minute communications of the oxyntic or parietal cells with the central duct are best brought out by the Golgi method.

The drawings of fragments found in the wash water of glandular proliferation with glands closely packed and connective tissue diminished, and of glandular atrophy, mucoid degeneration, vacuolization, and small cell infiltration are all explained by the text accompanying the illustrations. These drawings are taken from the author's text-book, "Diseases of the Stomach." We have seen that histologic changes approaching or actually representing pathologic states may be going on in perfectly healthy stomachs. Furthermore, the stomachs of diseased patients may on serial sections show a different pathologic state at different places of

cell, hyperplasia and hypertrophy. Edinger explains the anatomic changes in the spinal cord and nerves by a disproportion of the function of activity and the replacement of nervous elements; an increased activity of an organ leads to straightening of the same and increase in its volume; inactivity leads to a weakening and atrophic state of an organ. W. Roux ("Entwicklungs-Mechanick der Organismen," 1895) uses the expression "each cell gains a living by work, and the more it works the better it is nourished and the stronger it grows." In hyperacidity the elements that come more into play and do more work will increase in strength and numbers and gain supremacy. Dujardin-Baumetz and F. Sohlern (*Berlin. Klin. Wochschr.*, 1891, 20 and 21); Fleiner (Volkman's *Klin. Vortr.*, 103) and Rummo (*Terapia clin.*, 1892, Nos. 10, 11 and 12), recommended a largely carbohydrate diet in gastric diseases accompanied by abnormal production of acid. We have satisfied ourselves personally, after a very large number of quantitative chemic analyses of the gastric contents of healthy

Hypertrophy and proliferation of glandular elements, from a case of persistent hyperacidity.

the mucosa. Therefore, it must be borne in mind that although the findings in hyperacidity and anacidity appear to be in some relation to the disease, this kind of investigation must not be relied on as representing in a given fragment the condition of the entire mucosa. It represents the state of the location from whence it is sequestered, and that not being accurately known generalizations must be made with caution.

It is therefore rational to presume that there are at least two kinds of hyperchylia: One which is of neurotic origin and in which the acid elements of the mucosa are not hypertrophied or augmented, and the other in which the neurotic symptoms are absent, or at least not very marked, and in which a decided proliferation of the secreting cells of the gland ducts can be made out. It is conceivable also that in the neurotic type, although there may be no hypertrophy and increase in the number of gland cells at the beginning of the disease, the constant demand upon and stimulation to the increase of acid and ferments may, in the course of the disease, eventually bring about an oxyntic

individuals after they had taken weighed amounts of pure proteid and pure carbohydrates, that the proteid articles of diet are greater stimulants to the secretion of HCl than the carbohydrates or fats. They therefore stimulate and irritate the secretory layer to a greater formation of hydrochloric acid; this has been experimentally proven to a certain extent by v. Jacksch in children. It is also known that the great variations stated by different authors as to the amount of free HCl that is normal in their localities are explicable by the nature of the various test-meals used by them. The amount of free HCl after the complex test-meal of Riegel and Fleiner is much greater than after a simple test breakfast which contains very little proteid. It must not be overlooked that an exclusive proteid diet causes the formation of excessively large quantities of soluble peptones and albumoses, which undoubtedly have an exciting action on the nervous system and constitute a favorable basis for the development of all kinds of neuroses. Although a diet rich in meat food is perfectly logical and cor-

rect in hyperacidity, careful quantitative analyses of the urine will show that the indican, the ethereal and preformed sulphates are very much more increased under this kind of diet, and that they become reduced as soon as the amount of carbohydrates is increased. Beef, mutton, lamb, venison, fowl and various fishes, undoubtedly relieve the main complaints of hyperacidity for a time. The gastric pain and the pyrosis disappear for a time at least, because the albumin contained in the substances combines with the excess of HCl, but in cases in which a lasting improvement is not affected by these articles of diet, and where the urotoxic coefficient, the amount of indican and aromatic sulphates become increased it is advisable to try a diet in which the proteids do not constitute the preponderance, but at least 50 per cent. of which is composed of fats and carbohydrates.

The following tables show the state of the urine in a case of hyperacidity, both under proteid and under a carbohydrate diet. The case suffered as a rule from auto-intoxication when the proteid food was in excess.

URINARY ANALYSIS.

Case 19.—April 12, 1896. W. M., male, age 35 years, hyperchylia, on proteid diet exclusively; no medicine. Free HCl=48.5 after Ewald test breakfast.

Indigo blue	very strong
Urea	22.712 grams
Uric acid	0.451 grams
Ratio	49.0
Preformed sulphates	1.826 grams
Combined sulphates	0.236 grams
Ratio	7.7

The combined ethereal sulphates and the indigo are present here in great excess; headache and much flatulence.

Case 19.—May 12, 1896. W. M., hyperchylia, Free HCl=59. Mixed diet, containing rice, bread, oatmeal, together with small quantities of beef or soft boiled eggs, and milk and butter. Neutralization with alkalies and aiding the gastric amylolysis with ptyalin at one meal and taka diastase at another.

Indigo blue	very faint
Urea	42.641 grams
Preformed sulphates	4.000 grams
Combined sulphates	0.350 grams
Ratio	11.0

Patient feeling much better, less headache, debility and flatulence. This is an example of twenty analyses, showing that even with an excess of HCl the proteid putrefaction may be considerable and that the symptoms are relieved and the urine shows less of toxins on a mixed diet with treatment by ferments and alkalies.

These are only two examples of a series of eighteen analyses.

It is undeniable that carbohydrates are found undigested six hours after they are ingested, in the wash water from the stomach. This is what one should expect, since if the excess of hydrochloric acid is not neutralized by alkalies, the ptyalin can not act and it is impossible for carbohydrates to be converted into a soluble form, the HCl at the same time producing a pyloric spasm obstructing the passage into the duodenum. Even with a diet almost exclusively of a proteid nature, the alkalies are indispensable, as this food is in itself inadequate to combine with the excess of acid. We advise our patients to take their amylaceous foods in a very finely divided state when possible. Potatoes should only be allowed in form of purée. Beets, turnips and carrots should not be allowed at all. The artificially prepared soup meals (Kufek's, Maggi's flour, Malto-leguminose) and soups made from aleuronat, and oatmeal flour are in these cases easily digested with the aid of artificial diastase or ptyalin. It is a very important question whether the diastatic ferment of the saliva is destroyed permanently or only temporarily

by the excess of hydrochloric acid and whether after its action has been inhibited by the acid it can be restored by subsequent alkalization. Boas ("Diagnostik u. Therap. d. Magenkrankh.," p. 19) has demonstrated that when saliva is exposed to the action of 0.15 per cent. HCl for one hour its action can be restored by alkalization with bicarbonate of soda. In hyperacidity the percentage of acid reached is 0.2 per thousand of HCl; in severe cases 0.4 to 0.6 per thousand HCl. We have made a large number of experiments with the effects of hydrochloric acid on saliva, both with artificial test-tube experiments as well as experiments from stomach contents of patients suffering with hyperacidity, and have found that, although Boas's results are in the main true, the action of ptyalin after it has been exposed to an acidity of 0.3 per thousand for forty minutes can not be restored to the same efficiency it showed before it was subjected to the action of the acid; that is, alkalization restores the action only partially. The conversion of starches by this ptyalin which has been thus detrimentally influenced by HCl takes place only very feebly, in other words, the ptyalin does not recover perfectly, it has been permanently damaged. These experiments have been repeated so often by us and the results have been so constant that they could not possibly be due to accident. They justify us in the conclusion that the internal administration of alkalies in hyperacidity will not restore gastric amylolysis although the distressing symptoms of hyperchylia will be relieved by it. It is therefore necessary, if we wish to aid digestion of starchy foods and not simply relieve the distressing symptoms of hyperacidity, not to depend on restoration of the function of ptyalin by giving alkalies. The employment of artificial ptyalin or diastase is here unavoidable; ptyalin, one-half to one gram (8-15 grains) given fifteen minutes before meals together with alkalies, effectively converts the carbohydrates into dextrose. The more carbohydrate is artificially digested in this way, the less undigested starch will naturally remain in the stomach. We will naturally, therefore, give the preference to a ferment which can in the shortest time convert the greatest amount of carbohydrate into dextrose. We have made a number of experiments with ptyalin and the diastase contained in malt extract, but the most effective agent that we have worked with is a ferment known as taka-diastase. This has the property of being far less susceptible to the destructive action of hydrochloric acid than ptyalin or malt extract. The objection we have to malt extract is that we have to give it in a large bulk to get a very small quantity of diastase. The extract itself contains fermentable sugars which, in some cases of indigestion, may give rise to fermentation in the stomach. Taka-diastase is a substance obtained from a fungus known as *euotium oryzae*, which is sown on moistened sterilized wheat bran. A growth results which is found, under the microscope, to present peculiar crystals of pure diastase covering the roots of the *euotium* which have penetrated the bran. The function of these crystals is to convert the starch of the bran for the nourishment of the fungus. We have employed taka-diastase in hyperacidity and in these rare cases of deficiency of saliva, which are found in the final stages of chronic Bright's disease, and after severe diarrheas. Malt diastase and ptyalin cease acting in the stomach contents much sooner than taka-diastase with a percentage of HCl equal to 0.04 per thousand to 0.05 per thousand of HCl, taka-

diastase will still transform more than 50 per cent. of starch into maltose. The following experiments will throw some light on the digestive power of taka-diastase. A starch paste was made from 5 grams of dry arrow-root starch with 500 c.c. of water. Mixtures were then made as follows:

1. 0.5 gram of taka-diastase + 90 c.c. of water + 10 c.c. of starch paste.
2. 0.5 gram of ptyalin + 90 c.c. of water + 10 c.c. of starch paste.
3. 20 gram of malt diastase + 90 c.c. of water + 10 c.c. of starch paste.

These three mixtures were placed at 40 degrees C. and tested from time to time with iodine solution. In periods of time varying from five to eight minutes No. 1 had reached the achromic point, while No. 2 had not reached the achromic point until one hour had elapsed. At the end of three hours No. 3 still gave a bluish violet reaction with iodine. In the presence of 0.015 HCl the action of the ptyalin and the malt extract were arrested, whereas the action of the taka-diastase continued unhindered. We therefore give taka-diastase because of its more effective and permanent amylolytic power, because of its cheapness, and because we are enabled to permit a diet rich in carbohydrate in cases of hyperacidity. The most essential thing in the treatment of an irritable gastric mucosa is rest for a short time, two or three days in every week; no food but milk or thin oatmeal gruel. To give meat and egg with a view to combine with the excess of HCl is only symptomatic treatment. But an amylaceous diet aims at the cause of the hyperacidity, because it is not a stimulant to HCl secretion and does not add to the irritation already present.

In recommending a carbohydrate diet in hyperacidity we are aware that it will generally not be tried until the proteid diet has failed to give permanent relief. The amylaceous diet is to a certain extent antiseptic by its tendency to keep the intestinal chyme acid by formation of lactic acid. The quantitative analysis of indican and ethereal sulphates will keep one posted on the degree of intestinal putrefaction under the two kinds of diet.

It is well known that hyperchylia often occurs in neurasthenia due to uric acid diathesis. The diet must then naturally be adapted as far as is consistent with the gastric function to the uric acid diathesis underlying the whole complication of symptoms (see *Diät bei Gicht und Harnsäure dyskrasie*, Penzoldt u. Stintzing, Vol. ii).

BIBLIOGRAPHY.

- 1 Cohnheim, Paul: D. Bedeut. klein. Schleimhaut-Stückchen f. d. Diagn. d. Magenkrankh., *Archiv f. Verdauungs Kr.*, Bd. i, S. 274.
- 2 Schmidt, Adolf: Untersuch. über Mensch. Magenepithel, normal u. pathol., *Virch. Arch.*, Bd. 143, xix.
- 3 Langerhans: *Virch. Arch.*, Bd. 124, p. 373.
- 4 Hartung, O.: *Deutsche Med. Woch.*, No. 38, p. 847, 1890.
- 5 Virchow, R.: *Virch. Arch.*, Bd. v, p. 363.
- 6 Gerhardt, D.: *Virch. Arch.*, Bd. 127, p. 85.
- 7 Ewald: Ein Fall v. Atrophie d. Magenschleimhaut mit Verlust d. HCl Secretion Ulcus carcinomatosum duodenale, *Berlin. klin. Woch.*, 1886.
- 8 Fenwick: Ueber den Zusammenhang einiger krankhafter Zustände des Magens mit anderen Organerkrankungen, *Virch. Arch.*, Bd. 118, xii.
- 9 Ewald: Ein Fall chronischer Secretions-unfähigkeit des Magens (*Anadenia ventriculi*?) Das Benzonaphthol, *Berl. klin. Woch.*, 26 u. 27, 1892.
- 10 Boas: Beitrag zur Symptomatologie des chronischen Magencatarrhs und der Atrophie der Magenschleimhaut, *Münch. med. Woch.*, 41 u. 42.
- 11 Einhorn: Sur Achylia Gastrica, *Arch. f. Verdauungskrankh.*, Bd. 1, Heft 2.
- 12 Schmidt: Fortgesetzte Untersuchungen über die Secretion des Magenschleims, *Deutsch. med. Woch. Vereinsbeilage*, 13, 1895.
- 13 Klemperer: Ueber die Dyspepsie der Phthisiker, *Berlin. klin. Woch.*, 11, 1889.
- 14 Boas: Ueber Gastritis acida, *Wiener med. Woch.*, 1-2, 1895.
- 15 Stöhr: Zur Kenntniss des feineren Baues des menschlichen Magenschleimhaut, *Schultze's Arch.*, Bd. xx, p. 221.
- 16 Ebstein: Beiträge zur Lehre vom Bau der sogenannten Magenschleimdrüsen, *Schultze's Arch.*, Bd. vi, p. 530.

- 17 Kupffer: Epithel u. Drüsen d. menschlichen Magens, München, 1883.
- 18 Stintzing: Zum feineren Bau u. zur Physiologie d. Magenschleimhaut, *Münchener med. Woch.*, 46, 1889.
- 19 Sachs: Zur Kenntniss der Magenschleimhaut in krankhaften Zuständen, *Arch. f. exp. Pharma. u. Pathol.*, Bd. 24, 1888.
- 20 Hammerschlag: Zur Kenntniss des Magencarcinoms, *Wiener klin. Rundschau*, 23, 1895.
- 21 Meyer: Zur Kenntniss der sogenannten Magenatrophie, *Zeitsch. f. klin. Med.*, Bd. xvi, p. 366.
- 22 Jaworski u. Korczynski: Ueber einige bisher wenig berücksichtigte klinische und anatomische Erscheinungen im Verlaufe des sogenannten Magencatarrhs, *Arch. f. klin. Med.*, 47, p. 578.
- 23 Rosenheim: Ueber atrophische Prozesse an der Magenschleimhaut in ihrer Beziehung zum Carcinom u. als selbständige Erkrankung, *Berl. klin. Woch.*, 51, 1881.
- 24 Schmidt: Einfall von Magenschleimhautatrophie nebst Bemerkungen über die sogenannte schleimige Degeneration der Drüsenzellen des Magens, *Deutsch. med. Woch.*, 19, 1895.
- 25 Damaschino: Note sur un nouveau procédé pour l'étude de lésions de l'estomac, *Gaz. méd.*, 1880.
- 26 Hayem: Résumé de l'Anatomie Pathologique de la Gastrite Chronique, *Gaz. Hebdom.*, pp. 33-34, 1892.
- 27 Boas: Diagnostik u. Therap. d. Magenkrankh., *Allg. Th.*, iii ed, p. 220.
- 28 Ewald: Klinik d. Verdauungskrankheiten, iii edit, p. 191.
- 29 Einhorn: State of the Gastric Mucosa in Secretory Disorders of the Stomach, *New York Medical Record*, June 27, 1896.
- 30 Einhorn: Clinical Observations on Erosions of the Stomach and their Treatment, *N. Y. Medical Record*, June 23, 1894.
- 31 Ebstein: Ueber die Lösung eines Stückes d. Pylorus-schleimhaut mit d. Magensonde, *Berlin. klin. Woch.*, 1895.
- 32 Korczynski u. Jaworski: Klinische Befunde bei Ulcus u. Carcinoma Ventriculi, etc., *Deutsche. Med. Woch.*, pp. 47-49, 1886.
- 33 Crümer: Ueber d. Ablösung d. Mag. Schleimhaut durch die Sondierung, *Münch. Med. Woch.*, p. 52, 1891.
- 34 Hayem: Gastritis Parenchymatosa, *Allg. Wien. Med. Zeitung*, 1894, pp. 2-17.

DISCUSSION.

Dr. C. B. LOWE of Philadelphia—I had the pleasure yesterday of meeting the Japanese chemist, Dr. Takamine, the discoverer of taka-diastase. The remarkable activity of this amylolytic ferment was demonstrated by his placing a few grains in a test tube containing starch paste. After adding it, either in the dry form or as a liquid, the solution began immediately, and in a moment or two the contents of the tube were liquefied and converted. The experiment was a very striking one and demonstrated the great power possessed by this agent in the digestion of starchy articles of food.

Dr. K. G. ECCLES of Brooklyn—Did I understand the lecturer to say that maltose would be absorbed from the stomach? I was under the impression that all starchy foods passed into the intestine where it was converted into dextrose by the pancreatic juice and the intestinal secretion. Perhaps Dr. Hemmeter will set me right on this point. I am not sure whether I fully understand his position in reference to the use of taka-diastase in cases of hyperacidity. It would seem to me that most of the starchy foods are not digested in the stomach, although the proteids are acted upon there.

Dr. F. E. STEWART of Detroit—In reference to the question of the absorption of maltose into the blood from the stomach, I would say that I am under the impression that Professor Chittenden found that it was not essential that the starch be entirely converted before being absorbed. Dr. Roberts of London found that the glycogen of the liver is very much like the same products of starch digestion.

Dr. ECCLES—I would like to add that I have tried taka-diastase on myself and on patients and have found the results from it to be very good. I therefore was not criticising the treatment but merely desired information upon the physiologic and chemic basis of the method of treatment advocated by Dr. Hemmeter.

Dr. FRANK H. MURDOCK of Pittsburg, Pa.—I have been exceedingly interested in Dr. Hemmeter's paper. I think that the question of hyperacidity of the stomach is one of very great importance and one which offers the most difficulty to me in my practice. If we can have within our reach remedies which, aided by diet, will afford the patient relief, we would certainly all be glad to know it.

Dr. HEMMETER—I must say that I am very much pleased at the discussion of this paper and at the questions which have been asked. One question was whether or not maltose is absorbed from the human stomach and enters the blood as such. I have invented a method of studying this problem by plugging the pyloric end of the stomach in animals; maltose is then introduced into the stomach. The pylorus being closed by the plug, it can not escape into the intestines. Nevertheless, shortly afterward there is no maltose left in the stomach. It is absorbed into the blood through the walls of the stomach. As Dr. Stewart has said, Sir Wm. Roberts has shown that maltose can be absorbed directly into the blood. Not only maltose, but albumose may be absorbed, and I have no doubt that egg albumin may be injected into the intestine and be absorbed directly into the blood without being digested at all. I have been asked how long amylosis will go on in the human stomach after an ordinary meal. Now it is limited by only one thing, that is, the change to an acid reaction in the

stomach contents. If the food stays long enough and there is not too much acid, the starchy constituents will be entirely converted in the stomach. In dilated stomachs, where there is destruction of the hydrochloric acid cells, the digestion of starch may go on for many hours. As soon as the proportion of hydrochloric acid reaches 0.15 then the action of starch ceases. This generally occurs on the average in about forty minutes after the close of an ordinary meal, so that we have this length of time in which to digest the starchy portions of our food in the stomach. If too much work is thrown upon the amylolytic action of the intestinal glands, there may be intestinal failure and intestinal indigestion. Because the amylase of the pancreatic juice digests starch and the ptyalin of the saliva digests starch, we must infer that their actions or end products are the same. The cases which Dr. Murdock has mentioned and in which we find alternating states of the secretion, one day hyperchylia, the next achylia, and perhaps in the next few days hyperchylia again, have been termed by me *heterochylia*. In determining the degree of acidity in these cases it is essential to know the amount of proteid in the test meal, because the secretion of HCl is greater after proteid than after amylaceous foods. The condition causing heterochylia can only be cured by correcting an underlying neurasthenia, by good hygiene, proper diet, electricity, baths, etc. Moreover, I believe that in many of these cases, lithemia and gout are at the bottom of the condition of heterochylia. In cases where this persistently exists, the patient may be taught to pass the stomach tube and wash out the stomach for himself with diluted HCl or alkalies as may be indicated, but it is impossible to go into the details of the treatment or give any sweeping advice for the management of such cases.

A STUDY OF THYROID EXTRACTS.

Presented in the Section on Materia Medica, Pharmacy and Therapeutics, at the Forty-eighth Annual Meeting of the American Medical Association at Philadelphia, Pa., June 1-4, 1897.

BY W. H. NEILSON, M.D.

MILWAUKEE, WIS.

Interest in the thyroid body is not of recent origin. As far back as 1856, Schiff demonstrated, by his experiments on dogs, that this gland is an absolute necessity for the maintenance of life in these animals and his deductions have been abundantly confirmed by many observers since that time. After the removal of the thyroid gland a number of characteristic symptoms supervenes, as cachexia, emaciation, apathy and muscular tremor which may pass into convulsions before death ensues. The muscular phenomena appear to have a central nervous origin, for section of the motor nerves prevents them; and it is possible that the trophic changes noticed are also due to perverted nervous influence.

In rabbits, Hofmeister demonstrated that by removal of the gland, leaving the glandules of Gley, life might be maintained. The hair became coarse, the body short and thick, the growth of the skeleton was arrested, the epithelium of the convoluted tubules of the kidneys was altered, and enlargement of the hypophysis cerebri occurred (Crary). In monkeys, these changes occur more slowly than in the lower animals and resemble much the condition developed in man through destruction of the function of the thyroid. The pronounced symptoms of this condition are diminished mentality, muscular weakness, marked anemia, dry harsh skin and loss of hair, with an edema of the subcutaneous tissue which does not pit on pressure. If the individual so suffering receive hypodermatic injections of thyroid extract, or if it be administered by mouth, or even the raw gland be fed him, all of these symptoms, except the anemia, rapidly disappear and he regains his normal condition with the one exception. In the case of dogs, if a small portion of the thyroid gland, after its removal, be grafted into the peritoneum, the evil results are averted. So, too, if a small portion of the gland remain or the para-

thyroids be left the animal will apparently suffer no inconvenience. Thus it becomes clear that the thyroid body is intimately associated with the normal manifestations of life. There are two views obtaining as to the manner of this association. The one is that the thyroid is a destructive agent, destroying poisonous substances formed in other organs, which if undestroyed would cause the various symptoms found in myxedema. In support of this is given as a fact by some observers, contradicted by others, that products are found in the muscles, blood and urine of animals deprived of the thyroid, which are decidedly toxic in character. The other view is that the cells of the thyroid produce a true internal secretion which finds its way into the blood and plays an important part in producing normal metabolic changes in some or all of the organs and especially of the nervous system. This is supported by the facts that the injection of thyroid extract does good when the function of the thyroid is in abeyance; that microscopically it can be proven that the cells secrete the colloid material, and that it eventually finds its way into the blood by way of the lymphatics; that the beneficial portion of the secretion can by chemic means be separated from the gland, thus demonstrating that it is a secretion of that body. A great deal of work has been done to isolate the active principles of the thyroid secretion. Sigmund Fraenkel has been successful in isolating a substance which is intensely hygroscopic and soluble in water and alcohol, the empiric formula of which is $C_6H_5N_3O_5$, to which he has given the name of thyreo-antitoxin.

Experimenting on animals he found that under its administration there was acceleration of the pulse rate without the fall of blood pressure demonstrated by Schaeffer on intravenous injection of the extract. He also found that the heart of a frog, poisoned by muscarin, which had ceased to beat, was stimulated to activity by a few drops of a watery solution of thyreo-antitoxin dropped upon it. Moreover, he demonstrated that in animals deprived of the thyroid gland with the usual subsequent symptoms, that temporary relief, at least, was given by a subcutaneous injection of a 1 per cent. watery solution of thyreo-antitoxin. He maintains that this substance is an exact chemical and that eventually he will be able to give exact dosage. It is without odor and has a beefy taste.

Baumann and Roos make stronger claims than Fraenkel. By a series of experiments they have been able to isolate a brown amorphous substance which contains the remarkably large quantity of 9.3 per cent., dry weight, of iodine. It has been named thyro-iodine. It is almost insoluble in water, only slightly so in alcohol, but easily so in dilute alkalies. The discoverers claim that its administration is followed by the same results as follow the administration of the gland itself, or its extract, in myxedema, obesity and parenchymatous goiters, and indeed that it is more rapid in its action. The amount of this substance varies in different thyroids, large ones only containing a trace of iodine, while in children there is less than in adults. Sheep thyroids are relatively richer in iodine and the maximum amount in humans is found between the ages of twenty-five and fifty. Drechsel has isolated two crystalline basic bodies from the thyroid, one of which is apparently identical with Fraenkel's thyreo-antitoxin. Both of these when administered to thyroidectomized animals exert a beneficial influence and Drechsel suggests that there "may