There is a train of eye symptoms which we frequently see in children, comprising exophthalmos, chemosis, keratitis, etc., almost invariably due to a thrombophlebitis arising from empyema of one or more of the accessory sinuses. Such extreme conditions are frequently the result of the non-recognition of the seriousness of the so-called common cold in children. If the pediatrician or general practitioner does not know how to handle these cases, he should early seek the advice and help of the rhinologist.

Dr. Skillern did not mention suction as a form of treatment. Suction properly applied will give brilliant and satisfactory results. It can be used by the pediatrician as well as by the rhinologist, and not infrequently obviates all necessity for operative procedures that would otherwise be required.

Dr. Skillern says the Luc operation should not be performed. Personally, I feel that radical operations are safer than intranasal work. It is hard to do intranasal work in infants. It necessitates the use of ether, and the intranasal exenteration of ethmoids under general anesthesia is a dangerous operation even in adults.

DR. GREENFIELD SLUDER, St. Louis: It seems to me that Dr. Skillern is proving a difference with very little differentiation, but Dr. Coffin has emphasized that. That an inflammatory involvement of the lateral wall of the nose be differentiated from a sinusitis, it seems to me is pretty much of a nicety. I believe these cases are to be classed as a sinus inflammation quite as much as in the older patients with a pus cavity. They come under not so much a suppurative head, as Dr. Skillern emphasizes, but under a hyperplastic classification. Dr. Skillern spoke of all the sinuses but the sphenoid. The sphenoidal sinus—and in this statement I quote his authority, Warren Davis of Philadelphia—the sphenoid begins in the sixtieth day of fetal life. At the end of two years it has reached laterally to the foramen rotundum; at the end of six years it has reached the floor of the sinus and the Vidian canal. Sinus involvement in young persons seems to be difficult, and more so when in the posterior portion, according to age. I feel absolutely sure that sphenoid—and when I mention that I include postethmoid—involvement is of frequent occurrence in childhood. I have been in the habit of separating infancy at the second year from childhood beyond the second year. The diagnosis of postethmoidal sphenoiditis in children is exceedingly difficult. Postnasal examination is to be achieved rarely. In such cases suppuration is not profuse. There is pain and swelling and a more or less serous discharge, which continues. These cases I have seen repeatedly exploited, as Dr. Beifeld has said, as psychic vomiting. I have repeatedly seen them made the asthma of childhood, and have seen that asthma recover after postethmoidal interference.

DR. ROSS H. SKILLERN, Philadelphia: There is no question of the truth of Dr. Coffin's statement regarding the baby sinuses. I quoted directly from him in this respect. I do not know anything about the suction method of treatment in children. I have never tried it but should like to. I imagine it would be a rather difficult procedure, however. We all know that it is difficult enough to treat children anyway, and to give suction in these cases it seems to me would be a difficult thing.

Dr. Sluder says that in most cases the ethmoids are hyperplastic. That is true. He did not mention that these cases occur in infants under 2 years of age, but I suppose that is what he means. I do not recall seeing a case of hyperplastic ethmoids in a baby. Regarding the connection between the ethmoids and sinuses, that is true, but at the same time, I cannot help feeling that a mass such as the ethmoid, representing structurally a sponge covered with thin mucosa, is totally different from the smooth sinus with hard, bony walls, and it seems to me that it must be, and often is, affected in a totally dissimilar manner.

Signs of Death.—It is a sign of death when the forehead grows reddish and eyebrows sink, the left eye decreases, the tip of the nose becomes white, the chin falls, while the pulse runs, feet grow cold and the belly is relaxed.—Galen.

EXPERIMENTAL PYELITIS IN THE RABBIT*

H. F. HELMHOLZ, M.D.

AND

CAROL BEELER, B.S.

EVANSTON, ILL.

Numerous attempts have been made in the past to produce pyelitis by injecting bacteria into the bladder, with the idea of getting an ascending infection, and by injecting organisms into the circulation and expecting them to set up lesions in the kidney during the process of being filtered out. It has long been known, and it is becoming increasingly more evident, that bacteria during any infection, if they reach the blood stream, are likely to be excreted in the urine. This

Fig. 1.—Spontaneous pyelitis (distended ureters and pelvis).

applies to acute as well as chronic diseases. The kidney acts as a filter to remove bacteria from the circulation. If this is constantly taking place, we must naturally assume that the kidney has a great resistance to these organisms or that the bacteria are carried out so rapidly that they do not have an opportunity to multiply.

The two possibilities for infection are: (1) a loss of resistance of the kidney or an increase in the virulence of the organism; and (2) a stopping up of the natural washing-out of the urinary passages. The latter one of these factors has been studied by a great many observers, who have all showed that anything that obstructs the flow of urine makes possible an

* From the O. S. A. Sprague Memorial Institute Laboratory of the Children's Memorial Hospital.

* Read before the Section on Diseases of Children at the Sixty-Eighth Annual Session of the American Medical Association, New York, June, 1917.

infection of the kidney, whether the infecting organism is introduced from the bladder or by the blood stream. Furthermore, the work of Stewart¹ on ascending infection of the kidney by ureteral transplantation into the rectum has shown that the infection spreads upward by the lymph channels, and that the spread by the lumen of the ureter is less of a factor than formerly supposed. In this work, however, the lymph channels were cut and naturally offered excellent opportunity for the spread of infection. Hess² came to the conclusion that something more than the simple injection of colon bacilli into the bladder was necessary to produce ascending infection in the rabbit. The previous injection of turpentine or other irritant into the bladder was sufficient to bring about an involvement of the upper urinary tract. More recently Eisendrath and Schultz³ have attempted to show that infection of the bladder with *Bacillus coli*, streptococci and *B. proteus* travels up the periureteral lymphatics. They used rabbits and dogs in their work. They described very extensive changes in nineteen of thirty-eight animals but recovered organisms from the kidney in only six of their positive cases in their first series and in none of their second. It seems that in view of the negative bacteriologic results, the causal relationship of the lesions and the organisms injected must be left open.

Fig. 2.—Hematogenous cortical abscesses.

It is not our object here to go into a discussion of the etiology of pyelitis, but merely to point out that with an unobstructed urinary tract there have been practically no constantly positive results in the production of pyelitis experimentally, either by intravenous or intracystic injection. In a previous series⁴ of intravenous experiments we were able to produce kidney lesions in only eight out of sixty-six animals injected with colon bacilli, and in four out of eleven animals when injected with a mixture of colon bacilli and pneumococci, the tendency of the organisms to produce lesions in other organs being just as great as in the kidney. During the course of the experiments just cited, in controlling the urine of all animals before injection, we found a female rabbit with a constantly purulent urine. Cultures taken from a catheterized specimen of urine showed a pure culture of a gram-negative bacillus, that had the following characteristics:

The organism grew well on all mediums. There was a profuse yellowish growth on agar. Gelatin was not liquefied. Litmus milk was rendered acid and coagulated; but there was no digestion of the casein. Indol was produced. There was acid and gas formation in dextrose, lactose and saccharose. The organism thus belongs in the group of *Bacillus coli-communior*.

1. Stewart: A Study of Ascending Infection, etc., Univ. Pennsylvania Med. Bull., 1910, 23, 233.
2. Hess: Mitt. a. d. Grenzgeb. d. Med. u. Chir., 1913, 26, 135.
3. Eisendrath and Schultz: Jour. Med. Research, 1917, 35, 295.

4. Helmholtz, H. F., and Beeler, Carol: Focal Lesions Produced in the Rabbit by Colon Bacilli Isolated from Pyelocystitis Cases, Am. Jour. Dis. Child., July, 1917.

SUMMARY OF INJECTIONS

Series	No.	Kidney Lesions				Other Organs Involved
		Total	Pyelitis	Abs.	Bact.	
A. Intravenous Injection						
I	10	6	4	4	..	Hemorrhage into lung.
II	2	2	2	1	..	Abscess of seminal vesicle.
III	4	2	2	1	..	Hemorrhage of cecum and rectum.
IV	2	2	1	2	..	0
V	5	4	2	3	..	0
VI	2	2	1	2	..	Chronic nephritis.
VII	2	1	1	0	..	Generalized peritonitis.
VIII	2	1	1	1	..	0
IX	3	2	2	2	..	0
	32	22	16	16		
B. Intracystic Injection						
II	2	0	0	0	..	Infected Right Testis.
IV	2	2	2	0	..	0
V	5	3	1	0	2	0
VI	2	2	2	1	..	0
VII	2	2	2	0	..	0
VIII	2	2	2	0	..	0
	15	11	9	1	2	
C. Rectal Injection						
VII	2	0	0	0	..	0
VIII	2	0	0	0	..	Gram-negative bacillus.*
D. Oral Administration						
V	5	0	0	0	..	1x Bladder Gram-positive coccus; Gram-negative bacillus.
VI	4	0	0	0	..	2x Bladder Gram-negative bacillus.

* From bladder urine.

HISTORY

Feb. 17, 1917. The first urine examined showed about 150 pus cells to a low power field.

February 9. There was a large excess of pus cells.

February 10 to February 20. Daily examination revealed a large number of pus cells, single and in groups.

Fig. 3.—Ascending pyelitis (distended ureter).

February 20. A catheterized specimen of urine was cultured in litmus lactose agar, plain agar and broth.

February 21. There are innumerable colonies on both plates.

February 23. Specimens were expressed every other day showing albumin and casts in addition to pus cells.

March 3. Another catheterized specimen was cultured; pure culture same as obtained on February 20.

The animal was chloroformed, as it was getting very much weaker.

Necropsy.—On opening the peritoneal cavity the abdominal organs all appear practically normal. The bladder is distended with urine. The stomach, gastro-intestinal canal, spleen, liver, gallbladder, heart and lungs show nothing abnormal. On dissecting free the overlying organs the ureters on both sides are seen to be considerably dilated, as far down as the brim of the pelvis. Beyond this point they appear to be of normal diameter. No apparent pathologic constriction in the ureter at the brim of the pelvis was visible. The pelves of both kidneys were smooth, and of a reddish brown color; the capsule stripped readily. On section the cortex was of normal thickness, the striations regular. The outer portion of the medulla was deeply congested. The pelves in both kidneys were markedly distended, as seen in Figure 1. The left kidney pelvis was filled with thin purulent material, the right practically empty. The bladder urine showed only relatively few leukocytes, numerous casts and much debris. Cultures taken from the bladder and right kidney pelvis yield a pure culture of gram-negative bacillus.

Histology.—The bladder mucous membrane in the entire section is practically intact. In areas there are small clusters of lymphocytes; there are practically no polymorphonuclear leukocytes to be seen in the section. Kidney: There is considerable degeneration of the epithelium and dilation of collecting tubules; in areas focally there is considerable increase in connective tissue. In the medulla, especially toward the tip of the papilla, there is marked leukocytic infiltration about the tubules.

Eosinophil cells are much in evidence. The mucosa of the pelvis is apparently thickened and shows considerable infiltration with leukocytes. The pelvis is filled with masses of polymorphonuclear leukocytes.

The colon bacillus isolated from the catheterized specimen of urine was used in nine series of experiments. The technic used in the experiments was as follows: The intravenous injections were made with a bacterial suspension that was made by centrifuging 15 c.c. of a twenty-four hour broth culture and suspending the bacilli in 5 c.c. of physiologic sodium chlorid solution. One c.c. of this suspension was injected into the ear vein. The intracystic injections were made after carefully catheterizing the animals and then allowing 5 c.c. of the bacterial suspension to run into the bladder.⁵ In the feeding experiments 8 c.c. were injected. In the first series of ten animals which were injected intravenously, six showed lesions of the kidney; in four instances a pyelitis, and in four instances cortical or medullary abscesses. In two instances, as is evident from the table, both were present. In Series 3 and 9 also, all the injections were intravenous. In the other series the same culture was used in the injections given intravenously, intracystic and either rectally or orally, in order that the results could be more accurately compared.

In the thirty-two animals of nine series injected intravenously, twenty-two showed lesions of the kidney, in sixteen instances pyelitis and in sixteen instances cortical (Fig. 11) or medullary abscesses. In ten instances there were both abscesses and pyelitis. No doubt if the animals with abscesses had lived they would all have developed a pyelitis also. Of the thirty-two animals only three showed lesions outside of the urinary tract that could be ascribed to the colon bacillus: one instance each of abscess of the seminal vesicle, hemorrhage of the

Fig. 4.—Ascending pyelitis (cortical abscesses).

cecum, and generalized peritonitis. The peritonitis was due in large measure no doubt to extensive coccidiosis. The absence of lesions in other organs is just as striking as the localization of the organism in the kidney in twenty-two of the thirty-two cases.

The intracystic injections consist of six series, totaling fifteen animals. In four series of two animals each, pye-

5. Three control animals injected with a dilute solution of gentian violet showed that no fluid passed beyond the bladder.

litis was produced in 100 per cent. In the fifth only one had a definite pyelitis, but in two others gram-negative bacilli were isolated from the pelvis of the kidney. In the sixth series neither one of the two animals injected acquired a kidney infection. Figure 3 shows the swelling of the ureter, Figure 4 the one kidney with cortical lesions in the intracystic series.

Four animals that received rectal injection and nine that were repeatedly fed the colon culture showed no lesions of the urinary tract, except that in one instance in the rectal series and in three instances in the oral series, colon bacilli were isolated from the bladder but not from the kidney.

The serum of a rabbit immunized to the specific organism isolated from the urine agglutinated the organism in a dilution of 1:2,000. It did not agglutinate several strains of *B. coli-communior* isolated from human cases of pyelitis nor several strains of *B. coli* isolated from rabbit feces. A *B. coli-communior* isolated from the feces of the original pyelitis rabbit was agglutinated in the same dilution as the colon bacillus isolated from the urine.

COMMENT

The remarkable specificity of this organism is of interest from two very different points of view: first, it enables us to study better than ever before the etiology and probably the treatment of kidney infections, and secondly, from the purely biologic point of view, it shows us the extreme specificity the colon bacillus can acquire in localizing in the kidney, on intravenous injection.

These experiments prove without a doubt that pyelitis can be produced experimentally by intravenous injection as well as by injection into the bladder. What the route of the infection is from the bladder to the kidney we must leave an open question at this time. One point that we do wish to make, however, is that this work shows that if the organism has a specific affinity for the kidney its introduction into the bladder can be of very serious consequence to the individual. It will take further study to determine whether the bacilli reach the kidney via the lumen, the lymphatics or the blood stream. Experiments in that connection are at present under way.

If we compare these experiments with the results obtained by the injection of organisms isolated from human pyelitis cases, the specificity of this organism is even more striking. In sixty-six animals injected with different strains of *B. coli* only eight showed kidney lesions, and twenty-one lesions of the other organs; with this organism, of thirty-two animals injected twenty-two showed kidney lesions and in only three instances were the other organs affected.

The so-called epidemics of pyelitis that have been described from various European clinics might well be ascribed to an organism that, as this organism, has a specific affinity for the kidney tissue.

The specific localization of organism in tissues has been of great interest, especially since the work of Rosenow on the pneumostreptococcus group. Rosenow⁶ in his work on the colon bacillus and streptococcus in gastro-intestinal infection, especially of the appendix, showed that the combination had specific tendencies to localize but that the colon bacillus alone had no such powers. Our work on the colon bacilli of human pyelitis cases led to practically the same result. With this organism the specificity manifests itself not only in that the kidney is involved in about 70 per cent. of the injections, but that in less than 10 per cent. of the injections is there involvement of any other organ. The intracystic infections are not of

much importance in this connection because the distribution of the bacteria is not so widespread throughout the system as with the intravenous injections.

SUMMARY

1. Infections of the kidney were produced in 70 per cent. of all animals injected by the intravenous or intracystic route with a colon bacillus obtained from a rabbit with spontaneous pyelitis.

2. The organism, a *Bacillus coli-communior*, had a specific tendency to localize in the kidney tissues.

A CASE OF POISONING BY MUSTEROLE

DAVID I. MACHT, M.D.

BALTIMORE

The following case should be of interest not only to the pharmacologist and toxicologist but also to the general practitioner:

Mrs. S., aged 32, consulted me, May 31, 1917, on account of a rash. There was a scarlatiniform eruption over the neck, face, chest, arms and back. The rash resembled very much that of scarlet fever. Further examination, however, contradicted such a diagnosis. There was no history of exposure to scarlet fever; the patient's temperature was normal; the pulse rate was 76; there was no sore throat, "strawberry tongue" or any other sign of scarlet fever, and the general condition of the patient was good except that she complained of itching and burning. A very marked irritation of the conjunctivae was also noted. Physical examination of the chest and the abdomen was negative. There were no abnormal urinary findings.

On investigating the case more carefully, it was learned that the patient on the day before was complaining of pain behind the right ear and that some friends advised her to apply musterole to the painful spot. Accordingly the patient rubbed in thoroughly some of the nostrum in the right mastoid region, and on the following day the general exanthem appeared.

Musterole has been described¹ as being essentially a mixture of lard or some similar material, with oil of mustard, menthol and camphor.

Oil of mustard is known to the chemist and pharmacologist to be an exceedingly irritating drug attacking especially the eyes. When applied to the skin for any length of time it quickly draws blisters. Erysipelatous eruptions, general dermatitis, and even necrosis of the skin have also been described following its use. If absorbed into the circulation, oil of mustard may produce albuminuria and even nephritis.² Its use in therapeutics is for that reason very limited, and when it is employed, care should be exercised in handling it. Mustard plasters are for all purposes safer than the oil of mustard, and can be employed whenever the oil is indicated. A mixture of such an irritant oil as oleum sinapis and lard is very liable to be absorbed through the skin, and therefore the indiscriminate use of such oils in salves under fanciful names, without a physician's supervision, should be discouraged, as it may lead to untoward effects, such as in the case described above.

3218 Auchentoroly Terrace.

1. Musterole, THE JOURNAL A. M. A., Jan. 20, 1917, p. 214.

2. Compare Lewin: Nebenwirkungen der Arzneimittel, Berlin, 1899, p. 679.

6. Rosenow: Jour. Infect. Dis., 1915, 16, 240.