TRANSACTIONS

OF

THE LINNEAN SOCIETY.

I. *A Monograph of Recent Brachiopoda.*—Part I.
By THOMAS DAVIDSON, LL.D., F.R.S., F.L.S., F.G.S., &c.

Read 5th November, 1885.

(Plates I.–XIII.)

INTRODUCTORY REMARKS.

DURING the last hundred years the recent Brachiopoda have attracted considerable attention, and a large number of valuable memoirs and papers have been published upon them. Their shells, shell-structure, anatomy, embryology, and affinities have alike been carefully investigated. Observations on the living animals of several genera have also been recorded. The sea-bottoms have been dredged for Brachiopoda in many latitudes and over a wide geographical area, and their habitats and ranges of depth accurately ascertained to a very considerable extent. Four or five incomplete monographs, in which the shells only of a large number of species have been well illustrated and briefly described, have appeared during the present century; but no satisfactory general monograph treating of the shell and animal *conjointly* has yet been published. This omission I have now endeavoured to supply.

In 1843, Küster, in his new edition of Chemnitz's 'Conchylien-Cabinet,' described some twenty-six or thirty species, of which several are now known to be synonyms. These he figured in six quarto plates.

In 1846, G. B. Sowerby, in his 'Thesaurus Conchyliorum,' described and beautifully illustrated forty-seven species, of which number several are synonyms.

In 1859–62, Lovell Reeve, in his 'Conchyliorum Iconica,' described the shells of seventy-five species, of which some were synonyms, accompanied with a series of beautiful illustrations, drawn by G. B. Sowerby.

In 1873, in the 'Proceedings of the Academy of Natural Sciences of Philadelphia,' Mr. W. H. Dall published a catalogue of all the recent species of Brachiopoda known to

him up to that date. In this catalogue without figures, about one hundred species are enumerated, some of which are synonymous.

During the last thirty-five or more years, I have devoted much time to the study of the recent forms, in conjunction with that of the fossil species, and have lost no opportunity of making myself acquainted with all that has been done and written upon the subject, as well as in assembling all available material, so as to enable me to bring together in a single monograph the chief results of many independent researches published in a number of scattered papers and works often difficult of access. The literature of the subject is indeed voluminous, as may be realized by a glance at the 'Bibliography of the Brachiopoda,' compiled by Mr. W. H. Dalton and myself, and published in vol. vi. of my 'British Fossil Brachiopoda' (Palæont. Soc., 1886).

I have also, I believe, had advantages which few have possessed in being able to follow out the observations made with respect to the animal and its anatomy, and in having been able to draw a very large number of figures from the types of the best-preserved examples of almost all the known forms, as well as of a large series of individuals of the same species at different stages of development. The study of the adult condition of a species gives insufficient data, and it is requisite to follow out the modifications it has to go through during the different stages of its existence, and to note these differences.

The study of the embryo has also shown that the animal assumes a series of well-defined stages in its development, a fact that was but little known prior to the publication in 1861 of Prof. Lacaze-Duthiers's admirable memoir on *Thecidium mediterraneum*. These observations were subsequently followed by the excellent researches of Fritz Müller, Kowalevsky, E. Morse, H. Friële, M'Crary, Dall, Van Bemmelen, A. E. Shipley, M. A. Schulgin, and one or two more. The results obtained by these authors will be referred to in the sequel. It is very desirable that these important investigations should be continued, as much still remains to be discovered, described, and illustrated.

The shell-structure of the recent Brachiopoda has been admirably worked out by a number of accurate observers, such as Dr. W. B. Carpenter, W. King, Van Bemmelen, Hancock, and many others, and has led to very important results. To Herman Friële, E. Deslongchamps, and one or two others we are indebted for much accurate and important knowledge with respect to the development of the loop, of which but little was known previous to 1852.

The anatomy of the animal has also been admirably investigated and worked out, and it is sufficient to mention the names of Cuvier, Owen, Huxley, Hancock, Vogt, Gratiolet, Lacaze-Duthiers, King, Brooks, Dall, Morse, E. Deslongchamps, Van Bemmelen, Woodward, Shipley, Schulgin*, and others, to show how important and varied have been the additions to our knowledge with respect to this very necessary branch of investigation. In drawing up the description of each species, I have considered it desirable, whenever possible, to reproduce the words and illustrations of the authors, and thus give them all credit for their careful, painstaking researches.

* To these names Dr. Davidson would doubtless have added that of H. G. Beyer, who contributed an important paper on the shell-structure and anatomy of *Lingula (Glottidia) pyramidata*, Stimpson, to the Studies from the Biological Laboratory of the Johns Hopkins University, Baltimore, vol. iii. no. 5, March 1886.—[A. C.]

The perplexing question of the affinities of the Brachiopoda has given rise to much discussion, and great difference of opinion, especially with regard to their relationship to the group of worms. Now, although I do not admit the Brachiopoda to be worms, they may, as well as the Mollusca and some other groups of invertebrates, have originally diverged from an ancestral vermiform stem, such as the remarkable worm-like mollusk *Neomenia* would denote. In a recent paper on the development of *Argiope* or *Cistella*, Mr. A. E. Shipley observes, and, I think, with justice, that the Brachiopoda and Polyzoa are not so closely united as to form a natural phylum; and he adds, "I should propose to follow Gegenbaur in making a primary class of the Brachiopoda, and though in their development and adult structure they are widely separated from both Vermes and Mollusca, of the two classes I would place them nearer to the former class than to the latter"*. Prof. Huxley† says:—"All known Polyzoa are compound animals, that is to say, the product of every ovum gives rise, by gemmation, to great assemblages of partially independent organisms, or zooids. The Brachiopoda, on the contrary, are all simple, the product of each ovum not giving rise to others by gemmation. All the Brachiopoda possess a bivalve shell—a shell composed of two, more or less horny, or calcified, pieces, which are capable of a certain range of motion on one another, and are very commonly articulated together by teeth and sockets." The shell, the pallial lobes, the intestine, the nerves, and the atrial system, afford characters amply sufficient to define the class.

In this view of Prof. Huxley I entirely concur.

As many species of Brachiopoda live at considerable depths, it is not surprising that so small a number should have been known to early conchologists, and that for many years they should have been such great rarities in conchological collections. The numerous well-conducted dredging expeditions have, however, brought to light a large number of forms that were not previously known, and we may constantly expect to add to the number of species as dredging operations extend to regions not yet explored. It has been ascertained beyond doubt that Brachiopoda are much localized, and that where they occur they are generally abundant. It has also been found that the range in depth of one and the same species is often very variable, that abyssal forms have generally a very thin shell, and that species living at a great depth have a much greater geographical range, and are not nearly so localized as those species that live in shallow waters.

The study of the species brought home by the 'Challenger' Expedition, which I was privileged to examine and describe, has revealed much valuable information with respect to the bathymetrical and geographical distribution of many species. The greatest depth at which a recent species of the class has been found alive was 2900 fathoms. A number of forms inhabit and prefer rocky and stony parts of the bottom, or are attached to corals, and are therefore more difficult to obtain.

It is necessary briefly to refer to the difficult question of classification, upon which many different opinions have been entertained. In company with a larger number of

* "On the Structure and Development of *Argiope*." Mittheilungen aus der zool. Station zu Neapel, Band iv. Heft 4, p. 516 (1883).

† An Introduction to the Classification of Animals, p. 27 (1869).

malacologists and palæontologists, I have considered the interior skeleton that supports the labial appendages as a classificatory character that could be advantageously made use of, and have consequently grouped the recent species into the two great divisions *Arthropomata*, Owen (= *Clistenterata*, King), and *Lyopomata*, Owen (= *Tretenterata*, King), and into six families, as follows :

ARTHROPOMATA, Owen = <i>Clistenterata</i> , King.			
		Species.	Uncertain species.
1st Family TEREBRA- TULIDÆ.	Subfamily TEREBRATULINÆ {	I. Genus <i>Liothyris</i> , Douvillé	8 2
		II. Subgenus <i>Terebratulina</i> , d'Orbigny . .	10 5
		III. Genus <i>Waldheimia</i> , King	10 1
	Subfamily TEREBRATELLINÆ . . {	IV. Genus <i>Terebratella</i> , d'Orbigny	9 3
		V. Subgenus <i>Magasella</i> , Dall	6 6
	Subfamily MEGERLINÆ {	VI. Genus <i>Megerlia</i> , King	2 1
		VII. Subgenus <i>Laqueus</i> , Dall	3 1
	Subfamily MAGASINÆ	VIII. Genus <i>Bouchardia</i> , Davidson	1 1
	Subfamily KRAUSSININÆ {	IX. Genus <i>Kraussina</i> , Davidson	5 1
		X. Subgenus <i>Megerlina</i> , Deslongchamps .	2 1
	Subfamily ARGIOPINÆ {	XI. Genus <i>Argiope</i> , Deslongchamps	1 2?
		XII. Subgenus <i>Cistella</i> , Gray	8 1
	Subfamily not yet determined. {	XIII. ? <i>Gwynia</i> , King	1 1
		XIV. Genus <i>Platydia</i> , Costa	2 1
2nd Family THECIDIIDÆ		XV. Genus <i>Thecidium</i> , Defrance	2 1
3rd Family RHYNCHONELLIDÆ {		XVI. Genus <i>Rhynchonella</i> , Fischer	6 1
		XVII. Subgenus <i>Atrertia</i> , Jeffreys	2 1

LYOPOMATA, Owen = *Tretenterata*, King.

4th Family CRANIIDÆ	XVIII. Genus <i>Crania</i> , Retzius	4	1
5th Family DISCINIDÆ	{ XIX. Genus <i>Discina</i> , Lamarek	1	
		XX. Subgenus <i>Discinisca</i> , Dall	6
6th Family LINGULIDÆ	{ XXI. Genus <i>Lingula</i> , Bruguière	8	3
		XXII. Subgenus <i>Glottidia</i> , Dall	3

In 1884, M. E. E. Deslongchamps proposed a new scheme of classification for the Terebratulidæ, in which he objected to any arrangement based on either the exterior shape of the shell or of the supports of the labial appendage.

His first group includes the different forms in which the calcified brachial apparatus or loop does not undergo any important modifications from its first origin up to the adult condition. To the characters drawn from the brachial apparatus or loop is added that of the presence of spicula, more or less complicated, which occupy in the mantle all the parts connected with the organs of circulation (arteries, veins or veiny sinuses, &c.), the labial appendages and cirri which accompany them. In this group he places the recent genera *Liothyris*, *Terebratulina*, *Megerlia*, *Kraussina*, and *Platydia*.

In his second group he unites those forms in which the brachial apparatus or loop undergoes numerous transformations from the embryo up to the adult condition, and which have been distinguished by the names of *Platydiiform*, *Magadiiform*, and *Megerli form* stages. The mantle in this group is not provided with those calcareous spicula which occur so constantly in the first group. The colour also of the dried animal is yellowish, whilst in the first group the hue or tint of the dried arms and of the periph-
 eric portions of the mantle are of a very characteristic dim white. This, I may, however, remark, is not always the case, as I possess specimens of the dried animal of several species of *Terebratulina* that are of a decided yellow colour. The group would, according to M. Deslongchamps, comprise the recent genera *Waldheimia*, *Macandrevia*, *Terebratella*, *Laqueus*, and *Magasella*.

While fully appreciating the importance of all characters derived from a study of the animal, I am not convinced that the temporary modifications in the shape of the loop, or the presence or absence of calcareous spicula in the mantle, &c., are indications of sufficient importance or permanence to supersede those derived from the adult shape of the calcareous lamellæ supporting the labial appendages,—characters which are often accessible, and of important assistance in distinguishing the more numerous fossil members of the group. Moreover, Mr. W. H. Dall, in describing the animal of *Waldheimia floridana* (Bull. Mus. Comp. Zoöl. Harvard, vol. iii. p. 16, 1871), distinctly notes the existence of a few exceedingly delicate spicula in the floor of the great sinuses. It would seem therefore that these spicula occur in a genus which M. E. E. Deslongchamps, in his proposed new classification of the Terebratulidæ, places among those forms characterized by the entire absence of spicula.

After long and searching examinations of the recent forms, I have described in this monograph about one hundred so-termed species, some varieties, and about twenty-eight uncertain ones. Of course the vexed question as to what really constitutes a species remains the same, and is likely to remain so for a long time to come.

It will not be necessary to extend these introductory remarks, as all details have been fully given under each species. In conclusion I would tender my grateful thanks to the many kind friends who have in so zealous a manner supplied me with valuable information and specimens*.

ARTHIROPOMATA, Owen=CLISTENTERATA, King.

Family TEREBRATULIDÆ, (Gray) emend. Davidson.

Subfamily TEREBRATULINÆ, Dall, 1870.

During the last few years a strong desire has been manifested by those palæontologists who consider an extreme subdivision of genera desirable, to separate from *Terebratula* proper those forms characterized by a small short loop, of which the principal stems are united anteriorly by a slightly arched lamella, and of which *Liothyris vitrea* may be taken as the type.

* The drawings for the Plates were made by myself, on paper; but the state of my health would not allow of my reproducing them on stone.

As there are certainly some differences observable in the two groups, and as none of the recent species would agree in the characters of their loops and in certain other particulars with the forms referred to the genus *Terebratula* of Lhwyd and Klein, it may perhaps be better to adopt Douvillé's generic name *Liothyris* for the species we are about to describe.

In the larger number of the recent species, such as in *Liothyris vitrea*, *L. arctica*, *L. Moseleyi*, *L. uva*, *L. Bartletti*, *L. Wyvillii*, and *L. subquadrata*, the connecting band of the loop is narrow, while in *L. sphænoidea* = *cubensis* it is larger.

The specific claims of *Liothyris cernica*, and *L. ? Dalli* are still uncertain, only a single example of each of them having been hitherto discovered.

Very small, or scarcely any modifications in the shape of the loop have been observed; all the species have their shell minutely perforated by canals, and calcareous spicules are abundant in the mantle.

1. LIOTHYRIS VITREA, Born, sp. (Plate I. figs. 1-12.)

Anomia vitrea, Born, Testacea Musei Caes. p. 119, vignette, 1778; Linné, ed. Gmelin, p. 3347, 1788.

Gryphus vitrea, Megerle v. Mühlfeld, Berlin Mus. 1811.

Terebratula vitrea, Lamarck, An. sans Vert. vol. vii. p. 245, 1819; Payraudeau, Cat. p. 83, no. 160, 1826; G. Bronn, Italiens Tertiär-Gebilde, p. 125, 1831; Philippi, Enum. Moll. Siciliæ, vol. i. p. 95, t. 6. figs. 6-8, 1836, vol. ii. p. 66, 1844; A. Scacchi, Cat. Conch. Regni Neapolitani, p. 8, 1836; Küster, Martini & Chemnitz, Conch.-Cab. vol. vii. p. 22, tab. 2. figs. 11-13, 1843; E. Forbes, Report on the Mollusca of the Ægean Sea, Brit. Assoc. Report, p. 141, 1843; D. Galvani, Illustrazione delle Conch. Foss. 1845; G. B. Sowerby, Thes. Conch. vol. i. p. 353, pl. 70. figs. 56-59, 1846; Aradas (pars) Conchiglie fossili di Gravatelli, p. 14, 1851; O. G. Costa, Fauna del regno di Napoli, p. 33, pl. i. figs. 1-3, 1851-52; Davidson, Sketch of a Class. of recent Brachiopoda, Ann. & Mag. Nat. Hist. vol. ix. p. 364, 1852, and Br. Foss. Brach. Pal. Soc. vol. i., Introduction, p. 62, fig. 23, and pl. vi. 1852; S. P. Woodward, Manual of Mollusca, p. 215, 1856; L. Reeve, Conch. Icon. pl. 3. fig. 8, 1860, and Journ. de Conch. vol. ix. p. 124, 1861; Chenu, Man. de Conch. vol. ii. p. 201, 1862; Seguenza, Atti della Soc. Italiana di Scienze Nat. vol. i. p. 17, pl. 1. figs. 1-7, 1865; H. C. Weinkauff, Die Conch. Mittelmeeres, vol. i. p. 284, 1867; Davidson, Italian Tert. Brach., Ann. & Mag. Nat. Hist. vol. vii. pl. xvii. fig. 11, 1870; Jeffreys, Ann. & Mag. Nat. Hist. 5th ser. vol. x. p. 28, 1882.

Liothyris vitrea, Douvillé, Bull. Soc. Géol. de France, 3^e sér. vol. vii. 1879; E. Deslongchamps, Note sur la Classification des Térébratules, ou Études critiques sur les Brachiopodes, pp. 106 & 153, pl. xx. figs. 7-11, 1884.

Shell longitudinally oval or ovate, globose, widest about the middle, laterally rounded or more or less pinched in near the front, front margin nearly straight or gently rounded. Colour nearly white, surface smooth, semitransparent, glassy, marked with fine concentric lines of growth and perforated by minute canals. Dorsal valve tumidly convex, longitudinally flattened along the middle, from which the anterior lateral portions slope to the edge. Ventral valve slightly deeper than the dorsal one, longitudinally flattened along the middle; beak incurved, moderately produced and slightly overlying the umbo of the dorsal valve, obliquely truncated by a very small circular foramen with thickened margin and separated from the hinge-line by a small triangular deltidium in two pieces. Loop in the interior of the dorsal valve simple and short, attached by its

crura to the hinge-plate and not exceeding a fourth of the length of the valve, the two principal stems becoming soon united anteriorly by a transverse lamella bent upwards in the middle; no mesial septum; cardinal process small and prominent, hinge-plate disunited, four diverging grooves extending from under the cardinal process to about half the length of the valve, the central pair being the longest. In the interior of the ventral valve a similar number of grooves; muscular impressions small, situated at the bottom of the valve under the loop and in the rostral portion of the bottom of the ventral one. Animal attached by a peduncle; labial appendages united to each other by a membrane; brachial disk trilobed; central lobe elongated and spirally convoluted. Very delicate spicula form elegant star-like plates in the mantle. Length 1 inch 8 lines, breadth 1 inch 5 lines, depth 1 inch.

Hab. Abundant in the Mediterranean in depths of from 90 to 250 fathoms. Very numerous in the Bay of Naples at depths of from 100 to 300 metres. Vigo Bay, 40 fathoms (M^cAndrew). Dredged by Prof. Giglioli, during the Italian Expedition to the Mediterranean in 1881, at a depth of 800 fathoms (see report in the *Atti del iii. Congresso Geografico Internazionale*). Also for distribution of this and other species, see Jeffreys's papers "On the Mollusca of the 'Lightning' and 'Porcupine' Expeditions," *Proc. Zool. Soc. Lond.* 1878, 1879, 1881, 1882.

Obs. *Liothyris vitrea* is a beautiful, well-known, and abundant Mediterranean species. It varies considerably in its relative length, breadth, and degree of convexity; some specimens being quite elongated oval, 1 inch 8 lines in length by 1 inch 1 line in breadth, while other examples of the same length would have a breadth of 1 inch and 3 or 4 lines; some are much more pinched in anteriorly than others, and, lastly, some are nearly circular with an equal length and breadth.

Prof. E. Deslongchamps, in his instructive memoir on the classification of the Terebratulidæ, gives us the result of his studies in connection with the embryo of *L. vitrea*, as well as of its subsequent stages of development. He states, "I have been able to examine the embryo at two millimètres of size, that is to say from the first moments when the larval condition has ended and the shell has begun to be formed; its shape is then absolutely similar to that of the young of *Terebratulina*; the dorsal valve is rounded, and slightly convex; the ventral one shows a triangular hole, of which the summit, which does not yet show any trace of a notch, will eventually become the beak. No trace of a deltidium is to be seen on the sides of this foramen. On opening this little shell, one is at once struck by the dull white of the internal walls; and, by the aid of a lens, one recognizes granulous parts affecting a certain regularity. The same dull white condition is seen on the fragments of the labial appendages that have remained adhering to them, hiding to some extent the brachial appendages. On examination of these fragmentary labial appendages with an enlargement of 20 to 30 diameters, one immediately observes calcareous spicula, the sharp extremities of which form a most elegant border, encroaching on the brachial membrane. These spicula were formed therefore from the beginning of the formation of the shell and, already very complicated, entirely resemble those seen in the adult individual The cirri and the channel of the labial appendages are enveloped by a layer of spicula spread out on the interbrachial

membrane in a sort of border or festoon Having afterwards submitted the minute shell to the action of water, accompanied by a little caustic potash, I was able to isolate the brachial appendages without effecting any fracture. The calcareous appendage, or loop, is formed at this stage of growth by two little short calcareous processes only, which represent the origin of the crura and offer no traces either of the principal stems or of the transverse connecting lamina of the loop. In this first stage, the brachial appendages entirely resemble those of a *Rhynchonella*; and if one limited oneself to a superficial examination one would be more disposed to take the embryo of *Liothyris vitrea* for a minute *Rhynchonella*, the then triangular aspect of the foramen and the pointed beak heightening the illusion I next examined different examples of 6 and 7 millimètres in length; the shell then, although still quite young, had completely changed in aspect, its shape being essentially the same as in the adult condition. The loop occupies about a fourth of the length of the shell, and is complete in all its parts. I next examined specimens of 10 millimètres in length, in which the only differences, and very slight ones, were limited to the loop being a little broader anteriorly, and it never afterwards assumed any difference. Therefore the study of *Liothyris vitrea* offers three facts of great general importance, namely:—

“1. The brachial apparatus or loop follows in its development a regular progression. It is at first as simple as possible, composed of two little branches which unite afterwards so as to form a small apparatus in the shape of a crest.

“2. This apparatus once formed does not undergo any metamorphosis, and does not pass through the complications that one observes in *Terebratella*.

“3. As soon as the labial appendages or arms have developed themselves, and the brachial apparatus or loop has commenced to be formed, the mantle and the arms present in the interior a very complicated system of calcareous spicula, especially destined to protect the channel of circulation, whilst in the *Terebratulæ* with long loops, *Waldheimia*, and *Terebratellæ*, one can discover not a trace of similar spicula.”

The soft parts of the animal of *Liothyris vitrea* are very similar to those of *Terebratulina caput-serpentis*, to which we allude in the sequel.

It is to be regretted that the anatomy of *L. vitrea* has not yet been published, and still remains a desideratum. The intimate shell-structure of *L. vitrea* has been minutely described and illustrated by Van Bemmelen in his memoir on the anatomy of the Brachiopoda*; the circular perforations, or canals, are widely separated from each other (as may be seen in the figure) although very small on the surface of the shell itself. Malformations in *L. vitrea* are not common. M. E. Deslongchamps has, however, described and illustrated a very remarkable one (Plate I. fig. 10 of this work) in which there exists a large longitudinal septum in both valves, which has been caused by an accident similar to that which caused the formation of the hole in *Terebratula diphya*.

Liothyris vitrea is a common fossil in the Pliocene rocks of Sicily, and occurs at Trapani, Tremonte, Gravitelli, and also at Terreti, near Reggio, in Calabria.

The shell referred to *L. vitrea*, by Chemnitz, in his Neues Conch. Cab. p. 97, tab. 78,

* Over den Bouw der Schelpen van Brachiopoden en Chitonen, 1882.

fig. 707, 1785, cannot surely belong to Born's species, for he figures in it a longitudinal septum, which never occurs in that species.

Several varieties of *L. vitrea* have received distinctive names.

LIOTHYRIS VITREA, var. *MINOR*, Philippi. (Plate I. fig. 13.)

Terebratula vitrea, var. *minor*, Philippi, Enumeratio Moll. Siciliae, vol. i. p. 99, pl. vi. fig. 8, 1836; vol. ii. p. 66, 1844.

Terebratula affinis, Calcara, Cenno sui Molluschi viventi e fossili di Sicilia, p. 48, 1845.

Terebratula vitrea (pars), A. Aradas, Descrizione delle conchiglie fossili di Gravatelli presso Messina, p. 14, 1847.

Terebratula minor, E. Suess, Ueber die Wohnsitze der Brachiopoden, 1859; Davidson, On recent Terebratulæ, Ann. & Mag. Nat. Hist. 3rd ser. vol. viii. p. 35, 1861.

Terebratula affinis, G. Seguenza, Notizie succinte intorno etc. pp. 19, 26, 32, 1862; id. Sulla formazione mioc. di Sicilia etc. p. 7, 1862.

Terebratula minor, Davidson, Outline of the Geology of the Maltese Islands by Dr. Leith Adams; and Description of the Brachiopoda by Thos. Davidson, Ann. & Mag. Nat. Hist. 3rd ser. vol. xiv. p. 8, pl. i. fig. 8, 1864; Seguenza, Memorie della Soc. Ital. di Scienze Naturali, vol. i. p. 21, 1865; Kowalevsky, Observations on the Development of the Brachiopoda (in Russian). Nachr. Ges. Mosc. xiv. 1873.

Shell small, ovate, longitudinally oval, longer than wide, about 9 lines in length by 6 in breadth and $5\frac{1}{2}$ in depth. Valves uniformly convex and smooth, no fold or sinus; ventral valve somewhat deeper than the dorsal one; beak moderately incurved, and truncated by a small circular foramen slightly separated from the hinge-line by a narrow deltidium in two pieces; loops short and simple. Colour pale yellowish white.

Hab. Living in the Straits of Messina and off the Eolic Islands in the Mediterranean; off Cape of St. Vincent ('Talisman' Expedition), in 298–818 fathoms.

Obs. Considerable uncertainty has prevailed with respect to this small shell, which Philippi, in 1836, described from fossil Pliocene specimens as a variety of *Terebratula vitrea*. Suess and others declare it to be a distinct species, while others look upon it as a young stage of *Liothyris vitrea*. I have compared species of the fossil shell with species dredged alive by Prof. Seguenza in the bay of Messina, and found them to be identical. I cannot, however, get rid of the idea that *Liothyris minor* is more than a small race or variety of *Liothyris vitrea*; it occurs, associated with the last-named shell, in the same beds and localities in Calabria and in Sicily.

To Kowalevsky's memoir on the development of the Brachiopoda (1873) the reader is referred for observations relative to the embryology of *L. minor*.

LIOTHYRIS VITREA, var. *DAVIDSONI*, A. Adams. (Plate I. figs. 14–16.)

Terebratula davidsoni, A. Adams, Proc. Zool. Soc. London, p. 314, 1867.

Terebratula minor, Davidson, Proc. Zool. Soc. London, p. 302, pl. xxx. fig. 10, 1871.

Shell ovate, broadest anteriorly, tapering posteriorly, longer than wide, marginally rounded. Valves uniformly convex, no fold or sinus, surface smooth, with faintly marked concentric lines of growth; beak incurved, truncated by a small circular foramen margined laterally by two small deltidial plates; surface of shell finely punctured.

Colour light yellowish white; loop small and simple. Length 10, width 7, depth 6 lines.

Hab. Dredged by A. Adams at Satanomosaki, Japan, in 55 fathoms.

Obs. I am not quite certain that this small species is really a variety of *L. vitrea*. I have seen only two examples of the shell, and they much resemble the typical var. *minor*. One of the specimens bore some resemblance to young examples of *Liothyris uva* from the Gulf of Tehuantepec, but differs from it, according to A. Adams, in its more solid structure and globose form, and in the foramen being smaller and entire. More Japanese examples will have to be examined before the variety can be definitely accepted.

2. *LIOTHYRIS ARCTICA*, Friele, sp. (Plate I. figs. 17, 18.)

Terebratula arctica, Friele, Særskilt Aftryk af Nyt Magazin for Naturvidenskaberne, pl. i. fig. i., 1877.

Shell small, globose, broadly ovate, rather longer than wide. Valves smooth, glassy, semitransparent, whitish; dorsal valve convex, squarely circular, without fold or sinus; ventral valve very convex and deep; beak unusually short, slightly incurved and truncated by a very small foramen margined anteriorly by rudimentary deltidial plates; loop very small and simple. Length 7, breadth 6, depth 4 lines.

Hab. Dredged by Herman Friele some few miles south-west of Jan Mayen, in 263 fathoms depth. Shell abundant, but so brittle that most of the specimens were broken during the dredging-operation.

Obs. After having carefully compared a specimen of the shell under description, sent to me by Friele, with others of the var. *minor* to which it had been referred by Dr. Jeffreys, I could, as Friele had previously done, discover several differences which, although not very great, have induced me to follow its discoverer in considering it a distinct species. *L. arctica* is much more globose and squarely rounded than *L. minor*, which is more of an elongated oval. As stated by Friele, its form approaches most to *L. minor* of Philippi, but the deviation is shown in the shorter beak and by the position of the foramen, which, in *L. arctica*, is placed directly above the dorsal valve, the deltidium being almost hidden. The loop in *L. arctica* is very much weaker and thinner, and the crura processes are placed further apart than in *L. minor*. It is the first representative of the genus *Liothyris* that has been hitherto found in Arctic seas.

3. *LIOTHYRIS UVA*, Broderip, sp. (Plate II. figs. 5-7.)

Terebratula uva, Broderip, Trans. Zool. Soc. Lond. vol. i. p. 142, pl. xxii. fig. 2, 1833; Sowerby, Thes. Conch. vol. i. p. 353, pl. lxx. figs. 53-55, 1846; Reeve, Monogr. of the genus *Terebratula*, Conch. Icon. pl. iii. fig. ii., 1860; Dall, Cat. of the recent species of the Class Brachiopoda, Proc. Acad. of Nat. Sciences of Philadelphia, 1873; Davidson, Report on the Brachiopoda, Voyage of H.M.S. 'Challenger,' Zool. vol. i. p. 31, pl. ii. figs. 3 & 4, 1880.

Shell oblong-oval, posteriorly compressed on each side, longer than wide, slightly diaphanous, white or of a very light salmon-colour. Dorsal valve convex; marginal

line flexuously rounded in front ; ventral valve deeper than the dorsal one ; beak incurved, moderately produced, obliquely truncated by a large circular foramen separated from the hinge-line by a narrow concave deltidium. Surface smooth, marked by fine concentric lines of growth. Shell finely punctated. Loop short, simple. Length 1 inch 1 line, width 8 lines, depth 6 lines.

Hab. Mr. Broderip states that this species was obtained at Tehuantepec by Capt. Dare, while dredging for *Meleagrina margaritifera*, attached to a dead sea-worn bivalve, at a depth of 10–12 fathoms, and on a bottom of sandy mud. The type, formerly in Mr. Cuming's collection, is now in the British Museum. The shell was also trawled by the 'Challenger' Expedition off Twofold Bay, in 120 fathoms, and dredged off Buenos Ayres by the same expedition at a depth of 600 fathoms, and again off Heard Island, lat. 52° 4' S., long. 71° 22' E., at a depth of 150 fathoms.

Obs. *Liothyris uva* varies much in shape ; it is usually longer than wide, and oval, but in some examples the length and depth do not differ materially.

4. *LIOTHYRIS MOSELEYI*, Davidson. Plate II. figs. 1–4.

Terebratula Moseleyi, Davidson, Proc. of the Royal Society, vol. xxvii. p. 436, 1878 ; Report on the Brachiopoda, Voyage of H.M.S. 'Challenger,' Zool. vol. i. p. 30, pl. ii. figs. 12–14, 1880.

Shell broadly ovate, semiglobose, rather longer than wide, broadest anteriorly, slightly tapering posteriorly, marginally and laterally convex, nearly straight in front, margin sometimes thickened, surface smooth, white. Dorsal valve uniformly convex, without fold or sinus ; ventral valve slightly deeper than the dorsal one, uniformly convex ; beak moderately produced, slightly incurved and truncated by a circular foramen separated from the hinge-line by a very narrow and small deltidium, beak-ridges not defined. Loop in dorsal valve short and simple, labial appendages occupying about two thirds of the length of the valve, united to each other by a membrane, the central coil making about three turns. Shell-structure perforated by numerous small canals. Colour white. Length 11, breadth 10, depth $7\frac{1}{2}$ lines.

Hab. Dredged by the 'Challenger' Expedition, west of Kerguelen Island, at a depth of 210 fathoms. Types in the British Museum.

Obs. I have reproduced the description I have given in the 'Challenger' Report. Five examples of this shell were obtained, all of about the same dimensions. It seems to be a smaller and more circular species than *L. vitrea* and *L. sphenoidea*. It is less elongated, and not quite so convex as the last-named species, and does not present the flatness and angularity observable in the mesial and lateral portions of the ventral valve of *L. sphenoidea* or in its synonym *L. cubensis*.

I made an examination of the animal of one of the specimens, which did not differ materially from that of *L. vitrea*. The mantle is thin, and not furnished with setæ at its edges. On the dorsal lobe of the mantle I distinctly observed the ramified, bifurcated, fine, thread-like pallial nerves as well as the pallial sinuses, muscles, and brachial or labial appendages, these last occupying a much smaller space in the interior of the shell ; and while the labial branches are visibly shorter, the cirri are of considerable length.

5. *LIOTHYRIS SPHENOIDEA*, Philippi, sp. (Plate II. figs. 17-22.)

Terebratula sphenoidea, Philippi, Enum. Moll. Siciliae, vol. ii. p. 68, tab. xviii. fig. 6, 1844; G. Seguenza, Pal. Malac. dei Terreni Terziarii del distretto di Messina, Memorie della Soc. Italiana di Scienze Naturali, p. 24, pl. ii. figs. 1-5, 1865; also Studii Paleontologici sui Brachiopodi dell' Italia Meridionale, pl. i. figs. 18-26, Pisa, 1871.

Terebratula cubensis, Pourtales, Contributions to the Fauna of the Gulf Stream at great depths, Bull. Mus. Comp. Zoöl. vol. i. pp. 109 & 124, 1867; Dall, Report on the Brachiopoda obtained by the United States Coast Survey Exp., Bull. Mus. Comp. Zoöl. vol. iii. pp. 3-9, pl. i. fig. 2, 1871; Davidson, Report on the Brachiopoda, Voy. H.M.S. 'Challenger,' p. 28, pl. ii. figs. 10-11, 1880; Dall, Bull. Mus. Comp. Zoöl. vol. ix. p. 103, 1881.

Terebratula vitrea, var. *sphenoidea*, Jeffreys, On the Mollusca procured during the 'Lightning' and 'Porcupine' Expeditions 1868-70, Proc. Zool. Soc. p. 404, pl. xxii. fig. 6, 1878.

Shell longitudinally more or less trigonal, broadest and rounded anteriorly, tapering posteriorly, lateral marginal line flexuous, that of the dorsal valve forming an outward curve, flexuously varying with age. Dorsal valve uniformly convex, sometimes rather inflated; ventral valve somewhat deeper than the dorsal one, longitudinally broadly flattened, sides of the flattened portion sloping away rather abruptly on either side, giving the valve a somewhat subquadrangular aspect; beak moderately incurved and truncated by a circular foramen separated from the hinge-line by a narrow deltidium. Surface smooth, sometimes marked by fine radiating lines. Loop narrow, small, and simple, bent-up band connecting the principal stems of the loop long and narrow. Colour soiled white. Length 1 inch 3 lines, breadth 11 lines, depth 9 lines.

Hab. Recent. Atlantic (Jeffreys). West-African coast ('Travailleur' and 'Talisman' Expeditions). Gulf of Florida in depths of 100-200 fathoms, rarer towards east end of reefs (Pourtales). Coast of Cuba (Sigsbee), off Havana 270 fathoms. Barbados 100 fathoms, St. Vincent 88 fathoms, Martinique 210 fathoms. Off Ascension 420 fathoms ('Challenger' Expedition). Off Morocco ('Talisman' Expedition) in 298 to 818 fathoms.

Fossil. In Pliocene rocks of Calabria and Sicily (Philippi and Seguenza).

Obs. In 1844, Philippi described and figured, as his *Terebratula sphenoidea*, some fossil Pliocene specimens he had collected in the valley of Lamanto in Calabria. Subsequently Signor Seguenza found the same fossil in Philippi's locality, as well as in rocks of the same age in Sicily. In 1878, Dr. Gwyn Jeffreys described and figured as *Terebratula vitrea*, var. *sphenoidea*, a living specimen which he had dredged during the 'Porcupine' Expedition in 1870, from the Atlantic, at depths of 292, 374, and 994 fathoms, and remarks:—"After a protracted and very careful examination of my specimens, which I had considered the *T. sphenoidea* of Philippi, and having compared them with fossil specimens sent me by Prof. Seguenza as Philippi's species from the Sicilian Tertiaries, as well as with a series of *T. cubensis* which I received from Count Pourtales and Professor Alexander Agassiz, and also after a close comparison of all these specimens with the description and figures given by Philippi, Seguenza, Pourtales, and Dall, I am convinced the *T. sphenoidea* and *T. cubensis* are the same, and constitute a well-marked variety of *T. vitrea*. The loop in *T. sphenoidea* and *T. cubensis* is precisely similar." In this last remark, Dr. Gwyn

Jeffreys is clearly mistaken; for there exists a well-marked difference between the loops of *T. sphenoidea* and *T. cubensis* and that of *L. vitrea*. The loops in the fossil specimens of *Liothyris sphenoidea* and those of the recent *L. cubensis* are exactly the same, as had been clearly illustrated by Philippi in 1844.

As justly remarked by Prof. Dall, *L. sphenoidea* varies much in shape according to age and specimen: some are longer than wide, others almost as wide as long; some taper more than others posteriorly, while much difference is observable in the degree of convexity of their valves.

In his description of *T. cubensis*, Dall enters into minute details, in order to point out the differences which exist between this last-named species and *Liothyris vitrea*. He has also carefully studied the animal of *T. cubensis*, and points out the differences it presents from that of *Waldheimia floridana*. He says, "the mantle is of stouter consistency than in *W. floridana*, and may often be removed from the shell with but little injury if care be exercised. The muscles are similar in disposition to those of the other members of the Terebratulidæ, and present no new features. The peduncle is solid, cup-shaped at its extremity, and has the edge produced in cylindrical horny rootlets, which are attached to foreign bodies. The regular arrangement in layers of the muscles and corium, as well as the axial tube of the peduncle, found in *Lingula*, is less evident or absent in these forms. In this species the peduncle is very short and stout, broadly cordiform at its inner extremity when enveloped by its various tunics.

"The brachia are arranged as in *T. vitrea*, as figured by Woodward; the central coil makes about four turns. The cirrhi are very short behind the mouth, in front of the supra-oesophageal body. A striking feature in its anatomy, which I believe has not yet been noted in any publication on Brachiopods, is the absence of that great series of sinuses in the anterior part of mantle, which was termed by Hancock the 'great pallial sinuses.' So extraordinary did this appear to me, that I could not believe, at first, that I was not deceived by the translucency of the membranes, and it was only after an examination of many specimens that I became convinced that they do not exist in this species. There is in the free lobes of the mantle an extensive and extremely close and fine network of minute channels; or perhaps it might be said that the whole of the mantle-lobes form one great lacune, the upper and lower walls of which are held apart by a profuse number of pillars of tissue, which appear like dark spots under the microscope, and which are situated so close together that the spaces about them are reduced to minute channels. This system occupies the anterior lobes of the mantle, which in some species also contain large branching sinuses, here absent. . . . In the inner lining of the mantle are scattered, everywhere, delicate, branching spiculæ, looking more like briars than like deer-horns, and, while more or less interlocked, and here and there stout and thick, are still much more delicate and slender than those of *Terebratulina caput-serpentis* and *Megerlia truncata*, and do not often exhibit a stellar arrangement. They are much more numerous in some individuals than in others, and when present in abundance are found in almost every part of the epithelium, even to the brachial cirrhi, where the spicules are slender and not branched. . . . The oesophagus is wide and funnel-shaped, narrowest at its junction with the stomach, which it enters at an acute angle. The stomach is

small and oval, tapering towards the intestine, which is nearly twice as long as the œsophagus.

“The heart in most specimens was pyriform and of a moderately large size.

“The genitalia . . . are situated in a reticulated series of sinuses, on the surfaces of the sides of the perivisceral tissues.

“Above and behind the mouth, and directly in front of the anterior oclusor (retractor) muscles, the external tissues of the perivisceral membrane are thickened, or a mass of cellular tissue is interposed between the laminæ of the membrane.

“No peculiarities of note were observed in the shell-structure. The perforations appeared to be slightly further apart than in *T. vitrea*, but the difference was not much greater than that which may be observed in the shells of different individuals of the same species.”

Mr. Dall then describes what he believes to be the young of *T. cubensis*, dredged off the Samboes, on the Florida reefs, a minute, polished, hyaline shell, 4-100ths of an inch in length, and follows by describing the muscular system, brachia, and organs of digestion.

Liothyris sphenoides and its synonym *T. cubensis* appear to be very abundant in their especial haunts. I have also been able to examine the animal from one of the specimens dredged by the ‘Challenger’ Expedition. The mantle in the dorsal valve of one of the specimens showed in a most distinct manner the four principal pallial sinuses, which again branched as they approached the front margin of the shell, and bifurcated again before reaching the margin.

6. *LIOTHYRIS BARTLETTI*, Dall, sp. (Plate I. figs. 20, 21.)

Terebratulula bartletti, Dall, The American Naturalist, vol. xvi. p. 885, Nov. 1882.

Shell ovate, globose, longer than wide, broadest anteriorly, dorsal valve convex, with a wide flattened mesial fold of very small elevation, commencing at about two thirds of the length of the valve and extending to the front. Lateral and frontal margins sinuous, front line of fold nearly straight. Ventral valve slightly deeper and more convex than the dorsal one, with a wide, shallow, flattened, mesial depression or sinus near the front; beak short, much incurved, overlying the umbo of the opposite valve, and truncated by a small, oval-shaped, incomplete foramen. Surface smooth, marked with concentric lines of growth. Colour light yellowish brown. In the interior of the dorsal valve the loop is short and simple. Length $1\frac{1}{2}$ inch, breadth 1 inch 2 lines, depth 1 inch.

Hab. Dredged by the United States Coast Survey. Gulf Stream (‘Blake’ Expedition) near Vera Cruz, in 218 fathoms.

Obs. I have seen one specimen only of this species, kindly lent to me by Mr. Dall. It approaches much in general shape to some specimens of *Liothyris vitrea*. It differs from *Liothyris sphenoides* (= *cubensis*) in the shape of its loop, which is similar to that of *L. vitrea*.

7. *LIOTHYRIS SUBQUADRATA*, Jeffreys, sp. (Plate II. figs. 15, 16.)

Terebratulula subquadrata, Jeffreys, On the Mollusca procured during the ‘Lightning’ and ‘Porcupine’ Expeditions in 1868–70, Proc. Zool. Soc. London, 1878, p. 402, pl. xxii. fig. 3.

Shell somewhat subpentagonal or pear-shaped, broadest anteriorly, tapering posteriorly, longer than wide, rounded laterally, slightly indented in front. Dorsal valve moderately and evenly convex, somewhat flattened along the middle. Ventral valve deeper than the dorsal one; beak slightly incurved and moderately produced, obliquely truncated by a rather large circular foramen, separated from the obtusely angular hinge-line by a well-defined triangular area and long, narrow, deltidial plates; beak-margins sharply defined. Surface of valves marked with numerous, wavy, fine, radiating riblets, widely separated and crossed at intervals by concentric lines of growth. Colour ochreous white, cæcal tubuli minute and close-set. Loop small and simple. Length 1 inch 1 line, breadth 1 inch, depth 7 lines.

Hab. Off the coast of Portugal, in 500–600 fathoms; in the Bay of Biscay.

Obs. The only complete specimen of this shell that I have seen is the one described, which I figured for Dr. Jeffreys. It was given to me by Mr. Saville W. Kent, who had dredged it during his cruise in Mr. Hall's yacht 'Norna' off the Setubal coast, near the Tagus in 1870. Since then I have been informed by Dr. Jeffreys that young examples were obtained during the dredgings of the French ship 'Travailleur' in the Bay of Biscay. It seems to be a good and well-marked species.

8. *LIOTHYRIS WYVILLII*, Davidson. (Plate II. figs. 8–14.)

Terebratula Wyvilli, Davidson, Proc. Roy. Soc. vol. xxvii. p. 436, 1878, and *T. wyvillii*, Report on the Brachiopoda, Voyage of H.M.S. 'Challenger,' Zoology, vol. i. p. 27, pl. ii. figs. 7, 8, 1880.

Shell somewhat subpentagonal, variable in shape, about as broad as long, sometimes almost square, with rounded angles, with a concave sinus on the dorsal valve and corresponding fold in the ventral one. Shell very thin and exceedingly brittle, almost transparent, smooth, glassy, light-yellowish white. Length 7, breadth 9, depth $4\frac{1}{2}$ lines. Valves in the young slightly and evenly convex; dorsal valve moderately convex, with a depression of greater or less depth commencing close to the umbo, and gradually widening and deepening as it nears the front, front line wide, straight, or presenting an inward curve. Ventral valve deeper and more convex than the opposite one, with a wide *median convex* elevation or fold commencing near the beak, and one extending to the front. Beak very small, slightly incurved, truncated by a small, generally incomplete, circular foramen, laterally margined by deltidial plates. Surface of valves marked at intervals by concentric lines of growth; shell-structure with minute, widely separated perforations or canals. In the interior of the dorsal valve the loop is short and simple; the adductor and other muscular impressions are small and delicate. The labial appendages extend to two thirds of the length of the shell. In the interior of the ventral valve the muscular impressions are small, and occupy a limited area close to the beak.

Hab. This remarkable and very interesting species appears to abound over a wide geographical area, and at depths from 1035 to 2900 fathoms.

It was dredged by the 'Challenger' Expedition at six or seven different stations. In South Australia, in lat. $42^{\circ} 42'$ S., long. $134^{\circ} 10'$ E., depth 2600 fathoms: one example was attached to a manganese nodule. In lat. $12^{\circ} 8'$ S., long. $145^{\circ} 10'$ E., depth 1400 fathoms; bottom-temperature $1^{\circ} 3$ C. In lat. $33^{\circ} 31'$ S., long. $74^{\circ} 43'$ W., depth 2160

fathoms. Off coast of Chili (Valparaiso), along with *Waldheimia Wyrillii* and *Discinisca atlantica*. In lat. 42° 43' S., long. 82° 11' W., depth 1450 fathoms. Off coast of Patagonia. One small example from the net-weights, not far from Falkland Islands, at a depth of 1035 fathoms. And, lastly, in lat. 35° 22' N., long 169° 58' E., depth 2900 fathoms, the greatest depth at which any Brachiopod was obtained by the 'Challenger' Expedition; bottom-temperature 1°·2 C.; sea-bottom, red clay.

Obs. *Liothyris Wyrillii* is one of the most interesting species of deep-sea Brachiopoda. The shell is of such extreme thinness that it is almost transparent; indeed the valves, when separated, are really so, and the muscular impressions may be seen through its transparency. I separated the valves of a specimen in order to be able to study the animal and its loop. The latter, which I was much surprised to find short, is exactly similar to that of *Liothyris vitrea*, notwithstanding the outward *Waldheimia*-like appearance of the shell. It bears also much resemblance to several species of the last-named genus occurring in the Jurassic and Cretaceous formations. We meet with but few recent species with such a thin shell; but among others may be named the widespread *Discinisca atlantica*, King, *Atretia gnomon*, Jeffreys, *Waldheimia* or *Macandrevia tenera*, Jeffreys, *Rhynchonella lucida*, Gould, and one or two others.

In external shape and character of loop, *Liothyris Wyrillii* also strongly resembles the *Terebratula nucleata* of Schlotheim. Zittel's figure 544, on page 700 of his 'Handbuch der Paläontologie,' 1880, seems as if drawn from a specimen of the recent *L. Wyrillii*. For *T. nucleata* Douvillé proposes a genus *Glossothyris*; but this I am unable to admit or adopt.

Uncertain Species.

9. *LIOTHYRIS CERNICA*, Crosse, sp. (Plate I. fig. 19.)

Terebratula cernica, Crosse, Journal de Conchyliologie, vol. xxi. p. 285, and vol. xxii. p. 75, pl. i. fig. 3, 1873.

Shell longitudinally oval, somewhat pear-shaped, longer than wide, broadest anteriorly, tapering posteriorly. Valves uniformly convex and globose, lateral margins rounded and flexuous. Surface smooth, white, semitransparent and somewhat vitreous. Beak not much produced, incurved and truncated by a small circular foramen, separated from the hinge-line by a deltidium in two pieces. Loop not known, in all probability short and simple, as in *L. vitrea*. Length 1 inch 4 lines, breadth 1 inch 1 line.

Hab. Off Mauritius Island (Liénard).

Obs. Mr. Crosse, in his description, says that *T. cernica*, from the lateral compression of its dorsal valve, possesses affinities with *T. ura*, Broderip, but that it is less elongated and more globose, and approaches *T. vitrea* and, even more closely, *T. cubensis* of Pourtales. I have never seen the single example of the shell under description; but from inquiries I made in the Island of Mauritius, Mr. V. Robillard has informed me that it has not been dredged, but was found in the stomach of a fish taken at 80 fathoms depth, that it is the only specimen existing, and that there is but little chance of finding it again. He adds that he does not know of any other species of Brachiopod from the waters of

Mauritius, although he has been studying conchology there for the last forty-five years.

10. *LIOTHYRIS* ? vel *TEREBRATULINA* *DALLI*, Davidson. (Plate II. fig. 23.)

Terebratula ? *Dalli*, Davidson, Proc. Roy. Soc. vol. xxvii. p. 437, 1878; and Report on the Brachiopoda, Voyage of H.M.S. 'Challenger,' Zool. vol. i. p. 38, pl. ii. figs. 15, 15 *a, b*, 1880.

Shell small, thin, longitudinally oval, globose, glassy, semitransparent. Dorsal valve moderately convex, slightly depressed anteriorly. Ventral valve uniformly convex, a little deeper than the dorsal one. Beak small, slightly incurved, and truncated by an incomplete foramen, laterally margined by small deltidial plates. Surface of valves covered with fine, radiating, raised striæ, with shorter ones here and there, interpolated between the longer ones. Loop short and simple. Length 3, width 2 lines.

Hab. One example only, without the animal, was dredged by the 'Challenger' Expedition near Yeddo, off Japan, lat. 34° 37' N., long. 140° 32' E., depth 1875 fathoms.

Obs. I am not quite certain whether this small shell is a *Liothyris* or a *Terebratulina*. Its generic claims must be considered uncertain.

Subgenus *TEREBRATULINA*, d'Orbigny, 1847.

The subgenus *Terebratulina* is closely allied to *Liothyris*. When young, and up to a certain stage of its development, the loop is similar to that of *Liothyris*; but with age the crural processes become united and form a shelly band, which is never the case in *Liothyris*. Apart from the last-named peculiarity of the ring-shaped loop, the subgenus *Terebratulina* well defines a small group of shells distinguished and characterized by the presence of ear-shaped expansions on each side of the umbo, and by the fine radiating striæ that cover the surface of their valves.

11. *TEREBRATULINA* *CAPUT-SERPENTIS*, Linné, sp. (Plate III. fig. 12; Plate IV. figs. 1-11; Plate V. figs. 32-34.) [Figs. 35-37, var. *emarginata*, Risso = var. *mediterranea*, Jeffreys. See footnote, p. 25.]

Anomia caput-serpentis, Linné, Syst. Nat. ed. duodecima reformata, vol. i. p. 1153, Holmiæ, 1767.

Anomia pubescens, id. ibid. p. 1153.

Anomia retusa, id. ibid. p. 1151.

Animal Anomiæ nondum antea depictum, Ad. Murray, Fundamenta Testaceologiæ, Upsala, p. 43, pl. ii. fig. 23, 1771.

Anomia caput-serpentis, Pennant, (Linné) Nova Acta Regiæ Societatis Upsaliensis, vol. i. p. 38, tab. v. fig. 4, 1773.

Terebratula caput-serpentis (animal), Grundler, Naturforscher, Bd. i. p. 84, tab. iii. figs. 1-6, 1774; Beschreibung und Abbildung zweier natürlichen Terebrateln, Naturf. Bd. ii. p. 80, 1774; Born, Musci Cæsarei Vindobonensis, p. 119, 1780.

Anomia caput-serpentis, Chemnitz, Conchylien-Cabinet, vol. viii. p. 103, tab. 78. fig. 712, 1785.

Terebratula pubescens, Retzius, Nov. Gen. xv., 1788.

Terebratula caput-serpentis, Dillwyn, Cat. Recent Shells, 1817; Lamarck, An. sans Vert. p. 247, 1819.

Terebratula aurita, Fleming, Phil. of Zool. ii. p. 498, tab. iv. fig. 5, 1822, and History of British Animals, vol. i. p. 369, 1828.

- Terebratula costata*, Lowe, Zool. Journal, vol. ii. p. 105, 1825.
- Terebratula emarginata*, Risso, Hist. Nat. de l'Europe Méridionale, vol. iv. p. 388, pl. xii. fig. 175, 1826.
- Terebratula quadrata*, Risso, ibid. p. 389, pl. iv. fig. 176, 1826.
- Delthyris spatula*, Menke, Synopsis Methodica Molluscorum, 2nd ed. p. 96, 1828-30.
- Terebratula caput-serpentis*, Philippi, Enum. Moll. Siciliæ, vol. i. p. 84, tab. vi. fig. 5, 1836; Anton, Verzeichniss der Conchylien, p. 23, 1836.
- Terebratula Gervillei*, S. V. Wood, Ann. & Mag. Nat. Hist. 1st ser. vol. vi. p. 253, 1840.
- Terebratula Chemnitzii*, Küster, Martini & Chemnitz, Conchylien-Cabinet, vol. vii. p. 37, pl. 2 b. figs. 19 and 20, 1843.
- Terebratula caput-serpentis*, G. B. Sowerby, Thes. Conch. vol. i. p. 343, pl. lxxviii. figs. 1-4, and pl. lxxii. fig. 116, 1846.
- Terebratulina caput-serpentis*, d'Orbigny, Ann. des Sci. Nat. vol. viii. p. 67, pl. vii. figs. 7 & 8, 1848.
- Terebratulina cornea*, d'Orbigny, 1848.
- Terebratula caput-serpentis*, Forbes and Hanley, Brit. Moll. pl. lvi. figs. 1-4, 1849.
- Terebratulina caput-serpentis*, Davidson, Sketch of a Classification of Recent Brachiopoda, Ann. & Mag. Nat. Hist. 2nd ser. vol. ix. p. 365, 1852; Davidson, Br. Tert. Brach., Pal. Soc. vol. i. p. 12, pl. i. figs. 3-6, 1852.
- Terebratula striata*, Leach, Br. Moll. 1852.
- Terebratulina striata*, S. P. Woodward, Manual of the Mollusca, p. 215, 1854; Hancock, Phil. Trans. vol. cxlviii., 1858.
- Terebratula striata*, L. Reeve, Conch. Icon. pl. iv. fig. 15, 1860.
- Terebratulina striata*, E. Deslongchamps, Recherches sur l'Organisation du Manteau chez les Brachiopodes articulés, 1864; Brusina, Moll. Dalmati, Verh. zool.-bot. Ges. Wien, Bd. xvi., 1866; Dall, Proc. Acad. Nat. Sciences of Philadelphia, p. 179, 1873; Kowalevsky, On the Development of the Brach., 1874; G. O. Sars, Bidrag til Kundskaben om Norges Arktiske Fauna, Mollusca Regionis Arcticæ Norvegiæ, tab. i. fig. 5, 1878; Jeffreys, Brit. Conch. vol. ii. p. 14, and vol. v. pl. xix. fig. 2, 1873-79; E. Deslongchamps, Études critiques sur des Brachiopodes nouveaux ou peu connus, 1884*.

Shell ovate, somewhat pentagonal, longer than wide, broadest about the middle, slightly rounded or indented anteriorly; dorsal valve more or less uniformly convex, but with sometimes a mesial longitudinal depression, marginally flexuous on the sides; lateral sides of the umbo auricular; ventral valve convex, rather deeper than the dorsal one, anteriorly mesially depressed; beak rather short, gently incurved and obliquely truncated by an incomplete foramen, posteriorly margined by the substance of the beak, anteriorly by the extremity of the umbo, laterally by two small deltidial plates, no distinct cardinal area or beak-ridges; surface of shell densely covered with fine radiating striæ, few in number, coarse and simple when young, but rapidly increasing in number with age, by means of bifurcation, and by the interpolation of smaller and shorter ribs; both valves are also crossed by numerous concentric raised lines of growth, more prominent in the young, finer and closer as the shell increases in size. Loop short and simple in the young, with age becoming annular through the union of the oral processes. Colour whitish, with a tinge of yellow, sometimes rust-stained. Dimensions variable; a large example measured 1 inch 3 lines, breadth 1 inch, depth $\frac{1}{2}$ inch.

* In addition to the above references, the species has been alluded to by many other conchologists and palæontologists, such as by Gualtieri, in his Index Conch. tab. 96. fig. 3, 1742; by Davila, Cat. i. pl. xx.; Favanne, Conch. pl. 41. fig. 4^a, 1780, and by many others.

Hab. Spitzbergen and Davis Straits; north-east European seas; Oban and off Cumbræ Islands, Loch Torridon, Scotland; off Belfast; Finisterre and Croix de Gavie; Morbihan; Cape Breton; Adventure Bank; off Guetaria, Spain; north-east coast of Jamaica; Corea and Sagami Bay, Japan, &c.

Fossil. In the Upper Tertiaries of Sicily; Coralline Crag of England, Belgium, south of Spain, Azores, &c., &c.

Obs. No species of Brachiopoda has been more thoroughly or more carefully studied than the *Anomia caput-serpentis* of Linné. It is a very common and well-known species, abounding in the localities where it is found, and has since 1767 attracted the attention of a large number of malacologists.

The shell varies somewhat in shape and especially, according to age, in the number of its ribs. These modifications have been often described by myself and others, and most recently, in 1884, by M. E. Deslongchamps*. In the youngest condition the shell tapers posteriorly, and is widest anteriorly; the hinge-line is then almost straight, the auricular expansions comparatively larger, the foramen triangular or elongated oval. In a specimen, less than a line in length, forwarded for my examination by the Marquis de Folin, the posterior half of the valves was smooth, while on the anterior half some seven, scarcely developed, rounded radiating ribs were present. At one and a half line the surface of each valve was ornamented with about ten simple rounded ribs with interspaces of about equal breadth, concentrically crossed by equidistant projecting ridges, most prominent on the surface of the ribs. As the shell grows larger, the auricular expansions become smaller, the hinge-line obliquely or obtusely angular, the ribs more numerous and finer, the lines of growth less prominent, and the greatest breadth at about half the shell's length. Similar important modifications take place also in the interior of the valves. In the interior of the dorsal valve, when quite young, and up to a certain age, the crura are widely separated; but as the shell grows, the pointed extremities facing each other extend nearer and nearer towards each other until they become united and form a well-defined band, giving the loop an annelliform shape and character.

The intimate shell-structure of *T. caput-serpentis* has been described and figured by Dr. W. B. Carpenter, Oscar Schmidt, Prof. W. King, E. Deslongchamps, and others. Dr. Carpenter says that it is in *T. caput-serpentis* "that the canals are of smallest dimensions, their largest diameter being about $\frac{1}{500}$ of an inch, whilst their average distance from each other is about the same as in the preceding case [*Waldheimia flavescens*, $\frac{1}{400}$ inch]—their regular arrangement, however, being so modified, that the external orifices are principally seen upon the elevated parts of the plications, whilst they open internally in similar rows."

When alluding to the shell-structure of *T. caput-serpentis*, in his valuable memoir on the histology of the test of the Palliobranchiata (Trans. Roy. Irish Acad. vol. xxiv. p. 448, 1869), Prof. W. King says:—"The species just mentioned, which is longitudinally ribbed, has two kinds of perforations: those lying in the furrows are simple; while those

* 'Études, critiques sur des Brachiopodes nouveaux ou peu connus.' Caen, 1884.

belonging to the ribs are antler-shaped, that is, twice, thrice, or oftener subdivided. Occasionally the perforations are confined to the ribs, which causes them to lie in bands. It is only by the most careful examination, while grinding down the ribs to the level of the intermediate furrows, that the true forms of the perforations can be determined. As the modification in question is a very important one in many respects, I have given two representations of it taken from a section prepared by Dr. Rowney. Fig. 10 shows a band of antler-shaped perforations in the simplest state, magnified 60 diameters: it would have been difficult to have represented them under the complex form they sometimes assume. Fig. 11 shows a portion of one of the perforations, magnified 210 diameters, having each termination of its branches furnished with a brush-like bundle, which in this species, as in *Terebratula vitrea*, is smaller than usual." [See woodcut, fig. 1 A & B.]

After W. King.

The shell-structure of *T. caput-serpentis* being so very remarkable, I asked my friend Mr. John Young, of the Hunterian Museum, Glasgow, to carefully examine the shell-structure of this species. He writes me on the 20th of April, 1885:—"The shell-structure of this species is of great beauty. The numerous canals incline from the lower margin of the shell upwards towards the beak, but on the ribs of the shell they radiate from the centre of each rib to either side, there being a central row of tubes or canals that point nearly straight upwards, while the other rows diverge. It is this bending inwards of the tubes on each side of the ribs towards the interior of the shell that caused Dr. Carpenter, in the introduction to your monograph, to state that they open internally in rows. Dr. Carpenter is, however, wrong when he states that the pores or canals are principally seen *on the elevated parts of the plications* (in this species), my etched specimens show that they open quite as numerous in the hollows between the ribs. It is only internally that the perforations gather together into rows with bare spaces between. On the exterior surface of the valves the pores are more evenly distributed, owing to the divergence of the tubes on the ribs. The perforations are not so well seen on this species, previous to etching. The tubes on their passage through the shell-surface often have a wave outward and upwards."

The embryology of *Terebratulina caput-serpentis* and of *T. septentrionalis* has been admirably investigated by Prof. E. Morse* and Prof. Kowalevsky†. Prof. E. Morse says:—"In the first stage the embryo becomes widened at one end. The segments are barely indicated, the posterior end is the widest, the anterior portion is ornamented with a conspicuous tuft of long cilia, so peculiar to the embryos of many worms. The embryo is also clothed with vibratile cilia, and in this condition slowly moves along the bottom of the dish without rising from it, or remains quiet. In the second well-marked stage the embryo is divided into two prominent segments; these expand and contract upon each other slightly, and the cephalic segment has the power of partially bending from side to side. In this stage the embryo is most active, swimming rapidly in every direction and turning

* "Embryology of *Terebratulina*," Mem. Boston Soc. Nat. Hist. vol. ii. pp. 251-264, 1873; and "The Systematic Position of the Brachiopoda," Proc. Boston Soc. Nat. Hist. vol. xv. pp. 315-372, 1873.

† 'Untersuchungen über die Embryologie der Brachiopoden:' Moskau, 1874.

abruptly about. The œsophagus also becomes dimly defined. In the third stage the peduncular segment is developed and projects from the posterior portion of what can now be called the thoracic segment. At this stage the embryo either remains immovable upon the bottom of the dish or slowly moves about. In two cases delicately barbed setæ to the number of thirty-five projected directly backward from the peduncular segment. The embryo is still clothed with cilia, though the long pencil of cilia has disappeared. The head is closely drawn to the thoracic segment, which becomes wider in transverse diameter, so as nearly to hide the peduncle. In the fifth stage the thoracic ring commences to fold, or turn upward upon opposite surfaces of its circumference, so as to gradually enclose the head; one fold being made slightly in advance of the other represents the larger or ventral valve. In this stage appear clusters of barbed and deciduous setæ upon the anterior margin, and in a later portion of this stage the first hardened areas of the dorsal and ventral plates make their appearance, and the cirri appear as blunted papillæ about the mouth. In the sixth stage the shell becomes rounded, the peculiar scaled structure makes its appearance, and the formation of tubules perforating the shell and of permanent setæ takes place." The author adds, further on, "that the eggs not only fill the large pallial sinuses, but hang in clusters from the genital band; from these parts they escape by dehiscence, and float freely in the perivisceral cavity The eggs are not uniform either in shape or size." Prof. Morse then describes in detail each of the stages; but space will not allow me to proceed further with his elaborate investigations, as they would demand numerous illustrations. (See Plate V. figs. 1-31.)

Dr. Gwyn Jeffreys states, in vol. ii. p. 7 of his 'British Conchology,' that in the fry-stage the little creature can creep and swim; but, "After quitting the embryonic state, they become invariably and permanently fixed to other substances, being incapable of any other motion than making a half turn round the peduncle or pivot."

Mr. Lucas Barrett, who had an opportunity of examining *T. caput-serpentis* in life, says * :—"This species shows more of itself than any other, and protrudes its cirri further; it was met with everywhere [near the coast of Norway], in small numbers, in 30 to 150 fathoms, often attached to *Oculina*. The cirri on the reflected part of the arms are shorter than those on the first part, as shown in the woodcut. The cirri were almost constantly in motion, and often observed to convey small particles to the channel at their base. When placed in a small glass of sea-water, the valves gradually opened. Individuals remaining attached to other objects manifested a remarkable power and disposition to move on their pedicles. Detached specimens could be moved about without causing the animal to close its valves. If any of the protruded cirri were touched, the cirri were retracted and the valves closed with a snap, but soon after opened again. When the oral arms are retracted the cirri are bent up, but are gradually uncoiled and straightened when the shell is opened, before which the animal has often been observed to protrude

Fig. 2.

T. caput-serpentis,
after L. Barrett.

* Ann. & Mag. Nat. Hist. 2nd ser. vol. xvi. p. 257, 1855.

a few of its cirri and move them about, as if to ascertain if any danger threatened. Only on one occasion a current was observed to set in on one side between the two rows of cirri. I had been attempting to ascertain the existence of currents, by introducing small quantities of indigo into the water surrounding the animal with a camel's-hair brush; three times the water was forcibly drawn in, and the particles of indigo were seen to glide along the groove at the base of the cirri in the direction of the mouth." See likewise a paper by Hérourard "Sur les courants de nutrition des Brachiopodes," Journ. de Conchyl. vol. xxv. p. 229, 1877.

Hancock * says:—"The pallial lobes extend forward, but do not project beyond the side of the body, where they become united, the junction being marked by a groove, bordered by a ridge on each side. They are very delicate and transparent, so that the great pallial sinuses can be distinctly traced, even to their terminal ramifications" (p. 793). "The arms of *W.* [*Waldheimia*] *cranium* and *T. caput-serpentis* are disposed in the same manner as in *W. australis*; and in the former the calcareous loop is precisely similar to that of the latter; but in *T. caput-serpentis* it is very much reduced, the extended lateral portions having almost entirely disappeared, little more than the transverse portion existing; and this, together with the crural processes, which are united below across the median line, forms a collar upon which the bases of the arms rest. In this species, therefore, the expanded lateral portions of the arms are without any apophysary support, and accordingly other means are provided for sustaining them. The two produced lobules of the dorsal pallial lobe reach to the ends of these portions of the arms as in *W. australis*; and are stiffened with numerous, imbedded, calcareous spicula, to such an extent, that when the soft tissues are removed by maceration the form of the parts remains unaltered. The spicula extend also over the surface of the inner lamina of the pallial lobe, and pervade likewise the walls of the canal, and even the cirri; so that the brachial apparatus becomes firmly fixed, and in this way a substitute is found for the usual apophysary support." (*Loc. cit.* p. 808.)

It is, however, to Oscar Schmidt, so far as I am aware, that we are indebted for the first notice of these remarkable spicula †, and in the *Annals & Mag. of Nat. Hist.* 2nd ser. vol. xvi. p. 439, pl. x., I gave a translation, with figures, of Schmidt's observations. The author remarks "that the mantle, oral arms, and cirri in *Terebratulina caput-serpentis* contain an innumerable number of calcareous plates, generally flattened, dilated and irregularly denticulated, situated in close vicinity to each other. It is easily conceived [he adds] that these calcareous masses stiffen the parts which contain them, and seem particularly to serve this function in the hollow cirri, thus preventing their sides from sinking down."

We are also indebted to Prof. E. Deslongchamps for an admirable memoir 'Recherches sur l'Organisation du Manteau chez les Brachiopodes articulés' (Caen, 1864), in which

* Phil. Trans. vol. cxlviii., 1858.

† "Die neusten Untersuchungen über die Brachiopoden von Owen, Carpenter und Davidson, mit einigen Zusätzen," Zeitsch. f. gesammten Naturwissenschaften, p. 325, 1854. In 1856, Dr. S. P. Woodward exhibited, at a meeting of the Zoological Society, the spicula in the pallial lobe of *T. caput-serpentis* (Proc. Zool. Soc. p. 368, 1856).

he minutely describes and illustrates the arrangement of the spicula in *Terebratulina caput-serpentis* as well as in other genera in which they occur (see Pl. IV. figs. 10, 11). He tells us that each of the lobes of the mantle in this species has four vascular sinuses, of which the lateral ones are more developed than the median two, which last contain in their interior the organs of reproduction. The spicula cover these sinuses, and afterwards line the walls of the visceral cavity, producing two large convex surfaces, which become united and thus completely close this cavity. In the dorsal valve the mantle reflects itself on the brachial apparatus, formed here of calcareous laminae bent in the shape of a ring, and spread themselves on the labial appendages, of which they line the interior of the great canal, as well as the cirri; the cirri, he adds, which I have also carefully examined in many specimens, are stated by Hancock to be large, much-branched, colourless, glass-like, and pellucid, somewhat like the antlers of a deer, only the branches are all on the same plane and are flattened and depressed a little; that they are, however, frequently much complicated, forming a central network with irregular radiating branches; in others, again, the branches pass from a single network centre. The spicula lie in the outer layer of the inner lamina, and are crowded to such a degree that the edges of the branches are almost in contact, thus forming an extensive though incomplete network of calcareous matter over the trunks of the great pallial sinuses; elsewhere they are sparingly distributed towards the margin of the lobes, where they are rather numerous. The brachial disk is trilobed, the central lobe elongated and spirally convoluted.

In 1771, in the 'Fundamenta Testaceologiae' it is represented as the animal of *Terebratulina caput-serpentis* without its shell. In 1773, Pennant describes the same species, and gives a figure of its labial appendages. Grundler, again, in 1774, devotes a whole plate to enlarged figures of the same appendages, and in two of them points out the position of the loop. Since that period they have also been described and illustrated by several malacologists, and more particularly by Albany Hancock, in his classical memoir 'On the Organization of the Brachiopoda' (1858). Mr. Hancock informs us, at p. 793 of his work, that, "To facilitate the examination of the soft parts, it is necessary to reduce the valves by dissolving them in dilute acid. When the calcareous matter has entirely disappeared, and the calcified shell-membrane has been removed, the transparent tegumentary envelope of the animal is exposed to view; and this, when placed in water, assumes the form of the shell. Towards the posterior or umbonal region the enlarged extremities of the shell-muscles are always apparent, clustered together on each side of the antero-posterior line, forming reddish-coloured patches."

The muscles naturally divide themselves into two groups, the valvular and those for adjusting the shell to the peduncle. As the arrangement of the muscles is stated by Hancock to be essentially the same in all the articulated Brachiopoda, and they have been described further on in *Waldheimia flavescens*, it will not be necessary here to refer further to those details, although it may be added that Hancock mentions an important modification as occurring in *T. caput-serpentis*, which consists in the large and powerful dorsal adjustor muscles not being attached to the hinge-plate, as in *W. flavescens*, but having their insertions in the valve itself. The posterior extremities of these muscles are seen on the

surface of the animal, on each side of the median line, elongated in the antero-posterior direction, and extending between the oclusors, almost as far forward as their anterior margins. They are of irregular form, and enlarged posteriorly in *T. caput-serpentis*.

In Pl. IV. figs. 8 & 9, we give Hancock's two enlarged figures showing the dorsal and ventral views of *Terebratulina caput-serpentis* deprived of the shell, and exhibiting the respective positions of the various muscles above described.

While alluding to the digestive organs, Hancock states (*l. c.* p. 814) that "In *T. caput-serpentis* there appears to be only two hepatic ducts, and both in it and in *W.* [*Waldheimia*] *cranium* the intestine is very short, terminating in a blind sac before it reaches the ventral wall of the perivisceral chamber. It tapers gradually to a point which is rounded, and suspended in its place by the mesentery." The genitalia exhibit a somewhat different arrangement in *T. caput-serpentis* to what prevails in *W. flavescens*; "they are placed in large sinuses situated in the pallial lobes, one at each side. These sinuses are, however, nothing more than the enlarged trunks of the so-called pallial vessels or great pallial sinuses. They are four in number, two in each lobe, and the genital band, which is placed within them, forms a thick convolute layer, with small spaces between the folds."

"In *T. caput-serpentis* the heart is more decidedly pyriform than in the other species, and it is placed a little further back, the branchio-systemic vein passing for some distance down the stomach beyond the central gastro-parietal band . . . The setæ in *T. caput-serpentis* are placed rather far apart from each other, and issue from the mantle at the points corresponding to the marginal crenulations of the shell; these crenulations give to the pallial membrane a scalloped appearance. The setæ are rather robust; the marginal fold is deep, and the follicles are of considerable length and rather wide, with their bases surrounded with glandular matter, forming a roundish, red-coloured spot at the end of each seta" (*l. c.* p. 829). The blood-system of the brachial apparatus has likewise been well investigated by Hancock. He says:—"This is beautifully developed, and presents considerable variety in the character of the several plexuses of which it is composed. The walls of the great canal, the ridge supporting the cirri, the membranes that unite the upper and lower members of the loop, that which connects the spirals, and those which form the small canal or channel at the base of the cirri, as well as that forming the sheath of the apophysary support,—all have their system of lacunes which intercommunicate and compose the brachial system" (*l. c.* p. 831).

Dr. Van Bemmelen questions the lacunary system as described by Hancock. He concurs in the statement that in *T. caput-serpentis* and *T. septentrionalis* the sexes are separate*.

While treating of the perivisceral chamber, Hancock observes that in *T. caput-serpentis* the four trunks may be recognized. "Here the trunks are fused so as to form on each lobe two large, lateral, semilunar sinuses, in which the genitalia are placed. The external margins of these sinuses give off numerous, rather delicate branches, which dividing dichotomously run to the pallial margin; the branches next the middle line, which

* "On the Structure of the Shells of Brachiopods and Chitons," Ann. & Mag. Nat. Hist. 5th ser. vol. xi. p. 379.

correspond to the inner sinuses, pass off from their internal margins, and divide once or twice" (*l. c.* p. 840).

TEREBRATULINA CAPUT-SERPENTIS, var. *UNGUICULATA*. (Plate V. figs. 38-40.)

Terebratula unguicula, P. Carpenter, Proc. Zool. Soc. 1865, p. 201, figs. 1-4; Cooper, Geogr. Cat. California. Moll. p. 3, 1866; Dall, Amer. Journ. of Conch. vol. vi. p. 102, 1870, and Report on Brachiop. of Alaska, 1877; S. T. Whitcaves, On Some Marine Invertebrata from the West Coast of North America, Canadian Naturalist, n.s. vol. viii. p. 464, 1878.

Shell ovate, longer than wide; valves almost equally convex, rounded laterally and in front; lateral sides of the umbo strongly auricular, especially in young and middle-aged specimens. Ventral valve evenly convex, but sometimes very slightly depressed anteriorly; beak short, obliquely truncated by a rather large incomplete foramen, margined anteriorly by the umbo of the dorsal valve and by two small lateral deltidia. Surface of valves marked with numerous radiating delicate riblets, simple and stronger at their origin, but rapidly augmenting in number from bifurcation and by the interpolation of shorter riblets between the longer ones. Valves crossed with concentric raised striae. Loop short and simple, the crura disunited in the young, annelliform in the more advanced age. Colour light yellowish white. Length 13, breadth 10, depth 5 lines.

Hab. San Diego, Cal., to the Aleutian Islands, San Pedro, Cal. (Cooper); Neeah Bay (W. T. Swan); Victoria (V. W. Lord, J. Richardson); Port Etches; Shumargin Islands; Unalaska (Dall). Range: low water to 100 fathoms, those from deep water, the largest, adhering to shells and stones (Dall). At Race Island lighthouse, and Victoria Harbour, in 30 to 70 fathoms; end of Texada Island, in 40 to 70 fathoms, mud (Richardson).

Obs. Mr. Dall states, in the paper already referred to, that this species is readily separated from *Terebratulina caput-serpentis*, *T. japonica*, and other species, which strongly resemble it externally, by the broad loop, which is usually open, instead of being closed, as the genus requires. However, it finally becomes closed in fully adult specimens, which reach the size of *T. caput-serpentis*. I have also had the advantage of being able to examine a large number of young and adult examples of Carpenter's species, which were kindly lent to me by Mr. Whiteaves, as well as others which I purchased from Mr. G. B. Sowerby, who had obtained them direct from Vancouver Island. After having minutely compared their exterior and interior with a large number of European specimens of *T. caput-serpentis*, I arrived at the conclusion that *T. unguicula* is no more than a variety of Linné's species. Carpenter's specimens, as he himself admits, were all of small size, and consequently the crura of the loop were disunited. The specimens of *T. unguicula* I was able to open had not the broad loop described by Dall, but agreed well with the European specimens of *T. caput-serpentis* which I had previously opened and examined. In these specimens of *T. unguicula*, the front line was rounded, and not indented; but this is also the case with very many examples of *T. caput-serpentis*.

In a letter dated December 1884, Mr. Dall writes me that he would not wish to be quoted as believing *T. unguicula* distinct from *T. caput-serpentis**, and would be only

* [Dr. Davidson has given three excellent figures (see Pl. V. figs. 35, 36, & 37) of the well-marked variety of *Terebratulina caput-serpentis* from the Mediterranean Sea, to which Dr. Gwyn Jeffreys gave the varietal designation of *mediterranea*. He described it as longer and more slender than *T. caput-serpentis*, "more compressed or flatter than

willing to assert that the specimens he has seen seem distinguishable, and that as the distribution of the two forms is absolutely separate, their identity would seem to require proof.

12. *TEREBRATULINA CAILLETI*, Crosse. (Plate V. figs. 41, 42.)

Terebratulina Cailleti, Crosse, Journ. de Conch. vol. xiii. p. 27, pl. 1. figs. 1-3, 1865; Pourtales, Bull. Mus. Comp. Zoöl. (Harv.) vol. i. p. 109, 1867; Dall, Am. Journ. Conch. vi. p. 106, 1870, and Bull. Mus. Comp. Zoöl. (Harv.) vol. iii., 1871; Davidson, Report on the Brachiopoda, Voyage of H.M.S. Challenger, Zool. vol. i. p. 37, pl. 2. fig. 2, 1880; Dall, Bull. Mus. Comp. Zoöl. (Harv.) vol. ix. p. 103, 1881.

Shell ovate, longer than wide, broadest about the middle, rounded or slightly indented in front; dorsal valve evenly convex, no defined mesial fold, lateral sides of the umbo auricular; ventral valve moderately convex, with a strongly marked longitudinal depression or sinus, commencing at the beak and extending to the front; front raised into a rounded wave; beak short, and obliquely truncated by a moderate-sized incomplete foramen and two small disunited ætidial plates; surface of valves marked with about forty small radiating riblets, the larger number being shorter ones interpolated at various distances between the larger ones; surface marked also with fine concentric raised lines. Loop short, crura disunited in the young, annelliform in the adult. Colour of a slightly greyish white. Length 5, breadth 4, depth 2 lines; but Mr. Dall mentions having taken specimens 1 inch in length.

Hab. *T. Cailleti* was obtained by M. de Pourtales off Chorrera, Cuba, in 270 fathoms; near Cojima in 450 fms. Off Double-headed Shot Key in 471 fms., and near Tennessee Reef in 115 fms. Off West Florida 30 fms., St. Vincent 88 fms. (Pourtales and Dall). At Guadaloupe in 200 fms., by an Italian party who were searching for beds of coral. Mr. A. Agassiz sent me a specimen dredged by the Hunter Expedition off Barbados, taken at a depth of 100 fms., and I have also two young examples dredged by L. Barrett off Jamaica, and by Sigsbee off Havanna, in from 80 to 450 fms. I am likewise indebted to Mr. R. Rathbun for several young specimens obtained in 70 fms. in lat. 21° 48' S. and long. 40° 3' W. of Greenwich, which had been dredged by the Captain of the English Atlantic steamer 'Norseman.' A single example was obtained by the 'Challenger' Expedition, west of Pernambuco, in 350 fathoms.

Obs. This species has been minutely described by Mr. H. Crosse and by Mr. W. Dall. Crosse says that it is easily distinguishable from *T. caput-serpentis* by its stronger radiating riblets, these last being more separated from each other than in Linné's species. When quite young, and measuring 1 line in length, it has only from nine to ten simple ribs; at 2 lines already two or three short interpolated riblets have made their appearance close to the margin; at 3 lines in length the interpolated riblets become more numerous, and the whole surface is concentrically crossed by strongly marked, slightly projecting, equidistant, concentric ridges. The ridges become very much less marked as the shell approaches and attains the adult condition; but this feature is common to several species of the subgenus.

usual, and more or less cloven in front" ("Mollusca of the 'Lightning' and 'Porcupine' Expeditions," Proc. Zool. Soc. 1878, p. 401). From a MS. note attached to the types in the Davidson Collection, it is evident that Dr. Davidson considered this to be "a good variety of *T. caput-serpentis*," but identical with the shell named *Terebratula emarginata* by Risso in 1826, "of which the *Terebratula quadrata*, Risso, is probably a synonym." The variety *mediterranea* should therefore be henceforth known as *T. caput-serpentis*, var. *emarginata*, Risso.—A. C.]

Mr. Dall, in his paper in the Bull. Mus. Comp. Zoöl. already referred to, enters into many details to show that it is a distinct species. He has since then written me that, although *T. Caillieti* may be a southern form of *T. caput-serpentis*, he has seen probably nineteen twentieths of all the existing specimens, and that it seems to him to be as well distinguishable as any of the fossil forms, and that he considers it at any rate fully entitled to a varietal name, as it is well distinguished by its granulated ribs, and varies much in form.

Dall says (*l. c.* p. 10) "That the smallest specimens of this species which I was able to find among those sent by M. de Pourtales were nearly .1 inch in length. That the characteristic sculpture was developed upon them to the very apex of the shell. . . . The various muscles were already well developed. The mouth was as described in the young of the *T. cubensis*. The intestine was short, cylindrical, and straight. The lower portion was embraced by a few hepatic digitations. These lobes were very dark brown, the muscles of a deep reddish brown, and the brachia of a flesh-colour. The latter were in the shape of a horseshoe, with no trace of a median lobe. They were close set and marked with transverse lines as in *T. cubensis*. The membrane which covered the viscera was covered internally with irregular hyaline spots with well-marked boundaries, which no doubt are the limits of the lacunar channels of circulation. The mantle was quite transparent, with a brownish edge, and in each of the internal channels, corresponding to the ribs of the outside of the shell, was a single bristle, composed of longitudinal fibres of chitine, without any of the transverse markings which are seen in the setæ of the adult. The extreme tip of the bristle alone protruded from the mantle, and its inner extremity was slightly bulbous. It was of a glistening yellow color throughout. In those adults which I examined there were only five or six of these setæ in each mantle lobe. These specimens were obtained off Havana, in two hundred and seventy fathoms water. The very extraordinary manner in which all the soft parts were crowded and crammed with masses of calcareous spiculæ defied my best efforts to obtain any very satisfactory results from the two or three alcoholic specimens at my command. A flocculent mass of white matter resisted the action of acid, and filled all the interstices of the membranes so as to render them quite opaque. . . . The intestine was cylindrical, and ended much as it does in *T. caput-serpentis*. The mouth was surrounded by a dark-brown line. There were no structures above and behind the mouth, such as are described as existing in *T. cubensis*. The attached extremities of the muscles were of a very bright red-brown. . . . The peduncle is white, slender, and exceedingly long, the exposed portion sometimes equalling in length one third of the shell. A brownish tinge pervaded all the tissues of the adult. Transverse markings were noticed in the brachia, as described in other species by Hancock. One specimen, growing on a rock which had become covered with sponge, afforded an interesting observation. The peduncle was exceedingly long, and on cleaning off the sponge it was seen that the creature, on the growth of the sponge towards it, had apparently lengthened its peduncle to get out of the way; and while the original attachment still remained, . . . somewhat further on, nearer the shell, a second attachment of the peduncle had taken place by the outgrowth, from the underside, of a bunch of cylindrical rootlets, exactly resembling the attachment of an ivy to a stone."

13. *TEREBRATULINA SEPTENTRIONALIS*, Couthouy. (Plate V. figs. 1-31 & 43-52.)

Terebratulina septentrionalis, Couthouy, Boston Journ. Nat. Hist. vol. ii. p. 65, pl. iii. fig. 18, 1838.

Terebratula septentrionalis, G. B. Sowerby, Thes. Conch. p. 342, pl. xviii. figs. 5 & 6, 1846; Stimpson, Test. Moll. New England, p. 75, 1851.

Terebratula caput-serpentis, L. Reeve, Monogr. of Terebratula, Conch. Icon. 1861.

Terebratulina septentrionalis, E. Morse (On the early Stages of), Mem. Boston Soc. Nat. Hist. vol. ii. (read Oct. 1869); Gould, Invert. Mass., Binney's 2nd ed. p. 208, fig. 500, 1870.

Terebratulina septentrionalis, Dall, Cat. of Recent Brach., Proc. Acad. Nat. Sci. Philadelphia, 1873, p. 180; Sars, Fauna Moll. Regionis Arcticæ Norvegiæ, p. 10, tab. i. fig. 4, 1878.

Terebratulina caput-serpentis, var. *septentrionalis*, Davidson, Report on the Brachiopoda, Voyage of H.M.S. 'Challenger,' Zoology, vol. i. p. 33, pl. i. figs. 3-9, 1880.

Terebratulina caput-serpentis, W. F. Ganong, The Invertebrate Zool. of Passamaquoddy Bay, Nat. Hist. Soc. of New Brunswick, Bull. No. 4, p. 91, 1884.

Shell rather thin, longitudinally broadly ovate, pear-shaped or somewhat subpentagonal, narrow and tapering posteriorly, abruptly widening below the beak, broadest anteriorly; front line nearly straight, slightly indented or rounded; colour yellowish white, nearly diaphanous; dorsal valve gently convex, especially towards the middle, slightly depressed towards the front, eared at the umbo; ventral valve feebly convex, or slightly deeper than the dorsal one, somewhat depressed towards the front; beak short, attenuated, slightly incurved and truncated by a moderately large and incomplete semielliptical foramen, completed below by the umbo of the ventral valve, and laterally margined by small deltidial plates; surface of both valves covered with a variable number of fine rounded radiating raised striæ or riblets (240 in some specimens when counted at the margin), these increase in number at variable distances from the beaks by the interpolation of shorter riblets; shell-structure perforated by numerous small canals. In the interior of the dorsal valve the loop is short and simple, rendered annular, with age, by the union of the oral processes. The brachial appendages are united to each other by a membrane, cirrated and developed from each side of the mouth, divided into three lobes, the two lateral ones extending to a little more than two thirds of the length of the valve, the central one not exceeding half the length of the valve, and spiral at its extremities. Proportions variable. Length 15, breadth 11, depth 6 lines.

Hab. The geographical range of *T. septentrionalis* seems to be very extended. The 'Challenger' Expedition dredged it abundantly off the New-York coast at depths of 51 fathoms, also, at a depth of 83 fathoms, off Halifax. Again, according to the officers of the expedition, off the Cape of Good Hope (but I should like to have this last statement confirmed), at latitude 46° 40' S., long. 37° 50' E., at a depth of 150 fathoms. Mr. Couthouy, in his description of *T. septentrionalis*, says that it has been found at Lubec Bay by Dr. C. T. Jackson during his geological survey of the State of Maine, and that it is probably an inhabitant of deep water on the whole New-England coast. Dr. Gould mentions having found it in considerable numbers in the stomach of fishes, and occasionally on the sea-beach, and that its usual habitat is in the Laminarian or deep Coral zones of northern seas. At Eastport, at low water, it is common off the

Isle of Shoals, 20 fathoms; Grand Manan, common (Packard, Stimpson); Halifax harbour (Willis); Trias Cove, Passamaquoddy Bay, in 12 fathoms, where they occur clinging together in bunches, large and small together, and seem to prefer the clear fresh water on pebbly bottoms (W. F. Ganong).

Obs. The above description is the one I published in my 'Challenger' Report. Some difference of opinion has prevailed with respect to the specific claims of the species under description. Certain malacologists consider it a distinct species, others as only a simple variety of *T. caput-serpentis*. Gould, in 1838, believed it to be distinct; but in the 'Report on the Invertebrates of Massachusetts' (2nd ed. by W. H. Binney, 1870) we find stated at p. 208:—"An examination of the descriptions of *T. caput-serpentis* given by Linnæus, Müller, and Chemnitz, and a comparison of them with our shell, has well satisfied me of their correspondence. The downy epidermis is a character too rare and singular to be overlooked. This, however, is rubbed off very easily. The shell is much thinner, in general more elongated, and the striæ nearly twice as numerous, being about thirty to forty in the European, and fifty to sixty in the American specimens. No account of the internal bony processes is given in any description except that of Mr. Couthouy. These would afford the best possible specific character, were it not that they are usually more or less broken. But I have been relieved from all further speculation by the receipt of specimens from Dr. Lovén, which settle the identity of our species with the European *caput-serpentis*." Mr. W. H. Binney seems to be of a different opinion, for he adds, after Gould's observations just recorded:—"I have retained the above remarks from the former edition, because our shell is so generally still regarded as identical with the European species. But further examination of numerous specimens has led me to coincide with Dr. Stimpson, who has dredged extensively, both in the British and American seas, in his opinion that the species differs from the European *caput-serpentis*, sufficiently in both shell and animal."

G. B. Sowerby, on p. 344 of his 'Thesaurus Conchyliorum' (1846), observes that *T. septentrionalis* is distinguished from *T. caput-serpentis* by its much finer radiating striæ, its larger and less oblique foramen, and by its rather more extended and somewhat differently formed internal appendage.

Since publishing my description of this shell in the reports of the 'Challenger' Expedition, I feel more disposed to leave *T. septentrionalis* and *T. caput-serpentis* as separate species, the first being evidently more regularly oval and rounded in front than in Linné's species; and even in the young stage the riblets are more numerous than in *T. caput-serpentis*. I have been able to examine specimens from less than 1 line in length up to that of 1 inch and 3 lines. Up to about 3 lines, and sometimes even more, the ribs are very few in number, prominent and radiate from the extremity of the beak to the margin, and are crossed by strongly indented concentric lines, which give to the riblets the so-termed tuberculated appearance that has been so often described in the species of this subgenus. As the shell grows the ribs become more delicate and more numerous from repeated interpolations of shorter ribs, and the concentric ones become more and more faintly marked.

It is, however, to Prof. E. Morse that science is most indebted for the knowledge we

at present possess of the embryology, or early stages, of *Terebratulina septentrionalis**. This distinguished zoologist having dredged in 15 fathoms of water in the harbour of Eastport, Maine, in 1869, individuals of all ages, has minutely and elaborately described all the stages of its development, which he illustrates in the most complete and admirable manner. The details and illustrations he communicates are so numerous that we feel ourselves obliged to refer the reader to the author's memoir, and to limit ourselves to the reproduction of the following selected observations and illustrations; he says:—
 “The species occur in great numbers, at various depths, and have also been collected at low-tide mark, by Dr. Stimpson and Prof. Verrill. The specimens were found attached to stones brought up in the dredge, and also adhering to the lower valve of adult individuals, generally near the peduncle†. . . . The eggs were generally kidney-shaped, though very irregular as to form and size [see Plate V. figs. 1–4]; they were spermaceti-white in color, and opaque, though having a central area, translucent, and apparently depressed. . . . This form recalled the general proportions of *Argiope* and *Megerlia*, in being transversely oval, in having the hinge margin wide and straight, and in the presence of a proportionately wide foramen. This stage was exceedingly minute, . . . the shell showed nothing of the scale-like structure so characteristic in later stages. Between this stage and the next, the shell rapidly elongates, while the hinge margin remains nearly the same in width; this is also shown in the concentric lines of growth seen faintly on the surface, indicating a rapid increase in the length of the shell, while no corresponding increase takes place in the widening of the hinge margin. The peduncle is longer than the shell, having distinct walls apparently enclosing a clear interspace, the end slightly dilating and forming a pear-shaped adhering disk. The structure of the shell . . . showed clearly the scale-like structure, with the cæcal tubules of the pallial lobes perforating it. The anterior margin of the pallial lobes gave rise to seven setæ of variable lengths, all of them projecting forward [Plate V. fig. 21]. . . . The future position of the calcareous loop was indicated by a strongly arched process midway the length of the shell, from which sprang six short and stout cirri, all of them curving towards the mouth, which occupied the centre of the base from which the cirri sprang. The digestive sac hung from the mouth, and was twice as long as broad, having a strong constriction in the centre, forming two chambers, the lowermost one being globular in shape, and having its walls colored a light reddish-brown, this colored portion evidently indicating hepatic cells. The cavity next the mouth indicates the stomach, whilst the lowermost cavity indicates the future intestine or cul-de-sac. The cirri moved frequently, and in various directions, though generally performing a grasping motion, as if securing some bit of food. . . . In this stage, and several succeeding stages, the outline of the shell is remarkably like that of *Lingula*, and this resemblance is more striking from the proportionally long peduncle. [Plate V. fig. 31.]

“In another stage, the numerous irregular shaped calcareous spiculæ lined the outer

* “On the Early Stages of *Terebratulina septentrionalis*,” *Memoirs of the Boston Society of Natural History*, vol. ii. p. 29, 1871; and “Embryology of *Terebratulina*,” *ibid.* p. 249.

† On an adult specimen of *T. septentrionalis*, dredged by the ‘Challenger’ expedition, I counted no less than twenty-eight young shells of the species attached to the surface of both valves, and more especially to its peduncle.—T. D.

margin of the cirri, while the future position of the calcareous loop, or crura, was indicated by a row of irregular-shaped spiculæ. . . . In these slightly advanced stages, the peduncle becomes much shorter in proportion to the length of the shell. . . . In all these stages the peduncle has very slight adhesion to the rock or whatever substance it may be attached to, in this respect differing greatly from the adult, which often requires at this age great force to detach it. . . . I was fortunate in observing an individual [slightly larger] in motion. The animal whirled quickly on its peduncle; when at rest the shells were always closed, and rested on the rock; from this position it turned slowly more than half way round, raising the body at the same time almost erect; this movement being completed, the valves would very slowly open, and the cirri expand as if to perform a grasping motion; in no case, however, were they projected beyond the margin of the valves. The cilia lining the cirri produced gentle currents in the water. In this position, with the valves widely open and the cirri expanded, the animal would remain motionless for twenty or thirty seconds, and then, with an abrupt closing of the valves, suddenly assume its first position. . . . In watching these motions for a long time, one could not help being impressed with the fact that great caution was evidently indicated in the slow and careful movements made in elevating and opening the shell, while the prompt closing of the valves, and the alert manner in which the animal regained its first position, seemed to show that food had been secured, and further caution was unnecessary."

Prof. Morse then goes on to describe how in successive stages a fold on each side of the stomach was first noticed, these folds being the first appearance of the liver, then the upper portion of the digestive sac and that portion which answers to the œsophagus and stomach in the contracted state, while the lower portion is widely expanded. In a still more advanced stage, he says:—"The shell is now becoming proportionally broader and the cirri increase in number, though still forming a simple circle around the mouth. The crura have also begun to form. At a considerably more advanced stage "The cirri are more numerous, numbering thirty-one: two of them are seen encroaching upon the circular lophophore, and at this stage the lophophore has begun to assume its hippo-crepian character.

"The crura are plainly seen supporting the crown of cirri, and the liver already shows the first indications of its differentiations under the peculiar cæcal ramifications which become so numerous in the adult. . . . The divaricators are completely formed, and between these two muscles is seen the lengthened intestine, the blind extremity of which is held firmly to the shell by a membrane, called by Hancock the ventral mesentery." At this stage "Radiating ribs, to the number of fifteen, ornament the shell."

Successive stages are then described. At p. 35 of his admirable memoir, Prof. Morse enters into long details in connexion with the structure of the shell; he says:—"In the youngest stages of the shell the scale-like structure may be studied to the best advantage. The scales are few in number, but slightly overlapping, and form a layer quite distinct from the outer layer, which appears to be homogeneous, save the concentric lines of growth appearing like rows of oblong and flattened nodules. The terminal portions of the pallial cæca within or upon this outer layer, are brown in colour and distinctly

granulated. . . . the tubules radiate from the largest diameter of the cæca, and not from the periphery of the granulated disk, and this is in accordance with the admirable observations made by King*. The scales do not appear to encroach upon the walls of the cæca."

Then follow many very important details on the growth of the shell, as well as the gradual development of the crura and loop. The crura of the calcareous loop, although not connected by a calcareous shelly layer in the young animal at all stages, have portions nevertheless connected by a membrane charged with spicula. The peduncular opening also, with age, becomes more circular, while the cardinal process, which does not appear in the earlier stages, is latterly present.

In Prof. Morse's memoir in question a vast number of important details are given, to which we have only briefly alluded.

I have myself examined specimens at all stages of growth, brought home by the 'Challenger' Expedition, and I am able to confirm Prof. Morse's accurate observations. When half a line in length the dorsal valve is elongate semicircular, the hinge-line straight, and nearly as long as the breadth of the shell, only eight or nine rounded ribs ornamenting the surface of each of its valves; by degrees the hinge-line becomes shorter and obtusely angular, and the ribs that ornament the surface of the valves become more numerous.

14. TEREBRATULINA WYVILLII, Davidson. (Plate III. figs. 1-3.)

Terebratulina Wyvillii, Davidson, Proc. Roy. Soc. vol. xxvii. p. 436, 1878, and (*T. wyvillii*) Report on the Brachiopoda, Voyage of H.M.S. 'Challenger,' Zoology, vol. i. p. 32, pl. i. figs. 1-2, 1880.

Shell large, trigonal, longer than wide, broadest anteriorly, light yellowish; dorsal valve triangular, anterior angles rounded; hinge-line obtusely angular, moderately convex, somewhat flattened along the middle, and abruptly bent inwards close to the margin; lateral sides of the umbo auricular; ventral valve convex, deeper than the opposite one, flattened anteriorly, abruptly bent inwards close to the margin; beak incurved, truncated by a large oval-shaped foramen, separated from the hinge-line by a triangular concave depression sharply defined laterally; surface of shell nearly smooth to the naked eye, but marked by very fine radiating raised lines; perforations rather large. In the interior of the dorsal valve the loop is short and simple, rendered annular by the union of the oral processes. Length $2\frac{1}{2}$ inches, breadth 2 inches, depth 1 inch 4 lines.

Hab. A single specimen of this fine species was dredged on the 25th of March, 1873, by the 'Challenger' Expedition off Culebra Island, to the north-west of St. Thomas, in the West Indies, depth 390 fathoms. It was also obtained in some abundance by the French expeditions of the 'Travailleur' and 'Talisman' in 1882 and 1883, near the western coast of Africa. It is alluded to by Prof. A. Milne-Edwards in the 'Comptes Rendus de l'Académie des Sciences,' tom. xciii. Nov. 1881, and has also, according to Dr. P. Fischer, been dredged off the coast of Spain.

* "On the Histology of the Test of the Class Palliobranchiata," Trans. Royal Irish Academy, vol. xxiv.

Obs. This is the largest species of *Terebratulina* with which I am acquainted, either in the recent or fossil condition.

15. *TEEBRATULINA CROSSII*, Davidson. (Plate III. figs. 4-6.)

Terebratulina Crossei, Davidson, Journal de Conchyliologie, vol. xxx. p. 106, pl. vii. fig. 1, 1882.

Shell large, elongated oval, longer than wide; valves almost equally convex, mesial fold and sinus scarcely indicated; dorsal valve feebly auriculated; beak of ventral valve slightly incurved, obliquely truncated by a rather large foramen, margined in part by the umbo and by two small lateral deltidial plates; surface of valves covered with numerous fine radiating striæ, at times almost obsolete, and crossed by concentric lines of growth. Colour light yellowish or light salmon. Loop short, not quite a third or fourth of the length of the valve. Length 2 inches 1 line, width 1 inch 8 lines, depth 1 inch.

Hab. This magnificent species occurs in Sagami Bay, Japan, where it was dredged by Dr. L. Döderlein. The locality is situated in the neighbourhood of station No. 232 of the 'Challenger' Expedition, only a little nearer the coast, and in about 100 to 250 fathoms. Dr. Döderlein, who kindly lent me the fine series of Brachiopoda procured by himself in that locality, tells me that the sea-bottom is covered with mud, stones, and fragments of volcanic origin, with abundant animal life of astonishing variety, and that there is no better locality for dredging-purposes in Japanese waters.

Obs. I became first acquainted with this fine species in 1882, M. H. Crosse having kindly forwarded a specimen he had received from Japan, which I subsequently described and figured in his valuable 'Journal de Conchyliologie.' Soon after I was able to purchase from Mr. G. B. Sowerby another good specimen of the same shell, and subsequently two other fine examples were lent to me by Dr. Döderlein. I have given figures of three of the specimens. It does not appear to be a very rare species in the locality where it is found.

I may also mention that no single locality has hitherto afforded a larger or more varied assemblage of specific forms of Brachiopoda than the waters that surround the Japanese islands, Corea, and the neighbouring coast of China. I have been able from those seas to examine and describe the following species:—

1. <i>Terebratula</i> or <i>Terebratulina</i> <i>Dalli</i> , Davidson.	14. <i>Terebratella frontalis</i> , Middendorff.
2. — <i>Davidsoni</i> , Adams.	15. <i>Magasella Adamsi</i> , Davidson.
3. <i>Terebratulina Crossii</i> , Davidson.	16. — <i>Gouldi</i> , Dall.
4. — <i>caput-serpentis</i> , Linné.	17. <i>Megerlia</i> or <i>Ismenia sanguinea</i> , Chemnitz.
5. — <i>japonica</i> , G. B. Sowerby.	18. — <i>Reevei</i> , Davidson.
6. — <i>Cumingi</i> , Davidson.	19. — <i>pulchella</i> , G. B. Sowerby.
7. — <i>radiata</i> , Reeve.	20. <i>Laqueus picta</i> , Chemnitz.
8. <i>Waldheimia Raphaelis</i> , Dall.	21. — <i>rubella</i> , G. B. Sowerby.
9. — <i>Grayi</i> , Davidson.	22. <i>Platydia anomioides</i> , Scacchi.
10. <i>Terebratella spitzbergensis</i> , Davidson.	23. <i>Rhynchonella psittacea</i> , var. <i>Woodwardi</i> , Adams.
11. — <i>Blanfordi</i> , Dunker.	24. — <i>lucida</i> , Gould.
12. — <i>coreanica</i> , Reeve.	25. — <i>Döderleini</i> , Davidson (a species covered
13. — <i>Mariæ</i> , Adams.	with spines).

26. *Crania japonica*, Adams.
 27. *Discinisca stella*, Gould.
 28. *Lingula anatina*, Lamarck.
 29. — *Adamsi*, Dall.

30. *Lingula jaspidea*, Adams.
 31. — *smaragdina*, Adams.
 32. — *lepidula*, Adams.

16. *TEREBRATULINA JAPONICA*, Sowerby, sp. (Plate III. figs. 7–11.)

Terebratula japonica, Sowerby, Proc. Zool. Soc. 1846, p. 91; Thes. Conch. vol. i. p. 344, pl. lxxviii. figs. 7–8, 1846; Adams & Reeve, Voyage of the 'Samarang,' Mollusca, p. 71, pl. xxi. fig. 1, 1848.

Terebratula angusta, Adams & Reeve, Voyage of the 'Samarang,' Mollusca, p. 71, pl. xxi. fig. 2, 1848.

Terebratulina japonica, L. Reeve, Conch. Icon., Monogr. of *Terebratula*, pl. iv. fig. 16, 1860; A. Adams, Ann. & Mag. Nat. Hist. 3rd ser. vol. xi. p. 98, 1863.

Terebratulina caput-serpentis, var. *japonica*, Davidson, Proc. Zool. Soc. 1871, p. 303, pl. xxx. fig. 8.

Terebratulina japonica, Dall, Cat. of the Recent Species of the Class Brach., Proc. Acad. Nat. Sci. Philadelphia, 1873, p. 180.

Terebratulina caput-serpentis, var. *japonica*, G. Dunker, Index Moll. maris Japonici, p. 251, 1882.

Shell oblong, ovate, almond-shaped, widest about the middle, rounded in front, valves nearly equally and uniformly convex, no defined fold or sinus; beak slightly incurved and obliquely truncated by an incomplete circular foramen; deltidial plates rudimentary. Auricles in dorsal valve very small, marginal line nearly straight, inclining to the sides near the hinge. Surface of valves longitudinally striated, striæ numerous and radiating, augmenting in number at intervals through the bifurcation of some of the ribs and by the interpolation of shorter ones of variable thickness. Valves crossed by concentric lines of growth. Colour light salmon or nearly white. In the interior of the dorsal valve the loop is small and ring-shaped. Length 1 inch 4 lines, breadth 1 inch, depth 7 lines.

Hab. Gotto, 48 fathoms; Tsusalei, 55 fathoms (Adams); Sagami Bay, Japan (Döderlein).

Obs. This species is remarkable on account of its spindle-like shape; it is nearest related to *T. Crossii*, and easily distinguished from the other recent species of the genus. Lovell Reeve says, in his Monograph of *Terebratula*, that "*T. japonica* is very closely allied to *T. caput-serpentis* and is without doubt its representative in the Corcan and Japanese waters." I must, however, differ from Mr. Reeve in this matter, although in 1871 I expressed a similar opinion. Dr. Döderlein, it is true, dredged in Sagami Bay a specimen which I could not distinguish from Linné's *T. caput-serpentis*, but it differs in many respects from *T. japonica*, which occurs in the same locality. I am, however, quite prepared to concur with Mr. Reeve's statement that *T. angusta*, Adams & Reeve, is only a smaller form of *T. japonica*. Dall places *T. abyssicola*, Reeve, *T. angusta*, Reeve, and *T. Cumingi*, Davidson, among his synonyms of *T. japonica*. Reeve's figures of *T. abyssicola* would hardly lead me to consider it a synonym of the above-named species, and a similar doubt may be entertained with respect to *T. Cumingi*.

17. *TEREBRATULINA RADIATA*, Reeve. (Plate VI. figs. 9–14.)

Terebratula (Terebratulina) radiata, L. Reeve, Conch. Icon. pl. iii. figs. 7 a–b, 1860, and Ann. & Mag. Nat. Hist. 3rd ser. vol. vii. p. 173, 1861.

Terebratulina radiata, Dall, Proc. Acad. Nat. Sci. Philadelphia, p. 180, 1873.

Shell subpentagonal or ovate, longer than wide, indented in front, flexuously waved towards the margin. Dorsal valve convex, anterior half biplicated, a median concave depression commencing at about half the length of the valve, separating the two rounded ribs. Ventral valve convex, somewhat flattened along the middle and divided on the anterior half by a median rounded rib, with concave groove on either side. Beak short, obliquely truncated by a rather large foramen, completed by a part of the umbo of the opposite valve and by two small lateral deltidial plates. Surface of valves densely covered with very fine radiating striæ of unequal width, and increasing in number through bifurcation and the interpolation of shorter riblets. Colour dirty white, sometimes rayed with broad pencils of black. Loop short and simple, incomplete when young, but with age becoming annular through the union of the crural processes. Length 11 lines, by 9 in breadth and 6 in depth.

Hab. L. Reeve says "Mr. Cuming possesses two or three specimens, all exactly alike, procured, he fancies, from the dredgings of Sir E. Belcher in the Strait of Corca." I have seen and possess a number of specimens of this shell, which Mr. Sowerby assures me were obtained near the Cape of Good Hope, its probable habitat.

Obs. Mr. Lovell Reeve considers this to be a good and well-marked species, and in this statement I feel disposed to concur. I have seen a great many specimens of the shell, some as wide as long, others longer than wide, and all presenting a more or less well-marked biplication. In some exceptional examples there existed also an additional rib on the lateral portions of the dorsal valve. *T. radiata* is remarkable, as observed by Mr. Reeve, on account of being very often sparingly rayed with black. It is a smaller shell than *T. cancellata*, its nearest ally. In some specimens the foramen becomes complete by the union of the deltidial plates, but this appears to be the exception and not the rule.

18. TEREBRATULINA CANCELLATA, Koch, sp. (Plate VI. figs. 1-8.)

Terebratula cancellata, Koch, in Küster, Conch.-Cab. vol. vii. p. 35, pl. 2b. figs. 11-13, 1843.

Terebratula, species quædam ignota, Menke, Moll. Nov. Holland. specim. p. 35, no. 192.

Terebratula cancellata, Sowerby, Thes. Conchyl. p. 358, pl. lxxi. figs. 93-95, 1846.

Terebratula (Terebratulina) cancellata, Reeve, Conch. Icon. pl. iv. fig. 13, 1860.

Terebratulina cancellata, Dall, Proc. Acad. Nat. Sci. Philadelphia, p. 179, 1873; Davidson, Report on the Brachiopoda, Voyage of H.M.S. 'Challenger,' Zool. vol. i. p. 37, pl. i. figs. 11-16, 1880.

Shell rather large, elongated oval, livid brown or light yellowish with darker concentric bands, widest about the middle, nearly straight in front. Dorsal valve very convex, flattened longitudinally along the middle, from which the lateral portions slope away at a strong angle. Ventral valve convex, rather less deep than the opposite one and sometimes flattened towards the front; beak incurved, tapering, moderately produced and truncated by a rather large foramen, which is very slightly separated from the hinge-line by a very narrow deltidium, the beak often slightly overlying the umbo of the dorsal valve. Surface of both valves ornamented with a great number of delicate raised striæ or riblets, increasing in number at various distances from the beaks by the interpolations of shorter riblets. The surface is also crossed at intervals by fine concentric lines of

growth. In the interior of the dorsal valve the loop is simple and short, and rendered annular in the adult by the union of the crural processes. Shell-structure perforated by canals. Brachial appendages united by a membrane and divided into three lobes, the central shorter one spirally coiled. Length 1 inch 7 lines, breadth 1 inch, depth 10 lines.

Hab. Küster informs us that his types were derived from Western Australia. His figures are not good, and he colours them dark brown; some specimens, it is true, are of livid brown, but those brought back by the 'Challenger' Expedition were of a yellowish colour. It was dredged abundantly by the 'Challenger' Expedition in 1874 off Moncœur Island, Bass's Strait, in a depth of 35 to 40 fathoms, sea-bottom mud; many of the specimens being attached by their peduncle to *Pecten*, *Cardium*, *Arca*, spines of *Cidaris*, stones, &c. A dwarfed form was also dredged for the first time by Mr. John Brazier of Sydney, at Old Man's Hat Point, Inner North Head, off Port Jackson, N.S.W., near the rocks, in 7 fathoms, bottom of dead broken shells with coarse sand and stones; also at Sow and Pigs Reef, eastern side, Port Jackson, near the rocks, in 3 fathoms bottom. One example was found adhering to a valve of *Trigonia Lamarckii*, Gray, and is the only specimen Mr. Brazier ever found on *Trigonia*.

Obs. When young *T. cancellata* is often longitudinally spindle-shaped and oval, but with age shows more of a straight front. Some of the 'Challenger' specimens have also much exceeded in size those figured by Küster, G. B. Sowerby, and L. Reeve. There exists likewise in some exceptional specimens a median depression on the anterior portion of the dorsal valve, commencing at about the middle of the shell and extending to the front.

19. TEREBRATULINA (AGULHASIA) DAVIDSONI, King. (Plate VII. figs. 1-5.)

Agulhasia Davidsoni, King, Annals & Mag. of Nat. Hist. 4th ser. vol. vii. p. 111, pl. xi. figs. 1-7, 1871.

Terebratulina Davidsoni, Dall, Proc. Acad. Nat. Sci. Philadelphia, 1873, p. 180.

Shell very small, elongated triangular, much longer than wide. Dorsal valve about as wide as long, broadest and slightly rounded anteriorly, more tapering posteriorly, uniformly and moderately convex without fold, lateral sides of umbo gently auriculated. Ventral valve pyramidal, rather deeper or more convex than the dorsal one and with a slight mesial depression near the front, beak solid, nearly one third of the length of the valve, tapering to a small incurved point, area in form of an acute isosceles triangle; deltidium long and narrow, closed except at the cardinal termination by the internal plate, foramen incomplete and notch-shaped, lying close to the umbo of the opposite valve. Surface of valves radiately striated with about twenty small ribs, of which about half the number are shorter ones interpolated between the longer ribs. Loop in dorsal valve short and simple, furnished with long crural spurs. Colour white. Length 3 lines, by 2 in breadth and 1 in depth.

Hab. Agulhas bank, South Africa, and obtained from a depth of 45 to 60 fathoms.

Obs. In 1870 several specimens of this elegant lyre-shaped shell were placed in my hands by Mr. G. B. Sowerby, and having sent them to my distinguished friend Prof. W. King, he kindly offered to publish a description of the shell under the name of *Agulhasia Davidsoni*. Although possessing characters bearing relation to *Terebratulina*, Prof. King

arrived at the opinion that it could be generically distinguished. The chief differences are to be found in the shape and character of the beak. "The long narrow form of the deltidium," writes Prof. King, "causes the area to appear as if divided longitudinally by a linear groove; which part is at once striking and unique. The foramen, which appears like a notch in the centre of the cardinal edge of the large valve, is made entire by the juxtaposition of the cardinal edge of the small valve. The area is well defined laterally by each of the sutures which separate it from the inflexed sides of the beak. The loop agrees very closely in form and relative size with that of *Terebratula vitrea*; perhaps its crural spurs are more produced. The tubuli appear to run in rows, and to be most numerous in the ribs, from which they branch off laterally towards the furrows." Mr. Dall believes the specimens described by Prof. King to be the young of some species of *Terebratulina*, but the specimens I have in my possession would lead me to consider them the adult condition of a very remarkable small species.

Uncertain Species.

20. *TEREBRATULINA CUMINGI*, Davidson. (Plate V. fig. 53.)

Terebratulina Cumingii, Davidson, Proc. Zool. Soc. 1852, p. 79, pl. xiv. figs. 17-19; Reeve, Conch. Icon., Monogr. of Terebratula, pl. iv. fig. 12, 1860.

Shell small, squarely ovate, somewhat pentagonal, nearly as wide as long; valves almost equally convex; beak small, obliquely truncated by a large incomplete foramen, deltidial plates disunited, a small portion of the aperture completed by the umbo; auricular expansions on either side of the umbo, very small; valves rather gibbously convex. Surface ornamented by a great number of minute radiating striæ, augmenting rapidly by the interpolation of smaller riblets at variable distances from the beaks; front margin in ventral valve forming a convex curve, and slightly depressed near the front. Loop small, annelliform. Colour opaque white. Length $3\frac{1}{2}$, width 3, depth 2 lines.

Hab. Chinese Seas.

Obs. Mr. L. Reeve says that this is "An extremely characteristic well-defined species. The radiating ridges do not all diverge together from the beak; a new ridge commences between each former ridge as the space enlarges to receive it." *T. Cumingi* seems to be distinguished from other species of the subgenus by its size and relative width and length, being much more convex and globular. In 1871 (Proc. Zool. Soc.) I thought that *T. Cumingi* might perhaps be a variety of *T. caput-serpentis*, but am now not of that opinion, for having seen a number of specimens of the little Chinese shell, I found them all to be possessed of the same shape and character.

In his "Catalogue of the Recent Species of the Class Brachiopoda" (Proc. Acad. Nat. Sci. Philadelphia, 1873), Mr. Dall places *T. Cumingi* among his synonyms of *T. japonica*, but I cannot agree with him in this particular.

21. *TEREBRATULINA ABYSSICOLA*, Adams & Reeve, sp. (Plate V. fig. 54.)

Terebratula abyssicola, Adams & Reeve, Moll. Voyage of the 'Samarang,' p. 72, pl. xxi. fig. 5, 1850; L. Reeve, Conch. Icon. pl. iv. fig. 14, 1860.

"Shell triangularly pear-shaped, orange flesh-colour, radiately obsoletely ridged, ridges distant, with numerous bifurcated striæ coming between them, beak obtusely produced, foramen rather large, incomplete, deltidium obsolete; loop short" (Reeve). Length 5, breadth $4\frac{1}{2}$, depth $2\frac{1}{2}$ lines.

Hab. Cape of Good Hope (dredged at a depth of 120 fathoms by Sir E. Belcher).

Obs. I have seen three examples only of this species (?), one of which was attached to a specimen of *Kraussina rubra*. In the type figured by Reeve there is evidence of a longitudinal depression along the middle of the dorsal valve, and in an authenticated example in my possession the riblets seem somewhat fasciculated.

Dall, at page 180 of his "Catalogue of the Recent Species of the Class Brachiopoda" (Proc. Acad. Nat. Sci. Philadelphia, 1873), places *T. abyssicola* among his synonyms of *T. japonica*; but I fear I cannot concur with my friend in this identification. The Cape-of-Good-Hope form is evidently more nearly related to *T. caput-serpentis* than to *T. japonica*. More specimens will have to be examined before a positive conclusion can be arrived at with respect to its specific claims.

22. TEREBRATULINA ? INCERTA, Davidson. (Plate VI. figs. 23-25.)

Megerlia (?) *incerta*, Davidson, Proc. Royal Soc. vol. xxvii. p. 438, 1878; and Report on the Brachiopoda, Voyage of H.M.S. 'Challenger,' Zoology, vol. i. p. 49, 1880.

Shell small, nearly circular, somewhat broader than long, widest towards the middle. Hinge-line long, nearly straight or very obtusely angular, rather exceeding two thirds of the breadth of the shell, with obtuse cardinal angles. Dorsal valve somewhat semicircular, very slightly convex, most so at the umbo, lateral sides of the umbo auricular, no fold or sinus. Ventral valve a little deeper or more convex than the opposite one, slightly longitudinally depressed along the middle; beak short and truncated by an incomplete circular foramen margined by the umbo of the dorsal valve and by two lateral deltidial plates; beak-margins sharply defined, leaving between them and the hinge-line a narrow area. Surface of both valves marked by numerous fine rounded radiating riblets with concave interspaces, some of the ribs bifurcating near the front, other shorter ones intervening between the longer ones at variable distances from the beaks. Valves crossed by numerous irregular concentric lines of growth. Shell perforated by minute canals. Colour yellowish white. In the interior of the dorsal valve the cardinal process is long and narrow; the loop short and simple. Length 5, breadth $5\frac{1}{2}$, depth $2\frac{1}{4}$ lines.

Hab. One young example attached to *Limopsis aurita* (?), Brocchi, and two or three more were dredged by the 'Challenger' Expedition between Sierra Leone (Africa) and Fernando de Noronha (South Atlantic) in 1850 fathoms, associated with *Discinisca atlantica*.

Obs. In 1880, when describing this shell for my 'Challenger' Report, I felt very uncertain with respect to the genus to which it should be referred, especially as I was unacquainted with the shape and character of its loop, and did not like to run the risk of opening the shell and separating its valves. Since then a number of specimens having been obtained during the 'Blake' Expedition, Mr. Dall was able to examine the loop, which he found to be simple and short, as in *Terebratula* or *Terebratulina*; the lateral

portions of the umbo are decidedly auricular, and the character of the radiating riblets would, he thinks, warrant us in classing the shell in the last-named subgenus. It appears to be a good species.

23. *TEREBRATULINA MURRAYI*, Davidson. (Plate VI. figs. 15–17.)

Terebratula Murrayi, Davidson, Proc. Roy. Soc. vol. xxvii. p. 437, 1878; and (*Terebratulina murrayi*) Report of the Brachiopoda, Voyage of H.M.S. 'Challenger,' Zoology, vol. i. p. 39, pl. i., 1880.

Shell small, obscurely trigonal, about as broad as long, widest anteriorly, tapering posteriorly, white; surface of valves marked with about seventeen rounded ribs, of which, in some specimens, a few are due to the interpolation of smaller and shorter ribs between the longer ones, the whole surface crossed also by fine concentric lines of growth. Hinge-line obtusely angular. Ventral valve a little deeper than the dorsal one, beak very slightly incurved, foramen rather large and incomplete, margined laterally by small deltidial plates. In the interior of the dorsal valve the loop is simple. Length 2, breadth 2 lines.

Hab. Dredged by the 'Challenger' Expedition in 1874, lat. 38° 35' S., long. 177° 50' W., near Kermadoc Island, south of Fiji Islands, in a depth of 600 fathoms.

Obs. Some eleven examples of this small shell were dredged by the 'Challenger' Expedition, and none exceeded the dimensions above given. It varied a good deal with respect to the character of its ribs; in some they were all simple, while in others smaller and shorter ribs were interpolated here and there between the longer ones. The crura likewise in some examples in the younger individuals were disunited. In 1879 I sent a specimen to Mr. Dall for examination, and he wrote me, saying that "by devoting about half an hour to this little shell, I have cleared away all the animal matter, leaving the loop perfect. You will see at once that it is a young *Terebratulina*. I suspected this before I could see the loop, from the character of the punctuations, which, you will recollect, is peculiar to the group." I had also previously ascertained that the loop is simple, and that the mantle rises from the bottom of the shell near the loop and adheres to its sides, as was so well illustrated by E. Deslongchamps in *Terebratulina caput-serpentis*.

24. *TEREBRATULINA TUBERATA*, Jeffreys, sp. (Plate VI. figs. 18–20.)

Terebratula tuberata, Jeffreys, Mollusca of the 'Lightning' and 'Porcupine' Expeditions, 1868–70, Proc. Zool. Soc. 1878, p. 401, pl. xxii. fig. 2.

Shell somewhat triangular, broadest and rounded anteriorly, tapering posteriorly, longer than wide. Dorsal valve semicircular; hinge-line nearly straight, shorter than the width of the shell; valve moderately convex and somewhat compressed, lateral sides of umbo auricular. Ventral valve rather deeper than the dorsal one; beak pointed, very little incurved; area triangular; foramen oval-shaped and incomplete, margined anteriorly by the umbo of the ventral valve, laterally by small deltidial plates. Surface of valves traversed by about twenty radiating ribs, of which some are shorter and interpolated between the longer ones. Valves crossed likewise by numerous equidistant concentric raised lines, sometimes slightly projecting and prickly in the young. In the dorsal valve

the crural processes are separated in the young, arched in the adult. Colour dirty white. Length 4, breadth 3, depth 2 lines.

Obs. This small shell was discovered by Dr. J. Gwyn Jeffreys, who described it as a small or young specimen of *Terebratula*, which he had obtained during the 'Porcupine' Expedition in 1870, at a depth of 795 fathoms; and he adds that Prof. Lovén had lent him a perfect example from the Josephine Bank off the Straits of Gibraltar, dredged in 340 to 430 fathoms. Since then several larger and more adult examples were also obtained by the French dredging expeditions in 1882 and 1883, off the western coast of Africa, at a depth of about 4787 metres, and off Morocco and Sahara ('Talisman' Expedition), in 300 to 1261 fathoms.

25. *TEREBRATULINA TRIGONA*, Jeffreys, sp. (Plate VI. figs. 21, 22.)

Terebratula trigona, Jeffreys, Mollusca of the 'Lightning' and 'Porcupine' Expeditions, 1868-70, Proc. Zool. Soc. 1878, p. 402, pl. xxii. fig. 3.

Shell small, ovate, broadly rounded anteriorly, tapering posteriorly, moderately convex, semitransparent between the ribs and glossy; beak short, foramen rather large, incomplete, lateral deltidial plates small; hinge-line obtusely angular. Surface of the valves marked by about eleven simple rounded ribs, of which the three central ones are the largest. Cæcal tubuli numerous. Margins gently curved and scalloped in front. Surface of valves marked with numerous equidistant raised lines. Loop short, simple, and semicircular. Colour clear white. Length scarcely 2 lines, breadth $1\frac{1}{2}$.

Obs. We know so very little about this so-called species that its specific claims cannot be substantiated. I have only seen a single living specimen, that dredged by Mr. Saville Kent in Marshall Hall's yacht 'Norna' off the coast of Portugal, in about 500 fathoms. Dr. J. Gwyn Jeffreys, who described the shell in 1878, says:—"It may perhaps be immature; but I cannot identify it with the young of any other species." In this I feel disposed to agree with Mr. Jeffreys, as I examined the specimen with great attention while drawing the figures for his paper.

Genus *WALDHEIMIA*, King, 1850*.

This excellent genus was founded in 1850 by Prof. W. King for those *Terebratulæ* in which the dorsal valve was provided with a long deeply reflected loop, having its crura attached to the hinge-plate, as is seen in the recent type *Terebratula flavescens*, Lamarck. In the same year Prof. King proposed a genus *Macandrevia* for a shell with a similar loop, in which the umbonal cavity of the ventral valve is furnished with two dental plates, passing somewhat perpendicularly from the dental prominences to the surface of the valve, the umbonal cavity in the dorsal valve being also furnished with

* In 1846, four years prior to the publication of King's name *Waldheimia* for a Brachiopod, the same name had been given by Brullé to an insect; and both Prof. J. Hall and E. Deslongchamps question whether it can consequently be retained for a Brachiopod.

similarly directed plates, and for this genus he selected the recent *Terebratula cranium*, Müller, as his type.

In 1880 M. Douvillé proposed a genus *Neothyris* for another recent species with a similar long loop, of which the *Terebratula lenticularis*, Deshayes, would constitute the type.

I cannot, however, but consider these so-termed genera to be very nearly connected and as mere modifications of *Waldheimia*, and not sufficiently important to warrant their separation into distinct genera.

After a long examination and much thought, I determined to retain the following ten species and one uncertain one:—

1. <i>Waldheimia flavescens</i> , Lamarck.	6. <i>Waldheimia floridana</i> , Pourtales.
2. — <i>kerghuelensis</i> , Davidson.	7. — <i>venosa</i> , Solander.
3. — <i>Grayi</i> , Davidson.	8. — <i>lenticularis</i> , Deshayes.
4. — <i>septigera</i> , Lovén.	9. — (<i>Macandrevia</i>) <i>cranium</i> , Müller.
5. — <i>Raphaelis</i> , Dall.	10. — (—) <i>tenera</i> , Jeffreys.

Uncertain: *Waldheimia Wyvillii*, Davidson.

The characters of both *Waldheimia* and *Macandrevia* will be found fully recorded in our descriptions of *W. flavescens* and *W. cranium*.

Calcareous spicula are absent or almost so in the species above named. Numerous metamorphoses or changes take place in the shape or development of the loop during the different stages of its growth, as will be fully described in the sequel.

26. WALDHEIMIA FLAVESCENS, Lamarck, sp. (Plate VII. figs. 6–19.)

Terebratula flavescens, Valenciennes apud Lamarck, An. sans Vert. vol. vi. p. 246, 1819.

Terebratula australis, Quoy & Gaimard, Voyage de l'Astrolabe, Zool. p. 551, pl. lxxxv. figs. 1–5, 1834.

Terebratula incurva, Quoy & Gaimard, loc. cit. p. 554, pl. lxxxv. figs. 11 & 12, 1834.

Terebratula australis, G. B. Sowerby, Thes. Conch. p. 349, pl. lxix. figs. 25–33, 1846.

Waldheimia australis, King, English Permian Fossils, Pal. Soc. p. 145, pl. xx. figs. 10–12, 1849.

Terebratula australis, Davidson, Classification of Recent Brachiopoda, Ann. & Mag. Nat. Hist. 2nd ser. vol. ix. p. 365, 1852.

Waldheimia flavescens, Davidson, Brit. Foss. Brach. vol. i. p. 64, figs. 6 & 7, 1853; R. Owen, Davidson's Mon. Pal. Soc. vol. i. chapter 1, 1853.

Waldheimia australis, S. Woodward, A Manual of the Mollusca, p. 216, figs. 113 & 114, 1854; Gratiolet, Etudes Anatomiques sur la *Terebratula australis*, Journal de Conchyliologie, 1857; A. Hancock, On the Organization of the Brachiopoda, Phil. Trans. Royal Soc. vol. 148, p. 791, 1858.

Waldheimia flavescens, H. & A. Adams, The Genera of Recent Mollusca, vol. ii. p. 575, vol. iii. pl. cxxx. fig. 3, 1858.

Terebratula (Waldheimia) flavescens, L. Reeve, Monogr. of *Terebratula*, Conch. Icon. pls. 1 & 2, fig. 1 a, b, 1860.

Waldheimia australis, Huxley, An Introduction to the Classif. of Animals, p. 28, fig. 10, 1869.

Waldheimia flavescens, Dall, Amer. Journ. of Conch. vol. vi. p. 108, 1870, and Proc. Acad. Nat. Sci. Philadelphia, 1873, p. 181; Tenison Woods, Proc. Roy. Soc. of Tasmania, 1877, p. 57; Douvillé, Bull. Soc. Géol. de France, 3^e sér. pl. vii., 1879; Davidson, Report on the Brachiopoda, Voyage of H.M.S. 'Challenger,' Zool. vol. i. p. 41, pl. iii. figs. 10–12, 1880; Zittel, Handbuch der Paläontologie, pp. 645

& 702, 1880; E. Deslongchamps, *Études critiques sur les Brachiopodes*, pp. 172 & 177, 1884; Davidson, *Pal. Soc. Brit. Foss. Brach.* vol. v. p. 323, 1884.

Shell longer than wide, ovate or subpentagonal, broadest about the middle, straight in front, yellowish or light brown. Dorsal valve convex, sometimes flattened from about the middle of the valve to the front; ventral valve rather deeper and more convex than the opposite one, with a slightly raised longitudinal flattened elevation or fold along the middle. Beak moderately incurved, truncated by a circular foramen, somewhat separated from the hinge-line by a deltidium; beak-ridges sharply defined. Surface of both valves from the beaks generally smooth up to a certain age, then radiating and irregularly plaited, some ribs being shorter than others, the central one straight, those on the lateral portions of the valves somewhat curved. Shell-structure perforated by canals. In the interior of the dorsal valve the loop is long and simple; after being attached to the hinge-plate and its crural processes given off, the principal stems are outwardly

Fig. 3.

Fig. 4.

Waldheimia flavescens. Interior of both Valves, enlarged.

Fig. 3. Interior of the ventral valve: *f*, foramen; *d*, deltidium; *t*, teeth; *a*, adductor impressions (=occlusors of Hancock); *c*, divaricators (=cardinal muscles of King=muscles diducteurs principaux of Gratiolet); *c'*, accessory divaricators (=muscles diducteurs of Gratiolet); *b*, ventral adjustors (=ventral peduncular muscles, or muscles du pédoncule, paire supérieure, Gratiolet); *b'*, peduncular muscle.

Fig. 4. Interior of the dorsal valve: *c*, *c'*, cardinal process; *b*, *b'*, hinge-plate; *s*, dental sockets; *l*, loop; *q*, crura; *a*, adductor impressions; *ss*, septum.

curved, and on reaching to about three fourths of the length of the valve become reflected. The median septum extends along the bottom of the shell from under the hinge-plate to a little beyond one third of the length of the valve. Brachial or labial appendages largely developed and united to each other by a membrane. The principal lateral branches commence on either side of the mouth, curving outward and facing the bottom of the smaller valve, extend to within a short distance of the front; then becoming suddenly bent back upon themselves to within a short distance of the mouth, are by an elegant semicircular curve directed towards the centre of the larger

valve, and form the commencement of the shorter spiral central lobe. Dimensions very variable, a large well-shaped example measured—length 1 inch 7 lines, breadth 1 inch 4 lines, depth 1 inch.

Hab. *Waldheimia flavescens* (= *australis*) was picked up in great numbers by the 'Challenger' Expedition on the 3rd of June, 1874, at Port Jackson, N. S. W., on the shore, and in from two to ten feet depth of water. Mr. John Brazier, who has dredged extensively in the seas adjoining New South Wales, has kindly sent me specimens from Point Piper, Port Jackson, found under stones in clusters like grapes during low spring-tides, also on *Ostrea glomerata*, Gould, with a specimen of *Kraussina Lamarckiana* on the inside; a young or smooth variety on a piece of shell with *Ismenia* or *Megerlia pulchella*, from the inner north head of Port Jackson, at 10 fathoms, on a bottom of sand and broken shells; likewise from Bottle and Glass Rocks, Port Jackson; and also from off Shark Point, Port Jackson, at a depth of 14 fathoms. Quoy and Gaimard found this species, in 1834, in immense numbers at Port Western, Bass Strait. They observe that hundreds were brought up at each haul of the dredge, either grouped among themselves, or attached to other shells; also at Port Jackson, in four feet of water. Prof. Beete Jukes collected any number while boating in South Australia, among the reefs at Port Jackson; indeed this is one of the commonest species in the locality. The Rev. Tenison Woods observes, in his 'Census of the Marine Shells of Tasmania,' that *Waldheimia flavescens* is found off all South Australia, but only on the north coast of Tasmania.

Obs. I have nothing much to add to my description of this well-known species given in the 'Challenger' Report. The shell has received four or five different specific names, but the best known are those of *flavescens*, Lamarck, and *australis*, Quoy. In 1819, Lamarck having become blind, Valenciennes described for Lamarck the species in question, under the names of *Terebratula flavescens* and *T. dentata*; but he gives no figures of his species, and upon inquiring from Valenciennes, in 1882, I found that it was to a specimen from Port Jackson that he gave the name of *flavescens*. The larger number of malacologists have preferred the name of *australis*, given to it in 1834 by Quoy and Gaimard, who gave a number of good illustrations of the shell, including that of the loop*. Quoy's *T. recurva* is no more than a short variation in form of a specimen in which the beak is much incurved; and the same may be said of Lamarck's *T. dentata*.

Waldheimia flavescens is very variable in shape and ribbing. In some examples the beak is longer or more produced than in others, and the foramen much more distant from the hinge-line in some individuals than in others. Some specimens are also more or less subpentagonal, others elongated oval, some lozenge-shaped and circular. The valves in some examples are quite smooth in the young, and very often up to an advanced stage of growth. In other specimens the ribs are simple or bifurcating, and of different size and strength on the same individual, and begin to rise close to, or at some distance from, the extremity of the beak and umbo. In Pl. VII. some of these modifications in shape and character have been illustrated.

* Voyage de l'Astrolabe, Atlas, pl. 85 (1834).

The loop passes likewise through a series of metamorphoses from the very young state up to the period when it assumes the simple adult *Waldheimia*-character, as has been so elaborately described and illustrated by Herman Friele in the case of *Waldheimia septigera* and *Waldheimia* (or *Macandrevia*) *cranium*.

The intimate shell-structure of *Waldheimia flavescens* has been minutely described and admirably illustrated by Dr. W. B. Carpenter, in his chapter "On the Intimate Structure of the Shells of the Brachiopoda," which he kindly prepared in 1853 for vol. i. of my 'Monograph on British Fossil Brachiopoda,' and to which the reader is referred. But we may here mention that he found the perforations in *W. flavescens* to average a diameter of about $\frac{1}{800}$ inch, and the distance of their centres about $\frac{1}{400}$. Three of his illustrations have been given in Pl. VII. figs. 17-19.

In 1869, in vol. xxiv. of the Transactions of the Royal Irish Academy, Prof. W. King gave an excellent description of the shell-structure of the species under description, in which he observes that Quekett, in 1850, discovered that the extremity of the cæcal appendages, as they are now generally called, is more or less flattened or disk-shaped, and encircled with a fringe of exceedingly minute radiating lines or membranous filaments. Prof. King adds, "In position and arrangement the *filaments* belonging to the terminations of the cæcal appendages so completely agree with the *radiating lines* which intersect the apertural rims of the perforations, as to leave no doubt on my mind that the latter are *tubular* and enclose the former." The shell of *W. flavescens* and its intimate shell-structure is also described by Gratiolet, in his admirable memoir "Études anatomiques sur la Térébratule australe," Journal de Conchyliologie, 1857. He says, at p. 214 of his memoir, that the shell, when divested of the foreign objects that generally cover its surface, is of a tolerable transparency. One notices in it, besides the concentric lines of growth, diverging costæ which give to the sharp edge of the valves an undulated appearance; examined with a lens, its surface is finely granulous, so much so, that one might think it formed of very fine granulations; but this apparent granulation is due to a multitude of microscopic perforations with which the shell is riddled. These perforations are, besides, very irregularly disposed on parallel bands which intercept quadrilateral spaces. He adds that the substance of the shell is formed of small calcareous prisms, somewhat attenuated at their extremity and disposed in parallel layers, which give to the fractured shell a fibrous aspect; these elements are closer together towards the edges of the shell than in their central portions.

The soft parts of the animal of *Waldheimia flavescens* have been admirably described and elaborately illustrated by several eminent anatomists. First by Owen in 1853, in the Introduction to vol. i. of my work on 'British Fossil Brachiopoda;' subsequently, in 1857, by Pierre Gratiolet, in his memoir above quoted, and in the following year by Albany Hancock, in his classical memoir "On the Organization of the Brachiopoda" (Phil. Trans. Roy. Soc. vol. cxlviii.). To these works the reader is referred for more complete anatomical details than we are able to reproduce in this monograph.

The body proper is small when compared with the size of the shell, and has both its valves lined by a delicate bilobed integument or membrane, termed the 'pallium' or mantle; this secretes the shell and is fringed with horny bristles. The mantle is composed

of an outer and an inner layer, between which are situated the blood-channels or lacunæ. There are four principal arterial trunks in the dorsal lobe of the mantle, and these run direct to the front and bifurcate at intervals (see Pl. VII. figs. 15, 16 *d*). The aperture of the mouth is situated in the middle line between the pallial lobes, and on each side of it is a prolongation of the body provided with ciliated tentacula.

The muscles have been minutely described by the three anatomists above named. Five or six pairs are connected with the opening and closing of the valves, or with their attachment to, or movements upon, the peduncle. First of all the *adductor* or *occluser* consists of two muscles, which, bifurcating near the centre of the cavity, produce a large quadrangular impression on the internal surface of the dorsal valve. The function of

Waldheimia flavesceus (after Hancock).

M, ventral valve; N, dorsal valve; l, loop; v, mouth; z, extremity of intestine; a, adductor; c, divaricator; c', accessory divaricator; b, ventral adjustor; b', peduncular muscle; b'', dorsal adjustor; P, peduncle.

this pair of muscles is the closing of the valves. Two other pairs have been termed *divaricators* by Hancock, and have the function of opening the valves. The divaricators proper are stated by the same eminent authority to rise from the ventral valve, one on each side, a little in advance of and close to the adductors, and after rapidly diminishing in size, become attached to the cardinal process, a space or prominence between the sockets in the dorsal valve. The accessory divaricators are a pair of small muscles which have their ends attached to the ventral valve, one on each side of the median line, a little behind the united basis of the adductors, and again to the extreme point of the cardinal process. The two pairs of muscles, apparently connected with the peduncle and their limited movements, have been minutely described by Hancock as having one of their extremities attached to this organ. The dorsal adjustors are attached to the ventral surface of the peduncle, and are again inserted into the hinge-plate of the smaller valve. The ventral adjustors are considered to pass from the inner extremity of the peduncle, and to become attached by one pair of their extremities to the ventral valve, one on each side of and a little behind the expanded base of the divaricators. The function of these muscles is not only that of erecting the shell, but also that of attaching the

peduncle to the shell, and thus controlling the steadying of it upon the peduncle. These details and the figure taken from Hancock will sufficiently explain the functions of the different muscles. Hancock describes the peduncle as composed of a dense muscular, semi-cartilaginous mass of a cylindrical form.

The nature of the muscles has also been well described and illustrated by Owen, Gratiolet, and more recently by Van Bemmelen, in his anatomical memoir 'On the Structure of the Shells of Brachiopods and Chitons' *.

Owing to the strong and tight interlocking of the valves of *Waldheimia flavescens* and other species, by means of curved teeth and sockets, the Brachiopoda would seem to open their valves very slightly.

At p. 806 of the admirable memoir already alluded to, Hancock says that in *Waldheimia flavescens* and in other forms of the genus, the "loop originates in the hinge-plate in two necks or processes,—the crura, from which two points project downwards, called the crural processes. The upper or dorsal members of the loop, passing from these two lateral necks, stretch forward for about two thirds the length of the valve, then bend towards the ventral valve, and turning back upon their course, are united across the median line, a short way in advance of the crural processes. The lateral portions of the loop are curved a little outwards. The whole of this calcareous support, including the crura and crural processes, is a product of the inner lamina of the dorsal pallial lobe. This lamina, with the exception of a portion at each side, which is continuous with the similar lamina of the ventral lobe, forming with it the anterior wall of the perivisceral chamber, is turned downwards and forwards, and extends as far as the transverse portion of the loop. It then divides into two lobules, one passing to the anterior extremity of each of the lateral portions of this calcareous support, binding together the dorsal and ventral members. The loop in its sinuous course follows the margins of this bifurcated pallial process, and lies imbedded in its substance.

"The arms themselves taper to fine points, having taken their origin at each side of the mouth, which is situated at the back of the pallial chamber in a central position, opening downwards. They pass outwards and forwards in connection with the calcareous loop and the pallial lobules, running along the outer edge of the dorsal member of the former; they then turn back upon the reflected or ventral part of it, and on reaching the transverse portion bend inwards and doubling sharply upon themselves, again advance and go to form two vertical spirals turned towards the dorsal valve. The spirals are placed parallel to each other, and throughout the coil, consisting of two or three turns, they are united across the median line by a rather wide, stout, semicartilaginous membrane, which is attached behind the transverse portion of the loop."

The labial appendages have been magnificently illustrated by Owen in plates 1 & 2 of his chapter on the 'Anatomy of the *Terebratulina*' already referred to. They are also minutely described and illustrated by Gratiolet in his 'Études anatomiques sur la *Térébratule australe*,' 1857.

These appendages are, in *Waldheimia flavescens*, as well as in other species of the

* 'Over den Bouw der Schelpen van Brachiopoden en Chitonen,' Leiden, 1882, and of which an English note will be found in the Ann. & Mag. Nat. Hist. 5th ser. vol. xi. p. 379, 1883.

genus, a pair of very remarkable organs, eminently characteristic of the Brachiopoda. They are often and more correctly termed *labial* appendages, on account of each member being a prolongation of the lateral portions of the lips of the margins of the mouth. They occupy the larger portion of the cavity of the shell in front of the visceral chamber;

Fig. 6.

Fig. 7.

Waldheimia flavescens.

Fig. 6. Interior of dorsal valve, to show the position of the labial appendages. (A portion of the fringe of the cirri has been removed to show the brachial membrane and a portion of the spiral extremities of the arms.) Enlarged.

Fig. 7. Longitudinal section, with a portion of the animal. *d, h*, brachial appendages; *a*, adductor; *c, c'*, divaricator muscles; *s*, septum; *v*, mouth; *z*, extremity of alimentary tube. (The peduncular muscles have been purposely omitted.) Enlarged.

and are mainly composed of a membranous tube fringed on one side with long flexible hollow cirri, which are not capable of being protruded in those families and genera in which they are folded back upon themselves and supported by a skeleton, as in *Waldheimia*. Hancock says (Phil. Trans. vol. cxlviii. p. cxliii.) that the arms or labial appendages "are normally composed of a membranous tube or canal bearing a semicartilaginous grooved ridge. The latter stretches from end to end of the former, and gives support to the fringe of cirri. As far back as the commencement of the spirals the arms are as above stated; but for the entire length of the lateral portions of the loop, where the arms are doubled upon themselves, and where, of course, two tubes or canals might have been expected, there is only one, the two having, as it were, coalesced. These large canals at the roots of the arms are continuous with those of the spirals, and terminate in blind sacs, one at each side of the œsophagus, close to the mouth. On making a transverse section of this part of the arm, the enlarged terminal portion of the brachial canal is seen connected with the external edges of the dorsal and ventral members of the loop; and the pallial lobule, stretching between the inner edges, forms a sort of inner tube. This inner tube opens widely into the perivisceral chamber, is in fact a prolongation of

this chamber, and terminates at the anterior extremity of the lateral portion of the loop, forming what may be designated as the brachial pouches. The cirri are undoubtedly contractile to some extent, and are also endowed with the power of motion individually, as proved by the observations of Mr. Barrett. It would likewise appear that the entire fringe of cirri can be elevated or depressed, there being ample means provided for the purpose." Many other highly important details have been recorded by Owen, Gratiolet, and Hancock, which I regret I am unable to reproduce.

Mr. Hancock states, at p. 811 of his memoir, that "the mouth, in all the Brachiopods, as previously stated, is situated in the brachial groove, at the posterior junction of the arms, and is a simple, oval, transverse slit, or orifice devoid of any armature. In *W. australis* there is, however, a slight bulging-out of the posterior wall immediately within the orifice, which is somewhat like the valvular appendage of *Plumatella*, and may perhaps assist in swallowing. The alimentary tube assumes the form of a siphon bent in the vertical plane, the arch turned towards the dorsal valve; the œsophagus represents the short, the stomach and intestine the long arm."

There is no anal outlet in this species. According to the same author, the mesentery is divided into two portions, the dorsal and ventral: it sustains the alimentary canal in the vertical plane, and altogether with it divides the anterior portion of the chamber into lateral halves. The gastro-parietal bands are three in number, two lateral and one median.

In *W. flavesceus*, Mr. Hancock says, "the genitalia are formed of thick bands,

Fig. 8.

Lateral view of the Viscera of *Waldheimia flavesceus*, after Hancock.

a, anterior layer of mantle; *b*, posterior layer; *c*, anterior walls of the body between the mantle-lobes; *d*, arms; *p*, gullet; *q*, stomach, with cut biliary ducts of the left side; *r*, right hepatic mass; *s*, intestine ending caecally between *j* and *k*; *v*, so-called "auricle" of the right "pseudo-heart," the left being almost wholly removed; *w*, pyriform vesicle fixed to the back of the stomach, and probably performing the function of a true heart; *z*, œsophageal ganglia. (This figure is also given by Huxley on page 28 of his 'Introduction to the Classification of Animals,' 1869.)

somewhat convoluted and branched; they are of a full yellow colour, and are thrust into the trunks and main branches of the great pallial sinuses" (p. 817).

"The heart is a simple, unilocular, pyriform vesicle, suspended from the dorsal aspect, and projects freely into the perivisceral chamber" (*loc. cit.* p. 834).

Prof. Huxley, in his 'Introduction to the Classification of Animals,' 1869, says that "The precise characters of the true vascular system of the *Brachiopoda* probably require still further elaboration than they have yet received; and the same may be said, notwithstanding the valuable contributions of F. Müller and of Lacaze-Duthiers, of their development; but the shell, the pallial lobes, the intestine, and the nervous and the atrial systems, afford characters amply sufficient to define the class" (p. 30). He also says (p. 29), "In all *Brachiopoda* which have been carefully dissected, a singular system of cavities and canals situated in the interior of the body, but in free communication with the surrounding medium, has been discovered. This, which I shall term the 'atrial' system (from its close correspondence with the system of cavities, which has received the same name in the Ascidians), has been wrongly regarded as a part of the true vascular system, and the organs by which it is placed in communication with the exterior have been described as 'hearts.' There are sometimes two and sometimes four of these 'pseudo-hearts,' situated in that part of the body-wall which helps to bound the pallial chamber. Each pseudo-heart is divided into a narrow, elongated, external portion (the so-called 'ventricle'), which communicates, as Mr. Hancock has proved, by a small apical aperture with the pallial cavity; and a broad, funnel-shaped, inner division (the so-called 'auricle'), communicating, on the one hand, by a constricted neck with the so-called 'ventricle,' and on the other, by a wide, patent mouth, with a chamber which occupies most of the cavity of the body proper, and sends more or less branched diverticula into the pallial lobes. These have been described as parts of the blood vascular system; and the arterial trunks, which have no existence, have been imagined to connect the apices of the ventricles with vascular networks of a similarly mythical character, supposed to open into the branched diverticula. In fact, as Mr. Hancock has so well shown in his splendid and exhaustive memoir published in the 'Philosophical Transactions' for 1857, the true vascular system is completely distinct from this remarkable series of 'atrial' chambers and canals, the function of which would appear to be to convey away excretory matters and the products of the reproductive organs, which are developed in various parts of the walls of the atrial system."

In an exquisitely beautiful enlarged illustration in his 'Anatomy of the *Terebratula*,' Owen represents the brachial aponeurosis and spiral arms of *Waldheimia flavescens*, showing the central part of the nervous system, with the brachial and the beginning of the pallial nerves.

27. WALDHEIMIA VENOSA, Solander, sp. (Plate VIII. figs. 1-5; Plate IX. fig. 1.)

Anomia venosa, Solander, G. Dixon, A Voyage round the World, Appendix no. 1, p. 355 and fig., 1789.

Terebratula globosa, Valenciennes, apud Lamarck, An. sans Vert. vol. vi. p. 246, 1819, with reference for figure to Encyclop. Méthod. p. 239, figs. 5 a, b, 1789.

Terebratula dilatata, Valenciennes apud Lam., An. sans Vert. vol. vi. p. 245, 1819.

Terebratula Gaudichaudi, Blainville, Diet. Sci. Nat. 1824.

Terebratula globosa, Anton, Verzeichniss der Conchylien, p. 23, 1839.

Terebratula eximia, Philippi, Küster, Conch.-Cab. vol. vii. p. 39, pl. 2. figs. 9 & 10, 1843.

Terebratula Fontaineana, d'Orbigny, Voy. Amér. Mérid. vol. v. p. 675, pl. 85. figs. 30 & 31, 1847.

Terebratula (Waldheimia) globosa, Reeve, "Monogr. of *Terebratula*," Conch. Icon. pl. ii. fig. 3, 1860, and pl. vi., 1861.

Terebratula (Waldheimia) globosa, Reeve, Ann. & Mag. Nat. Hist. vol. vii. p. 173, 1861.

Terebratula (Waldheimia) dilatata, Reeve, Conch. Icon. pl. vi. fig. 2, 1861.

Terebratula physema, Valenciennes, MS. ; Reeve, Conch. Icon. pl. vi. fig. 23, 1861.

Waldheimia venosa, Davidson, Ann. & Mag. Nat. Hist. 3rd ser. vol. viii. p. 36, 1861.

Terebratula venosa, Cunningham, Voyage of H.M.S. Nassau, Trans. Linn. Soc. vol. xxvii. p. 488, 1871.

Waldheimia venosa, Dall, Amer. Journ. of Conch. vol. vi. p. 109, 1870, and Proc. Acad. Sci. Philadelphia, p. 182, 1872.

Shell large, longitudinally oval, ovately globose, widest about the middle, longer than wide. Dorsal valve convex, with a shallow flattened longitudinal depression or sinus along the middle, commencing at about half the length of the valve and extending to the front. Ventral valve a little more convex than the opposite one, with a slightly produced and flattened mesial fold commencing near the beak and extending to the front; beak moderately incurved and truncated by a rather large circular foramen, with thickened margin and completed anteriorly by a deltidium in two pieces; beak-ridges sharply defined. Surface of valves smooth, marked only by concentric lines of growth. Shell-structure punctate. Colour light olive-horny. Loop long and reflected; cardinal process and hinge-plate large. Under the hinge-plate a rather strong mesial septum extends to a third or a little more of the length of the valve, and on either side on the bottom of the shell are situated the adductor and other muscular scars. Length 3 inches 2 lines, breadth 2 inches 8 lines, depth 2 inches.

Hab. Tierra del Fuego; Falkland Islands (Dixon); Port William, Falkland Islands (B. J. Sullivan); Coquimbo; off Cape Possession in about 15 fathoms; Port Famine (Cunningham).

Obs. This is the largest recent Brachiopod at present known, and its history has been somewhat confused. It was, however, well described and figured in 1789 by Captain George Dixon, who says, at p. 355 of his work 'A Voyage round the World':—"At Falkland Islands we met with a curious kind of shell of the *Anomia* genus of Linnæus, of which, though the species are numerous in a fossil state in most parts of the world, few have been discovered recent, or fresh from the sea. Only one of this sort was before known in Europe, which was brought over by my late worthy commander, the much regretted Captain Cook, in his first voyage round the world. It was in the Portland Museum, and was named by the late celebrated Dr. Solander, in his MSS. description of the shells of that splendid cabinet, *Anomia venosa*; the specimen is now in the collection of Mr. Calonne, of London. This kind (as do all that are properly of this genus) adheres to coral rocks by a ligament that comes from the animal through the hole in the larger valve. The internal structure peculiar to the shells of this genus is very singular, and consists of two testaceous rays, which commence near the hinge in the lesser valve, where they adhere; from whence, leaving the shell, they

proceed to near the edge, then bend towards the other valve, and turn back to their commencement, where they unite. This internal part is very delicate, and breaks upon the smallest touch, but is thicker in the parts nearest to the large valve. The shell takes its name from certain parts of the animal, which run in a branched form along the inside of the shell, which being held to a strong light, or a candle, gives a beautiful veined appearance. The outside is smooth and of a pale brown colour. The specimen from which the engraving was made, is in the private collection of Mr. George Humphrey, dealer in natural curiosities, London" (Plate VIII. fig. 1).

In 1819 Valenciennes briefly described the same species by the name of *Terebratula globosa*, referring at the same time to plate 239, fig. 2, of the 'Encyclopédie Méthodique' for a figure of his species. The figure in the 'Encyclopédie' is not well drawn; but the specimen from which it was taken is in the collection of the Jardin des Plantes, and was lent by M. Valenciennes to Lovell Reeve, who gave a good figure in plate vi. fig. 3 of his 'Monograph of *Terebratula*' in 1861. In 1843 the same species received the name of *T. eximia* from Philippi and Küster; that of *T. Fontaineana* from d'Orbigny, and to a large specimen of the same form in the collection of the Jardin des Plantes, measuring $2\frac{1}{2}$ inches in length, M. Valenciennes gave the MS. name of *physema*. Under this name it is also described and figured by Reeve in his 'Monograph of *Terebratula*' in 1861. The specimen was brought from Coquimbo by M. Gaudichand in 1833 (Plate VIII. fig. 6). In his description of this specimen Mr. L. Reeve observes that "It would be satisfactory if the species could be confirmed by the discovery of further specimens. It is intermediate in its characters between *T. dilatata* and *T. globosa*, inclining rather to the latter species, of which it may prove to be a colossal broadly inflated variety," and this I have no doubt it is.

It was Valenciennes, however, who described and gave the names to all the recent Brachiopoda published in 1819 in the 'Animaux sans Vertèbres' of Lamarck, that celebrated zoologist having unfortunately lost his sight at that period.

I am quite convinced, after the inspection of the original specimen and figure of *T. dilatata*, Val., as well as from Reeve's figures in plate vi. fig. 2 of his 'Monograph' (the type having been lent to him by M. Valenciennes out of the collection of the Jardin des Plantes), that it is only a smaller and more transverse form of *Waldheimia venosa*. Several other specimens of the same form which have also passed through my hands have tended to confirm this view, which had already been expressed by Mr. W. H. Dall in his memoir published in the 'American Journal of Conchology,' vol. vi. p. 109, where he notes also that the *T. Gaudichaudi* of Blainville is another synonym of *W. venosa*. In a paper I published in the 'Ann. & Mag. Nat. Hist.' for August 1867, I described and figured the largest example of *W. venosa* that had been discovered, which had been dredged in 1843 or 1844 by Rear-Admiral B. J. Sullivan at Falkland Islands, in the same locality whence the type of the species had been procured by Dixon. Admiral Sullivan informed me, when lending me his specimen, that the depth at which the animal lived was from 6 to 7 fathoms, the bottom on which the shell lay being a compact quartzose sand only, as no mud ever came up with the dredge, although a stiff muddy clay underlies the sand, on which anchors hold very firmly.

Waldheimia venosa, like all its congeners, is variable in external shape. The foramen is larger in some specimens than in others. As is the case with *W. lenticularis* when quite young and sometimes until half its growth, *W. venosa* is either almost circular or transversely oval. *W. dilatata* seems to be a half-grown individual with a large foramen (Plate IX. fig. 1).

W. venosa has often been confounded with *Laqueus californicus* of Koch, both species attaining large dimensions; but the loop in both is entirely different, and in the last-named species the beak is much more incurved, the foramen very small, and separated to a greater or less extent from the umbo of the dorsal valve by a deltidium in two pieces.

In Vol. clxviii. of the Phil. Trans., 1879 (Transit of Venus Exped.), Mr. Edgar A. Smith mentions *Waldheimia* (*Terebratula*) *dilatata*, Lamarck, as having been obtained at Observatory Bay at a depth of 4 fathoms. I may, however, remark that the 'Challenger' Expedition did not bring back a single specimen of Solander's species.

28. WALDHEIMIA LENTICULARIS, Deshayes, sp. (Plate IX. figs. 2-13.)

Terebratula lenticularis, Deshayes, Revue Zoologique par la Soc. Cuvérienne, p. 359, 1839, and Mag. de Zoologie, d'Anatomie comparée et de Paléontologie, p. 41, figs. *a, b, c, d*, 1841; G. B. Sowerby, Thes. Conchyl. vol. i. p. 360, pl. lxxii. figs. 108-110, 1846; Davidson Ann. & Mag. Nat. Hist. 2nd ser. vol. ix. p. 365, 1852; L. Reeve, Monogr. of *Terebratula*, Conch. Icon. pl. 2. fig. 4, 1860.

Waldheimia lenticularis, Gray & Woodward, Brit. Mus. Cat. of the Brachiopoda, p. 58, 1853; Dall, Proc. Acad. Nat. Sci. Philadelphia, p. 182, 1873.

Neothyris lenticularis, Douvillé, Bull. Soc. Géol. de France, 3^e sér. t. vii., 1879.

Waldheimia lenticularis, E. Deslongchamps, Notes sur les Modifications à apporter à la Classe des Terebratulidæ, pl. xvii. figs. 5-7, 1880.

Shell large, orbicular, elongated oval, globose, longer than wide, thick, rounded laterally, less so in front. Dorsal valve uniformly convex or slightly depressed at or close to the frontal margin. Ventral valve deeper, and a little more convex than the dorsal one and slightly longitudinally carinated; beak rather sharply incurved, overlying the umbo of the opposite valve, and truncated by a small circular foramen, separated from the hinge-line by a wide, narrow, concave deltidium in two pieces; beak-ridges very sharply defined. Surface smooth, marked by concentric lines of growth, punctate. Colour pinkish red or yellowish horny. Loop simple, long, and reflected; cardinal process rather large and prominent; under it a mesial septum extends to about one third of the length of the valve on either side the adductor and other muscular scars. Length 2 inches 2 lines, breadth 1 inch 11 lines, depth 1 inch 3 lines.

Hab. Lives abundantly attached to rocks in the Straits of Foveau, New Zealand, in 15 fathoms. It occurs also in great numbers in the Pleistocene rocks of New Zealand.

Capt. F. W. Hutton, in his 'Catalogue of the Tertiary Mollusca of New Zealand,' p. 35, 1873, quotes the following localities:—"Wanganui, The Deans, Waipara, Waitotari, Chatham Islands." The fossil is also well described and figured by Prof. E. Suess from the 'Novara' Expedition, in his fine memoir "Fossile Mollusken und Echinodermen aus Neu-Seeland," Palæontologia, p. 56, pl. x. figs. 3, 4 (1866).

Obs. I have in my possession a very large series of specimens of this fine species from

2 up to 23 lines in length. When young and up to nearly half its growth the shell is almost circular and broadest anteriorly, tapering at the beak, and nearly as wide as long; with age the shell becomes elongated and gibbous.

I have carefully observed and drawn the modifications that take place in the shape of the loop, of which I give some illustrations in Plate IX. figs. 11, 12, 13. My figures agree well with those published by M. E. Deslongchamps in his 'Etudes Critiques' above quoted, and they do not differ from those peculiar to other species of the genus. In a specimen two lines in length the whole loop is small, not extending further than to about half the length of the valve, its principal lamellæ, as well as the reflected portion, becoming attached to the extremity of the upper sides of a perpendicularly elevated septum, as in *Magasella*. After a time the septum becomes less elevated, the principal stems of the loop acquire greater length, and give off at about half their length a horizontal lamella, which is also fixed to the septum, as in *Terebratella*; later on this horizontal lamella detaches itself from the septum, the principal stems of the loop become more separate, and the reflective portion assumes the appearance of that of a full-grown *Waldheimia*.

Waldheimia lenticularis was well described and figured by Deshayes in 1841, and also by G. B. Sowerby and L. Reeve. It has had the good fortune to preserve the same specific name, although located in the genera *Terebratula*, *Waldheimia*, and *Neothyris*. As observed by Reeve, in his 'Monogr. of *Terebratula*,' "Were it not that the habitats of this and the preceding species [*Terebratula* (*Waldheimia*) *globosa*] are well authenticated, it would be difficult to believe that there is any specific difference between them. This is rather lighter [with an] inflated growth, and the colour is remarkable, but the characters in other respects are the same in detail throughout. The foramen is probably as small as in *T. globosa*; in all the specimens of *T. lenticularis* that I have seen it is more or less eroded, according to the circumstances of its attachment. The loop is precisely the same in both species." There are, however, differences between *W. venosa* and *W. lenticularis* which must not be overlooked. In well-preserved specimens of the last-named shell the foramen is small, in fact much smaller than in *W. venosa*, and separated from the hinge-line by a well-developed deltidium in two pieces, in this respect more nearly resembling *Laqueus californicus*. The beak is more incurved, and the dorsal valve more uniformly convex. The colour is also very different in the two species. It is also a smaller shell.

29. WALDHEIMIA KERQUELENENSIS, Davidson. (Plate X. figs. 7-17.)

Waldheimia kerguelensis, Davidson, Proc. Roy. Soc. vol. xxvii. p. 437, 1878, and (*W. kerguelensis*) Report on the Brachiopoda, Voyage of H.M.S. 'Challenger,' Zool. vol. i. p. 40, pl. iii., 1880.

Shell ovate, ventricose, longer than wide, light yellowish white, smooth. Dorsal valve most convex near the umbo, a slight mesial depression commencing about the middle of the valve extends to the front, and is more or less distinctly margined on either side by a faint raised line or ridge; front line slightly depressed or nearly straight. Ventral valve rather more convex than the opposite one, and more or less distinctly keeled by the presence of a wide, convex, but slightly raised fold, which corresponds with the depression in the dorsal valve. Beak moderately produced and incurved, truncated by a small

circular foramen laterally margined by two small disunited deltidial plates. Cardinal process prominent, and formed of three distinct parts. In the interior of the dorsal valve, and under the incurved extremity of the umbo and hinge-plate, there rises a wide short and massive septum, and on either side on the bottom of the valve the muscular scars are visible. Loop delicate, elongated and reflected. Shell-perforations numerous. Length 2 inches, breadth 1 inch 4 lines, depth 1 inch 2 lines.

Hab. It was dredged by the 'Challenger' Expedition in great numbers off Marion Island, west of Kerguelen Island, at a depth of 100 fathoms. Two examples of *Platydia anomioides* were attached to one of the specimens. Also at Balfour Bay, near Kerguelen Island, in 20 to 60 fathoms, south of the same island at a depth of 150 fathoms. Three examples of *Rhynchonella nigricans*, var. *pixidata*, were obtained with it.

Obs. I have examined several specimens of the animal that had been dredged alive by the expedition, as well as of the shell from 1 line up to 24 lines in length. Some examples were nearly circular and as broad as long, but the larger number were of an elongated oval shape, becoming ventricose with age. When the peduncle by which the shell adhered to foreign bodies was sufficiently long, and did not interfere with the animal's limited movements, the beak was much incurved and the foramen small; but in most cases the peduncle was exceedingly short, and the shell came into contact with the hard bodies to which it was moored, causing erosion and the consequent enlargement of the foraminal aperture. The animal closely resembles in general character that of *Waldheimia flavescens*, so admirably described by Owen and Hancock.

Some specimens bear a certain resemblance to *Waldheimia lenticularis*; but this last is smaller, more ovate or regularly oval, and more especially in the young and intermediate stages of growth. Having forwarded two examples for Mr. Dall's examination, he informs me:—"I have carefully compared it with d'Orbigny's *Waldheimia Fontaineana*, and feel more sure than ever of the correctness of my reference of his species to *Waldheimia venosa*. It is certainly not this fine species (*Waldheimia kerguelenensis*)," though it is possibly identical with the shell erroneously described and figured by G. B. Sowerby, at p. 359, and plate xxi. figs. 99-101, of his 'Thesaurus Conchyliorum,' in 1846, as the *Terebratula globosa*, Lamarck. The specimen Sowerby figured under that name (said to have been taken from Lamarck's collection) is now in the British Museum, and Sowerby was certainly mistaken in stating "It agrees perfectly with the representation in the 'Encyclopédie Méthodique,' tab. 339. fig. 2"!

30. WALDHEIMIA GRAYI, Davidson. (Plate X. figs. 1-4.)

Terebratula Grayi, Davidson, Proc. Zool. Soc. 1852, p. 76, pl. iv. figs. 1-3.

Terebratula (Waldheimia) Grayi, L. V. Schrenck, Reisen und Forschungem in Amur-Lande, p. 465, 1854-56.

Waldheimia Grayi, E. Suess, Sitzungsab. Akad. Wissenschaften, Bd. xxxvii. p. 201, 1859.

Terebratula (Waldheimia) Grayi, L. Reeve, Conch. Icon. pl. 2. figs. 5 a, b, c, 1860, and Ann. & Mag. Nat. Hist. 3rd ser. vol. vii. p. 175, 1861; Journal de Conchyl. p. 123, 1861.

Waldheimia Grayi, Carpenter, Rep. Brit. Assoc. 1863, p. 636.

Waldheimia Grayi, A. Adams, Ann. & Mag. Nat. Hist. 3rd ser. vol. xi. p. 99, 1863; Davidson, Proc. Zool. Soc. p. 304, pl. xxxi. figs. 7 & 8, 1871; Dall, Am. Journ. of Conch. vol. vi. p. 110, 1870, and Proc. Acad. Sci. Philadelphia, 1873, p. 182; G. Dunker, Index Moll. Maris Japonici, p. 252, 1882.

Shell variable in shape, squarely subpentagonal or subtriangularly semicircular, longer or broader than wide. Hinge-line either nearly straight or very obtusely angular, shorter or longer than the breadth of the shell; lateral margins rounded, nearly straight or gently curved in front. Dorsal valve very moderately convex, somewhat flattened and more or less longitudinally depressed along the middle. Ventral valve very much deeper and more convex than the dorsal one; beak short, bent backwards, nearly straight or very little incurved, and obtusely truncated by a large, transversely oval, incomplete foramen, always more or less eroded, margined anteriorly by the dorsal valve and by two small labial rudimentary deltidial plates; beak-ridges strongly marked, leaving between them on the hinge-line a wide triangular almost flat area. Surface of valves marked by a number of strong bifurcating radiating angular costæ, of which the central one is usually the largest. Valves traversed by numerous more or less prominent scale-like zigzag concentric lines or ridges of growth. Shell-structure punctate. Colour yellowish, with concentric bands of crimson colour, deeper in tint at or near the projecting concentric ridges of growth. Proportions very variable. Two large examples measured:—

Length 1 inch 4 lines, breadth 1 inch 3 lines, depth 10 lines.

„ 1 „ 2 „ „ 1 „ 7 „ „ 9 „

In the interior of the dorsal valve the cardinal process is wide and narrow, the hinge-plate large, mesial septum strong, extending to about half the length of the valve. Loop long and simple; the principal stems attached to the base of the hinge-plate extend to about four fifths of the length of the valve before becoming reflected.

Hab. Hakodadi, Mososeki, Japan, in 7 fathoms (Adams). Strait of Corea (Belcher and St. John), in 37 fathoms. Dall mentions also Catalina and Monterey.

Obs. This very remarkable and beautiful shell is, as already stated, very variable in shape, so much so, indeed, that some malacologists have felt disposed to divide it into two species; and, as observed by Dall, it is sometimes hardly possible to distinguish it exteriorly from some specimens of *Terebratula transversa*, Sow., var. *caurina*, Gould, or *T. occidentalis*, Dall, except by the loop, which in *W. Grayi* is simple, while in the others it is three times attached, as in *Terebratella*.

Prior to 1852, the shells of the species under description had erroneously been referred to the *Terebratula* (*Kraussina*) *rubra* of Pallas, to which they also bear some external resemblance; and it was only after careful examination of the interior calcareous support of the labial appendages that I discovered that they belonged to two very distinct genera. As observed by Lovell Reeve, in his 'Monograph of *Terebratula*,' the foramen of the shell naturally becomes more or less eroded and enlarged according to the roughness with which the creature is buffeted about in the place of its attachment, owing to the shortness of its peduncle. He also observes that the deltidial plates meet in the middle in very young examples, and that the ribs of *W. Grayi* are not really scaled, but scales

are formed on them by the overlapping of the concentric lines of growth. The ribs are also somewhat irregular in some specimens.

Waldheimia Grayi is extremely abundant in its favourite haunts. In Corean waters it was dredged in great abundance by Admiral Sir Edward Belcher during the voyage of H.M.S. 'Samarang.' A. Adams and Capt. St. John obtained it in similar abundance, and say that it could be collected by thousands in certain places where it had been washed on shore. Mr. Dall has observed that the blood in this species is red.

31. *WALDHEIMIA SEPTIGERA*, Lovén, sp. (Plate XI. figs. 1-10.)

Terebratula septigera, Lovén, Index Molluscorum Litora Scandinaviæ occidentalia habitantium, Öfversigt af K. Vet.-Akad. Förh. 1846, no figure.

Waldheimia septigera, Gray & Woodward, Cat. of Brach. in the Brit. Mus. p. 59, 1852; Davidson, Ann. & Mag. Nat. Hist. 2nd ser. vol. xvi. pl. x. fig. 1, 1855; E. Suess, Ueber die Wohnsitze der Brachiopoden, Sitzungsber. k. Akad. Wissensch. Wien, Bd. xxxvii. p. 200, 1859.

Terebratula (Waldheimia) septigera, Reeve, Monogr. of *Terebratula*, Conch. Icon. pl. iii. fig. 10, 1860.

Terebratula peloritana, Seguenza, Notizie succinte, p. 19, 1862.

Waldheimia peloritana, Seguenza, Memorie della Soc. Ital. di Scienze Naturali, vol. i. p. 49, pl. vi. fig. 10, 1865.

Terebratula septata, Jeffreys (not Philippi), Brit. Conch. vol. ii. p. 14, 1863.

Waldheimia septigera, Seguenza, Mem. della Soc. Italiana di Scienze Nat. vol. i. p. 44, pl. vi. figs. 1-10, 1865; Bull. Soc. Géol. France, 2nd ser. vol. xxv., 1868; Bull. Malacologica Italiana, anno iii., 1870.

Terebratula septata, Jeffreys, Proc. Zool. Soc. 1878, p. 407, pl. xxiii. fig. 1 (not Philippi); G. O. Sars, Bidrag til Kundskaben om Norges arktiske Fauna: Mollusca Regionis Arcticæ Norvegiæ, Christiania, pl. i. fig. 2, 1878 (not Philippi).

Terebratula floridana, Jeffreys (not Pourtales), Proc. Zool. Soc. p. 407, 1879.

Waldheimia septigera, Dall, Bull. Mus. Com. Zool. Harvard, vol. iii. p. 13, pl. 1. fig. 4, pl. 2. fig. 9, and Proc. of the California Acad. of Sciences, p. 182, 1872; H. Friele, "Bidrag til Vestlandets Mollusk-fauna, Vidensk.-Selsk. Forhand. p. 57, pl. 1. fig. 2, 1875, and The Development of the Skeleton of the genus *Waldheimia*, Archiv for Mathematik og Naturvidenskab, Bd. ii. p. 380, pl. iii. figs. 5, 6, and pl. iv. figs. 12-14, 1877; E. Deslongchamps, Études Critiques sur les Brachiopodes, p. 98, pl. i. figs. 17 & 18, pl. xv. figs. 4-6, and pl. xvi. figs. 1-3, 1884.

Shell somewhat square, ovate or subpentagonal, broadest anteriorly, more attenuated posteriorly, longer than wide, lateral edges rounded, wide and nearly straight in front, anterior margin sinuated, corners obtusely rounded anteriorly. Dorsal valve evenly convex to about half its length; anterior half divided into three lobes, the central one by far the largest and widest, forming a slightly raised, rounded, wide mesial fold. Ventral valve deeper than the dorsal one, with a wide concave sinus commencing at a short distance from the beak, and gradually widening until it reaches the front; beak incurved and truncated by a rather large circular foramen, completed and separated from the hinge-line by a deltidium in two pieces; lateral portions of the beak rounded. Surface of valves smooth, but marked by numerous fine concentric lines of growth. Colour light yellowish or greyish white. Shell-structure punctate. Dimensions variable, two large specimens measured:—

Length 1 inch 8 lines, breadth 1 inch 4 lines, depth 1 inch.

„ 1 „ 8 „ „ 1 „ 1 line, „ 10 lines.

In the interior of the dorsal valve, the cardinal process is so small that it can hardly be differentiated from the hinge-plate, and from under which a mesial septum extends to one half the length of the valve. Loop long, simple and reflected, the lamella of the reflected portion being wide, the principal stems nearly parallel.

Hab. Off Norway, Finmark, &c., in 100 to 300 fathoms. Atlantic coast of Spain. Dredged by the 'Porcupine' Expedition at various northern stations, at depths from 75 to 755 fathoms (Jeffreys). In British Seas between Shetland and Faeröe, by Jeffreys and J. Murray, in 345 to 570 fathoms. Vigo Bay (M'Andrew). Off Marocco and the Canary Islands ('Talisman' Expedition) in 331 to 861 fathoms.

Obs. The shell and some parts of the animal have been carefully examined. Mr. Dall says that its visceral area is very small, the muscular attachments being even smaller than in *Waldheimia floridana*. The stomach, produced into a point without differentiation of the intestine, is very much shorter than in *W. floridana*.

The development of the skeleton or loop in this species has been well studied and illustrated in two papers by Herman Friele in 1875 and 1877, and by E. Deslongchamps in 1884. Dr. Gwyn Jeffreys and myself have also followed out the same investigations. Herr Friele begins by informing us that he has not been able to examine the first stages in the development of the loop in this species, as he had done for that of *Waldheimia cranium*; that the youngest examples of *W. septigera* he has seen measured 1, 2, and 3 lines in length, and represented the second stage of *W. cranium*, in which the united lamellæ begin to split apart at the anterior end; and that the only essential difference at that age is in the form of the septum, being 4.5 millimetres and 5.5 millimetres respectively. The next stage he terms the *Megerlia* stage, the shell having attained 4 lines in length. In this stage in *Waldheimia cranium* the lateral walls were broken down by an aperture appearing in the middle of each and widening backwards. In *W. septigera* the break occurs, on the contrary, on the posterior end of the walls, and extends in a forward direction, the similarity at this stage to *Terebratella* being striking. The connection between the process of the lamellæ and the septum is severed in a specimen 6 lines in length, and in a specimen 8 lines in length the lamellæ are separated and the character of the loop is that of the adult *Waldheimia*. In Plate XI., figs. 7, 8, 9, 10 are from Friele's memoir, showing the modifications above described.

Waldheimia septigera was described for the first time by Lovén, in 1846; but having neglected to give a figure of his species, the shell was very little known until 1855, when I gave a figure of it in the Ann. & Mag. Nat. Hist. In 1863, in vol. ii. of his 'British Conchology,' and subsequently in a paper in the Proc. Zool. Soc., Dr. Gwyn Jeffreys erroneously maintained that Lovén's *Waldheimia septigera* was a synonym of Philippi's *Terebratula septata*, and this mistaken view he maintained to the last. In 1846, Signor Seguenza, after much trouble and skill, examined the perfect loop in several adult examples of Philippi's *T. septata*, and found it to be three times attached, as in *Terebratella*, while the loop in *W. septigera* is only twice attached, as in *Waldheimia*. The same indefatigable palæontologist also examined the loop of his Pliocene *Terebratula peloritana*, and found it to agree with that of *Waldheimia septigera*, of which he admits it to be a synonym.

My valued and distinguished friend Dr. Gwyn Jeffreys was again mistaken when he said, at p. 408 of his paper in the 'Proceedings of the Zoological Society' for 1878, that *Waldheimia peloritana* and *W. floridana* appear to be the same variety of *Terebratula septata*. Having had all the developed specimens in my hands, I am able to give my full support to Prof. Seguenza's statement.

Waldheimia septigera and *W. Raphaelis* are much more nearly related than either of them are to *W. floridana*. *Terebratella Mariæ* (Adams), from Japanese waters, is also said by Dr. Gwyn Jeffreys to be possibly a variety of *W. septigera* (his *T. septata*); but in the small specimen of *T. Mariæ* I have been able to examine the loop was three times attached, as in *Terebratella*.

Waldheimia septigera, under the name of *W. peloritana*, has been well described and illustrated by Signor Seguenza, in his Monograph of the Brachiopoda from the Tertiary deposits of Messina in Sicily, p. 49, pl. vi. figs. 1-10 (1865). It occurs plentifully in the Upper Pliocene rocks at Gravitelli.

32. WALDHEIMIA RAPHAELIS, Dall. (Plate XI. figs. 11-13.)

Waldheimia Raphaelis, Dall, Am. Journ. of Conch. vol. vi. p. 3, pl. vii. figs. *a, b, c, d*, 1870; Davidson, On Japanese recent Brachiopoda, Proc. Zool. Soc. 1871, p. 303, pl. xxxi. fig. 9.

Shell elongated, longer than wide, somewhat subtrigonal, widest about the middle and anteriorly, tapering posteriorly, nearly straight or gently curved outwardly in front. Dorsal valve moderately convex, with a very wide mesial fold, commencing at about a third of the length of the valve, uniformly and gently convex, or divided longitudinally by a small mesial groove. Ventral valve deeper and more convex than the dorsal one, with a broad deep mesial sinus commencing at a short distance from the extremity of the beak, and becoming deeper and wider as it reaches the front margin; on either side of the sinus the valve slopes abruptly to the lateral margins. Beak slightly incurved and truncated by a large circular foramen, separated from the hinge-line by deltidial plates. Surface smooth, marked only by concentric lines of growth; punctures oval-shaped outside, smaller, more distant, and circular on the interior surface of the valves. Colour light yellowish, with a slight rufous tinge. In the interior of the dorsal valve the cardinal process is small, quadrate, and stout; under it commences a small septum that does not reach to quite half the length of the valve. Loop long, reflected portion broad; crura much curved at their extremity. Length 1 inch 10 lines, breadth 1 inch 5 lines, depth 1 inch $1\frac{1}{2}$ lines.

Hab. Japanese coast, near Yeddo (Plumpelly and Dall). Sagami Bay, depth 100 to 200 fathoms (Döderlein).

Obs. I have seen five good examples of this remarkable species. In Mr. Dall's and Dr. Döderlein's specimens the fold in the dorsal valve is gently convex throughout; in a fine example I obtained from Mr. G. B. Sowerby the fold was probably exceptionally divided into two lobes by a median groove. The shell is named after Prof. Raphael Plumpelly, its nearest ally being the *Waldheimia septigera* of Lovén.

Waldheimia septigera, *W. Raphaelis*, and *W. floridana* belong, as was justly remarked by Mr. Dall, to a peculiar group in the genus *Waldheimia*, and are specifically distinct. The first is from northern European seas, the second from those of Japan, and the third from the Florida coast or reefs.

33. WALDHEIMIA FLORIDANA, Pourtales. (Plate XII. figs. 1-5.)

Waldheimia floridana, Pourtales, Bull. Mus. Comp. Zoöl. Harvard, vol. i. p. 127, 1868; Dall, Bull. Mus. Comp. Zoöl. Harvard, vol. iii. p. 12, pl. i. figs. 1-3, 1871.

Eudesia floridana, Bull. Mus. Comp. Zoöl. Harvard, vol. ix. p. 103, 1881.

Shell triangular, widest anteriorly, tapering posteriorly, generally as wide or wider than long; anterior margins strongly flexuous. Dorsal valve uniformly convex posteriorly, anteriorly divided into three lobes, the central by far the largest, forming a wide moderately elevated fold; lateral edges of valves flexuous, anterior corners rounded. Ventral valve deeper than the dorsal one, with a large wide mesial sinus commencing at a little distance from the extremity of the beak and gradually widening as it reaches the front; lateral lobes nearly flat or gently concave, sloping rapidly from the outer side of the fold and sinus to the edges of the shell. Beak incurved and truncated by a small circular foramen, very little separated from the hinge-line by a narrow deltidium in two pieces; beak-ridges rounded. Surface smooth. Colour greyish or brownish white with a slight touch of yellow. Length 11, width 12, depth 8 lines.

In the interior of the dorsal valve the cardinal process is small, the hinge-plate rather large, and from under its base a mesial septum of small elevation extends to about two thirds of the length of the valve. Loop simple, very broad and separate anteriorly, narrow posteriorly. Stomach spherical, with a long cylindrical intestine.

Hab. Very abundant, attached to Florida Reefs, depth 100 to 200 fathoms, on rocky bottom (Pourtales); off Sand Key, 125 fathoms (Sigsbee); off Havana, 175 fathoms (Dall).

Obs. Thanks to the great liberality of Prof. A. Agassiz, I have been able to examine a large number of specimens of this remarkable and well-characterized species from the dimensions of five up to eleven lines in length. When quite young and up to a somewhat advanced period of its growth, the shell is longer than wide, the fold and sinus being scarcely indicated in very young examples. Dall, in his excellent description of this species, states that it belongs to a peculiar group including *W. floridana*, *W. septigera*, and *W. Raphaelis*; he adds:—"Thus it is seen that the smallest species [*W. floridana*] is by far the widest and most inflated proportionately; the second species is the flattest in proportion to its length; and the third the most elongated."

"The greater portion of the mantle of *W. floridana* is of the most extreme tenuity and perfect transparency. It is furthermore so closely attached to the shell as to render its removal intact—even with the aid of acid—a matter of great difficulty. With this exception, the examination of its anatomy is easy. . . . The soft parts are mostly of a translucent whitish color. The number and disposition of the muscles are similar to those of *W. australis*, already described by various authors. The muscles themselves are of a glistening tendinous appearance, except at their points of attachment, where they are of a more or less dark yellowish-brown. The peduncle is moderately long, and the portion which is external or contained in the foramen is covered with a dark, horny, reddish-brown membrane or skin, and the attached extremity is trumpet-shaped. Upon opening the shell in its normal position, the median spires of the brachia are seen to be somewhat widely separated, and between them is stretched a fine translucent membrane extending

forward from the under lip of the mouth and following the downward curve of the median lobes. In this great extension of this membrane this species differs from *T. caput-serpentis* and *W. australis*, in which species the cirrhi of the median lobes touch at their extremities, and are separated by only a very narrow strip of membrane between their bases, so that the appearance is almost as if there was but a single broad band of cirrhi in the median line. This intervening membrane in ordinary specimens of *W. floridana* is about .24 of an inch in width at its narrowest visible portion. The upper and lower bands of cirrhi in the lateral loops are also much more widely separated by a similar membrane than in *W. australis*. The spiral portion in the middle lobe makes about two complete turns. The mouth is, as usual, just in front of the posterior junction of the brachial bands, and is in a rather long flexuous groove, the edges of which are of a dark brown colour, and somewhat thickened. The œsophagus is about half as long as the intestine, and has a slight curve, of which the convexity is anterior; it is transversely flattened close to the mouth, and is a little compressed laterally, behind that portion. It is of a nearly uniform calibre throughout. It has quite a thin lining membrane, which becomes thicker, though still smooth, in the stomach, and quite thick and longitudinally plaited in the intestine. The stomach is well differentiated from the alimentary canal and intestine, and is of an oval shape. It is embraced by the hepatic digitations, which are of a greenish-yellow color, and empty into the stomach by four ducts. The orifices of these ducts are of a compressed oval shape, obliquely inclined, and the anterior pair, which correspond to the right and left anterior congeries of hepatic digitations, are twice as large as the posterior pair, which similarly correspond to the anterior lobes or bunches of digitations. The intestine is twice as long as the œsophagus, of uniform calibre, and perfectly straight. It leaves the stomach abruptly without any dilatation of the portion adjacent to the latter organ, and reaches about half-way to the dorsal valve. The heart is situated behind the junction of the stomach and intestine. The termination of the intestine is abruptly rounded off and not at all pointed. It is entirely closed, and is upheld by the mesentery. It is also of a much darker color than the rest of the alimentary canal, being of a deep chestnut brown hue."

"The great pallial sinuses and their ramifications in *W. floridana* are of much less extent and disposed in quite a different manner from that which obtains in *W. australis*. The hæmal pallial system consists essentially of four branches which are remarkable for their straight course and the paucity of their ramifications. The neural pallial system is very similar, with a greater number of small sinuses about the perivisceral cavity, but in both lobes the narrowness and small extent of the sinuses, as compared with those of other species, is very marked, and the same is true with regard to the ovaries. But a very few exceedingly delicate spiculæ were observed in the floor of the greater sinuses. The heart consists of a very minute pyriform vesicle situated behind the intestine at its junction with the stomach, and sending one vessel in the hæmal direction along the median line of the stomach, and another on each side laterally. The ovaries are very limited in extent and principally confined to that portion of the sinuses which surrounds the visceral cavity, only their ultimate extremities entering

the larger branches of the great sinuses. The ova were visible in all stages of growth. Those floating free in the lacunes were nearly spherical, and of a flesh-colour; their substance seemed of a granular consistency, due perhaps to the action of the spirit in which they were preserved. The immature ova were pyriform, attached to the ovary by their pointed ends. No spermatophoræ or spermatozoa were observed in any of the specimens examined. The oviducts are situated as in *W. australis*. The lining-membrane of their trumpet-shaped portion was drawn into thin plicæ. Their apices were teat-shaped, with very small orifices. The setæ are very slender and fine, irregularly marked with transverse lines, but smoother towards their outer ends. They protrude from their follicles, hardly more than one third of their length. The mantle edge was brownish and seemed to have a slightly villous epithelium."

I have reproduced this long extract from Mr. Dall's admirable memoir, as I have done in similar cases from the works of other authors, because all that relates to the animal of different species of the same genus is of the utmost importance. The animals of species of the same genus no doubt agree in their general character, but often differ considerably in their respective details, and these differences should always, when possible, be carefully noted. The animal of every species should also under favourable circumstances be anatomically examined and described.

In his "Contributions to the Fauna of the Gulf Stream at great Depths" (Bull. Mus. Comp. Zool. vol. i. p. 127), Count L. F. de Pourtales informs us, in his description of *Waldheimia floridana*, that "Very young specimens are flatter, rounder, and have a straight margin; they could scarcely be distinguished from the young of *Terebratula cubensis*, if it was not for the loop and septum seen by transparency. There is also some variety of form in the old; in some specimens the length is greater than the breadth, and there is considerable diversity in the sinuosity of the frontal margin. It is always associated with *Terebratula cubensis*, the latter being still more common (in the proportion of about three to one), and making its first appearance in 100 fathoms."

34. WALDHEIMIA (MACANDREVIA) CRANIUM, Müller, sp. (Plate XII. figs. 11-23; Plate XIII. figs. 1, 2.)

Terebratula cranium, Müller, Zool. Danicæ Prodromus, p. 249, no. 3006, 1776.

Anomia vitrea, Chemnitz? (not Born), Conch.-Cab. vol. viii. p. 97, t. 78. figs. 707 & 709, 1795.

Terebratula cranium, Montagu, Trans. Linn. Soc. vol. xi. p. 188, tab. xiii. fig. 2, 1811; Turton, Conch. Dict. of British Isles, 1819.

Terebratula vitrea, Fleming (not Born), Phil. of Zool. vol. ii. p. 498, pl. iv. fig. 4, 1822; Schumacher (not Born), Essai d'un nouveau Système des Hab. des Vers Testacés, p. 133, pl. ix. fig. 1, 1817.

Terebratula euthyra, Philippi, Enum. Moll. Siciliæ, vol. ii. p. 68, tab. xviii. fig. 8, 1844.

Terebratula cranium, G. B. Sowerby, Thes. Conch. vol. i. p. 354, pl. lxx. figs. 60-62, 1846; Lovén, Index Moll. Scandinaviæ, Cefvers. K. Vet.-Akad. Förh. p. 183, 1846; Davidson, Monogr. Brit. Foss. Brach. Tertiary Species, vol. i. pl. i. fig. 8, 1852, and Ann. & Mag. Nat. Hist. 2nd ser. vol. xvi. pl. x. fig. 8, 1855.

Terebratula glabra, Leach, Brit. Moll. p. 359, pl. xiv. figs. 3, 4, 5, 1852.

Terebratula subvitrea, Leach (*teste* Reeve).

Waldheimia cranium, Barrett, Report Brit. Assoc. Glasgow, 1855, p. 107; Hancock, Phil. Trans. Roy. Soc. vol. cxlviii. 1858; Suess, Ueber die Wohnsitze der Brachiopoden, Sitzungsab. k. Akad. Wissensch. Wien, p. 200, 1859.

Macandrevia cranium, King, Proc. Dublin Univ. Zool. and Bot. Assoc. vol. i. p. 261, 1859.

Terebratula (*Waldheimia*) *cranium*, Reeve, Conch. Icon., Monogr. of *Terebratula*, pl. iii. fig. 6, 1860.

Terebratula cranium, Jeffreys, Brit. Conch. vol. ii. p. 11, and vol. v. p. 163, pl. xix. fig. 1, 1863.

Waldheimia euthyra, Seguenza, Pal. Mal. Class. Brach. p. 46, pl. v. figs. 6-14, 1865.

Waldheimia cranium, Dall, Am. Journ. of Conch. vol. vi. p. 110, 1870; Friele, Vidensk. Selsk. Forhandl. pl. i. fig. 1, 1875, and Archiv for Mathematik og Naturvidenskab, pl. i. figs. 1-4, pl. ii. figs. 5-7, pl. iii. figs. 1-4 and 7, 1877; G. O. Sars, Mollusca Region. Arct. Norveg. (Christiania) p. 10, tab. i. fig. 3, 1878; Jeffreys, Proc. Zool. Soc. 1878, p. 405.

Macandrevia cranium, Douvillé, Bull. Soc. Géol. de France, 3^e sér. vol. vii., 1879; K. A. Zittel, Handb. der Paläontologie, p. 703, 1880; E. Deslongchamps, Études critiques sur les Brachiopodes, pl. xiii. figs. 13-16, pl. xiv. figs. 4-9, and pl. xv. fig. 5, 1884.

Shell lenticular or ovate globose, longer than wide, broadest about the middle, lateral margins rounded, anteriorly subquadrate or rounded, sometimes much thickened at the margins. Dorsal valve evenly convex, without fold or sinus. Ventral valve deeper than the dorsal and evenly convex; beak slightly incurved, short, and truncated by an incomplete circular foramen, margined anteriorly by the umbo and by two rudimentary deltidial plates, which are in many specimens absent; beak-ridges not very sharply defined. Surface smooth, marked only by concentric lines of growth. Colour yellowish white or light horny. Shell-perforations separated by interspaces of about equal size. Dimensions variable—length 1 inch 1 line, width 10 lines, depth 7 lines. In the interior of the dorsal valve there is no defined cardinal process or mesial septum, but two deviating septa commence under the extremity of the umbo and extend to a little more than one fourth of the length of the valve; loop long, attached to the base of the hinge-plate, and extending to about four fifths of the length of the valve before becoming reflected*. In the interior of the ventral valve the teeth are supported by strong dental plates.

Hab. *W. cranium* is an abundant shell near the coast of Norway and in the northern seas. It was dredged on several occasions by R. MacAndrew and L. Barrett between Vigten Islands and the North Cape, in 25 to 160 fathoms, attached to stones; only abundant at Omnæsøe. Dr. Gwyn Jeffreys dredged it many times during the 'Lightning' and 'Porcupine' Expeditions off the Faerøe Islands, in 164 and 208 fathoms, and as deep as 690 fathoms during the 'Porcupine' cruise in 1869. MacAndrew obtained it outside Vigo Bay in 30 fathoms, Wallich in Greenland, and the Marquis de Folin and Dr.

* So delicate and brittle is the loop at the point where the primary lamellæ become reflected, that both Dr. J. Gwyn Jeffreys and Dr. W. B. Carpenter for some time entertained the mistaken opinion that at that point they were naturally disunited. This led Dr. Jeffreys to remark on p. 13 of the second volume of his 'British Conchology':—"Having carefully cleaned the inside of a specimen of *T. cranium*, containing the dried remains of the animal, with a weak solution of potash, and examined several other perfect shells of different ages, I could not perceive the least appearance of a loop, which is so evident in *T. australis*. The lamellar processes in the lower valve of *T. cranium* are equal in length, and end in sharp points. They may be compared to the chariot-blades used by the ancient Scythians, and they somewhat resemble the falciform apophyses of *Teredo* and *Pholas*. In the young of *T. cranium* these processes are extremely short. Their arrangement and shape are so dissimilar in species closely allied in other respects, that I should be inclined to consider their importance, as characters of generic distinction, somewhat doubtful." It was some considerable time before I could convince my distinguished friend that the loop of *Waldheimia cranium* differed but little from that of *W. flavesceus*.

P. Fischer off the south-west of France, in from 5 to 650 fathoms. It was also dredged by Prof. Sars, Friele, and others off the Norwegian coast; east coast of Shetland (Fleming and Jeffreys); North Hebrides by Dr. W. B. Carpenter and Sir Wyville Thomson, in 170-650 fathoms, &c.

Waldheimia cranium occurs fossil under the name of *Terebratula euthyra*, Philippi, in the Upper Pliocene rocks of Valle Lamato in Calabria, and near Catancaro near Reggio, at Gravitelli, Rometto, and other places near Messina in Sicily (Seguenza). Prof. Sars and others quote it from the glacial and post-glacial deposits of Norway. *Waldheimia cranium* has been recorded as from Halifax in Nova Scotia by the late Mr. Willis; but I have been informed by Mr. Dall, Mr. Whiteaves, and others, that what he took for the shell under description was a specimen of *Terebratella spitzbergensis*.

Obs. This species, its shell, animal, and different stages of development have been carefully investigated and admirably described by several competent zoologists. It has, however, been classed in the genera *Terebratula*, *Waldheimia*, and *Macandrevia*. In 1859, Prof. W. King considered that the species should be separated from *Waldheimia*, on account of the deviating septa and want of a mesial septum in the dorsal valve, as well as by the absence of dental lamina in the ventral one. I, however, with Mr. Dall, question whether the differences brought forward are of sufficient value to warrant us in admitting it as a separate genus; they are, at most, we think, sub-generic differences.

To Herman Friele we are indebted for a most valuable and important investigation into the modifications assumed by its loop, from the very youngest age up to the adult condition, which I now propose, in part, to transcribe from his memoir published in 1877. I also reproduce some of his figures on Plate XII., regretting that space will not permit me to give them all. He says:—"The earliest stage at which a coherent apophysary system of the *Waldheimia cranium* is observed, has a size of a little less than three millimetres [1 line], the hæmal valve being two millimetres [Plate XII. fig. 16]. Two long thin lamellæ project from the crura, connect with a filiform septum and run together in an acute angle in front, where they unite. By the connection of the lamellæ two close-set walls are given off, which by a reflection form a tube, the posterior end of which is closed. The size of the hæmal valve has reached 2·5 millim., but the state of things remains the same, save that both the loop (or the tube) and the lamellæ have expanded. [See Pl. XII. figs. 17, 17 *a*] *a* the lamellæ, *b* the two issuing vertical walls, or the lateral walls of the tube, *c* the closed tube (or the loop).

"The first visible change occurs *by an opening in the closed end of the tube* [figs. 18, 18 *a*]. The united lamellæ then begin to split apart at the anterior end. The hæmal valve has now attained the size of 4·5 millim. A continuous expansion of the loop-complex causes *a perforation in the lateral walls* [Pl. XII. fig. 19], and the septum becomes thinner and thinner. The next sizes observed were 5·2 millim. and 5·6 millim. (hæmal valve) [Pl. XII. fig. 21]. The same characteristics as those seen in [Pl. XII. fig. 19] are still in the main prevailing, but *the connection with the septum is broken off*, and there remains but a little lump of the latter at the bottom of the valve. The wedge-

formed separation of the lamellæ have progressed so far [Pl. XII. fig. 21], that only a short band of connection is left between them; the apertures in the walls have likewise widened, and a narrow slip is all that still combines the lower and the upper parts.

"The stage of growth illustrated by [Pl. XII. figs. 19-21], may properly be designated the *Megerlea Stage*, and it would be difficult to recognize in this state either the individual described in [Pl. XII. figs. 16 and 17], or the fully developed *Waldheimia*. In order to become a mature *Waldheimia* the band between the lamel-processes and the loop must vanish and the connected lamellæ must separate." In another figure, which we have not reproduced, "the *hinder connection between the upper and lower parts is completely severed*, but the lamellæ are still, though slightly, connected." In the next stage the connection is cut, Pl. XII. fig. 22, and in fig. 23 we have the most mature *Waldheimia cranium*. Here Friele gives figures from one fifth of a line in length up to one line, the smallest of which is no bigger than a dot.

Waldheimia cranium has sometimes been confounded with *Liothyris vitrea*, to which it occasionally bears some general external resemblance; but the two forms are not only specifically, but completely generically distinct. Mr. Lucas Barrett was able to examine the animal alive, and states that "the oral arms are so fixed to the calcareous skeleton as to be incapable of motion, except at their spiral terminations. . . . It has been supposed that these conjoined spiral ends can be unrolled like the proboscis of a butterfly; I never saw any disposition of the kind manifested. This species is more lively than *caput-serpentis*, moving often on its pedicle, and is more easily alarmed." The cirri are not protruded beyond the margins of the valve; when the shell is closed they are bent up; no currents were detected by Barrett, though frequently sought for. Dr. Gwyn Jeffreys states likewise (*op. cit.* vol. v. p. 163) that he has also frequently observed *W. cranium* turn round its peduncle, apparently in order to improve its position for the purpose of feeding, and that the valves on being touched close with a snap.

In Van Bemmelen's "Over den Bouw der Schelpen van Brachiopoden en Chitonen," 1882, and in his "Untersuchungen über den anatomischen und histologischen Bau der Brachiopoda Testicardinia," 1883, the author enters upon many anatomical and structural details with respect to *Waldheimia cranium*; he says that the number of cæca on the same part of the shell-structure in very old and young specimens was found to be the same; this fact, showing that the distance between two cæca (perforations) does not change with age, led him to the conclusion *that no intussusception occurs during the growth of the shell*. The bases of the calcareous prisms were found to be very regular-shaped at the margin of the shell (especially in *Terebratula* and *Terebratulina septentrionalis*); they became very irregular towards the older parts. The concentric lines of growth occurring on the outer surface were totally absent on the inner surface, which is explained by supposing the apposition at the margin to stop for some time, while the formation of new layers on the whole under surface continued. In *Waldheimia cranium* he found the sexes were separate.

Mr. Albany Hancock describes and figures the soft parts of the animal in his admirable memoir already so often referred to, and from which I now give a few extracts, with reproductions of his admirable figures of the dorsal and ventral views of *Waldheimia*

cranium (Pl. XIII. figs. 1, 2). He says that in *Waldheimia cranium*, and in *Terebratulina caput-serpentis*, "the dorsal adjustor muscles are not attached to a hinge-plate, as in *W. australis*, but have their insertions in the ventral valve itself, and are very large and powerful. In both species the superior extremities of these muscles are seen at the surface of the animal, on each side of the median line, elongated in the antero-posterior direction and extending between the occlusors almost as far forward as the anterior margins. . . . The divaricators and the accessory divaricators in *W. cranium* are likewise united in the same manner as in the species before alluded to." Mr. Hancock further observes that "the arms in *W. cranium* and *T. caput-serpentis* are disposed in the same manner as in *W. australis*; and in the former the calcareous loop is precisely similar to that of the latter; but in *T. caput-serpentis* it is very much reduced, the extended lateral portions having almost entirely disappeared, little more than the transverse portion existing; and this, together with the crural processes, which are united below across the median line, forms a collar upon which the bases of the arms rest.

"In *W. cranium* the intestine is very short, terminating in a blind sac before it reaches the ventral wall of the perivisceral chamber. It tapers gradually to a point, which is rounded, and suspended in its place by the mesentery. The mucous membrane, lining the intestinal tube of *W. cranium*, is exceedingly thick, and produced into five or six excessively stout, longitudinal folds, which in transverse sections exhibit a pyramidal contour, their apices almost meeting in the centre of the tube. . . . In *W. cranium* the genitalia are arranged as in *W. australis*, only the bands do not extend so far forward, and are of a pale yellow colour. The red matter is also present, sprinkling the surface with distant round spots. The bands are very finely granular, and in no instance have I detected eggs in them. It is therefore probable that those examined were out of season." Mr. Hancock also observes:—"There is no modification to note in the perivisceral chamber in any of the Terebratulidæ that I have had an opportunity of examining. The pallial sinuses, however, vary in several species. Thus in *W. cranium*, though there are still four such sinuses in each lobe, the trunks are proportionately smaller, and more nearly of a size; the branches are fewer and more attenuated, but, as in the other species, divided dichotomously twice or thrice, without any very marked symmetry."

The animal of *Waldheimia cranium* although differing in some unimportant details from that of *Waldheimia flavescens* is essentially similar. In both the loop is the same, and there is no difference, of any great importance, in the shape of the shell. I therefore question the necessity of burdening the nomenclature by placing *W. cranium* and *W. flavescens* in different and distinct genera. But as the contrary view has been expressed by such excellent observers as Prof. W. King, Douvillé, Zittel, Waagen, Deslongchamps, and others, I would not wish to press my own views too far, in opposition to theirs. M. Douvillé seems to think that *W. cranium* possesses a special development of its loop; while M. E. Deslongchamps, at page 141 of his *Études critiques sur les Brachiopodes*, 1884, differs from M. Douvillé, and expresses his opinion that the general appearance of the loop of *Macandrevia cranium* is actually the same as that attributed to *Waldheimia*. The shape of the principal stems shows the greatest

resemblance to those of *Zeilleria*; but the position of the larger-sized adductor muscles is no longer the same as in *Waldheimia*, while the complete absence of a mesial septum and the shape of the largely divided hinge-plate would, in his opinion (Deslongchamps'), be valid grounds for separating the genus *Macandrevia* from *Waldheimia*.

35. *WALDHEIMIA* (*MACANDREVIA*) *TENERA*, Jeffreys, sp. (Plate XII. figs. 6-10.)

Terebratula tenera, Jeffreys, Ann. & Mag. Nat. Hist. 4th ser. vol. xviii. p. 250, 1876, and Proc. Zool. Soc. 1878, p. 405, pl. xxii. fig. 7.

Shell rather small, extremely thin and delicate, oval, longer than wide, rounded laterally, less so in front; surface smooth, marked only by fine concentric lines of growth, and perforated by minute canals. In the interior of the dorsal valve no cardinal process is observable; hinge-plate comparatively large, and divided into two parts by a narrow mesial depression, between which, in the middle, a very narrow delicate longitudinal ridge or rudimentary septum is present, which extends to about one fifth of the length of the valve. The inner slopes of the hinge-plate extend likewise in the shape of two septa of small elevation to about the same distance, first diverging, then converging to the anterior extremity of the small median ridge. At the base of the hinge-plate the principal delicate stems of the loop are attached, and these extend to about two thirds of the length of the valve before they are reflected. Colour light brown. Length 6, breadth 4, depth 3 lines.

Hab. Dredged by Dr. Gwyn Jeffreys during the return voyage of H.M.S. 'Valorous' from Davis Strait, in lat. 56° 11' N., long. 37° 41' W., at a depth of 1450 fathoms, on a sea-bottom of *Globigerina*-ooze and stones.

Obs. I have seen two or three perfect examples of this extremely delicate species, brought home by Dr. Gwyn Jeffreys. The shell seemed full-grown. A specimen about one line in length showed traces of one of its early stages of growth, in the presence of a small vertical septum, at about half the length of the valve, to which the loop at that age had an attachment, as Friele has so well described to be the case in *Waldheimia cranium*.

Dr. Gwyn Jeffreys observes that "this species differs from *T. cranium* in being only half the size in exact measurement, and consequently one fourth in bulk; it is of a different shape, texture, and colour, compressed instead of convex, having a much shorter beak and smaller orifice, with not half the proportionate number of tubercles; and the blades are closer together, and do not extend so far towards the front. In the young of each species the comparative number of tubercles and prominence of the beak are distinctly marked; and the septum in the present species is shorter, although conspicuous and gnomon-shaped."

Uncertain Species.

36. *WALDHEIMIA* *WYVILLII*, Davidson. (Plate X. figs. 5, 6.)

Waldheimia Wyvilli, Davidson, Proc. Roy. Soc. vol. xxvii. p. 438, 1878; and (*W. wyvillii*) Report on the Brachiopoda, Voyage of H.M.S. 'Challenger,' Zool. vol. i. p. 44, pl. iii. figs. 13 a, b, 1880.

Shell ovate or longitudinally oval, very thin, semitransparent, light brownish yellow, smooth, marked at intervals by concentric lines of growth. Dorsal valve moderately

convex longitudinally, slightly flattened along the middle. Ventral valve deeper and more convex than the dorsal one, without sinus; beak incurved, truncated by an incomplete foramen, margined laterally by small deltidial plates. In the interior of the dorsal valve the loop is long and simple, the principal branches extending to a little beyond two thirds of their length before becoming reflected. Length 9, width 7, depth 5 lines.

Hab. Only one incomplete example was dredged by the 'Challenger' Expedition, off Valparaiso, at a depth of 2160 fathoms. *Terebratula Wyvillii* and *Discinisca atlantica* were obtained at the same time.

Obs. In external shape this species approaches *Waldheimia cranium*, which is, however, a thicker and more convex shell. The extremely delicate shell of *W. Wyvillii* is very remarkable, and reminds us of that of *W. tenera*, from which, however, it seems to differ in size and in some other particulars.

The fact that several of the species obtained at such depths as *Terebratula Wyvillii* and *Discinisca atlantica* possess such extremely thin and delicate glass-like shells, is certainly worthy of notice.

NOTE.—In accordance with the verbally expressed wishes of the late Thomas Davidson, LL.D., F.R.S., "The Davidson Collections of Recent and Fossil Brachiopoda" were presented to the nation, with the manuscripts and original drawings for many of his publications on the Brachiopoda, and the extensive series of works and pamphlets he had accumulated during nearly fifty years' investigation of this group of animals. The recent specimens are associated with the fossil species in the Geological Department of the Natural History Branch of the British Museum at South Kensington. The literature now forms part of the library of the same department in that Institution. [A. C.]

DESCRIPTION OF THE PLATES.

PLATE I.

- Figs. 1-6. *Liothyris vitrea*, Born, sp., at various stages of growth. Mediterranean Sea (Davidson Collection, Geological Department, British Museum). 6. Interior of the dorsal valve, enlarged. *cp*, cardinal process; *sl*, short loop; *dg*, diverging grooves for the attachment of the pallial sinuses.
- Fig. 7. Exterior of ventral valve of the same species, showing the four internal radiating furrows which serve for the attachment of the pallial sinuses, seen through the transparent shell.
- Fig. 8. Interior of the ventral valve after E. Deslongchamps. *cv*, visceral cavity; *ap*, raised portion of the mantle forming the walls of the visceral cavity; *svl*, furrow corresponding to the lateral venous sinus; *svm*, furrow corresponding to the median venous sinus.
- Fig. 9. Interior of the ventral valve showing the position of the muscular impressions. *a*, adductor or ocluser; *b*, ventral adjustors; *c*, divaricators.
- Fig. 10. Malformation of *Liothyris vitrea* (after E. Deslongchamps), seen in profile, showing a longitudinal frontal septum in both valves.
- Fig. 11. Full-grown spiculum, much magnified (after E. Deslongchamps), from a venous sinus near the raised part of the mantle of *Liothyris vitrea*.
- Fig. 12. Portion of the shell of *Liothyris vitrea*, magnified (after Van Bemmelen), to show the perforations and imbricated arrangement of the prisms.

- Figs. 13–13 *b*. *Liothyris vitrea*, var. *minor*, Philippi, from the Straits of Messina (Davidson Collection, Geol. Dept., British Museum).
- Figs. 14–16. *Liothyris vitrea*, var. *Davidsoni*, A. Adams, from Satanomosaki, Japan. 14. (Davidson Collection, Geol. Dept., British Museum.) 16. Interior of dorsal valve.
- Figs. 17, 18. *Liothyris arctica*, Fricke, sp. 17. Exterior, of natural size. 18. Interior of dorsal valve, enlarged. (Davidson Collection, Geol. Dept., British Museum.)
- Fig. 19. *Liothyris cernica*, Crosse, sp. : after Crosse's figure, Journ. de Conch. vol. xvi. pl. i. fig. 3. The specimen was obtained near the Island of Mauritius, from the stomach of a fish.
- Figs. 20, 21. *Liothyris Bartletti*, Dall, sp., from near Santa Cruz (Museum of Comp. Zool., Harvard).

PLATE II.

- Figs. 1–4. *Liothyris Moseleyi*, Davidson. 1–1 *b*. Exterior of the shell. 2. Interior of dorsal valve, to show the loop. 3. Interior of ventral valve, enlarged, showing the muscles; *a*, oclucor; *b*, ventral adjustor; *c*, divaricator; *m*, mantle; *n*, dorsal pallial nerves; *d*, dorsal pallial arteries. 4. Interior of dorsal valve, showing the labial appendages. 'Challenger' Expedition. West of Kerguelen Island. (Zoological Department, British Museum.)
- Figs. 5–7. *Liothyris uva*, Broderip, sp. 5–5 *b*. Type from Gulf of Tehuantepec (Zoological Department, British Museum). 6–6 *b*. A smaller specimen, from the same locality (Davidson Collection, Geol. Dept., British Museum). 7. Interior of dorsal valve, to show the loop.
- Figs. 8–14. *Liothyris Wyvillii*, Davidson, at various stages of growth. 12. Interior of dorsal valve, to show the loop. Dredged by 'Challenger' Expedition at different places named in the description of species. (British Museum and Davidson Collection.)
- Figs. 15, 16. *Liothyris subquadrata*, Jeffreys, sp. 15, 15 *a*, 15 *b*. Exterior of the shell after the type, in Davidson Collection, Geol. Dept., British Museum; 15 *c*. Shell-sculpture enlarged. From Setubal, coast of Portugal, near the mouth of the Tagus. 16. Interior of dorsal valve of the same species.
- Figs. 17, 18. *Liothyris sphenoidea*, Philippi, sp. : 'Travailleur' Expedition; off Morocco. (Muséum d'Histoire Naturelle, Paris.) 18. Interior of dorsal valve, to show the loop.
- Figs. 19–22. *Liothyris sphenoidea*, Philippi (= *cubensis*, Pourtales). 19, 19 *a*, 19 *b*, 21, 22. Florida reefs, dredged by M. de Pourtales (Davidson Collection, Geol. Dept., British Museum). 20. Ascension Island; dredged by 'Challenger' Expedition (Zoological Department, British Museum).
- Figs. 23–23 *b*. *Terebratulina* (or *Terebratulina*?) *Dalli*, Davidson. 23. Type, natural size: dredged by 'Challenger' Expedition, lat. 34° 37' N., long. 140° 32' E. 23 *a*, 23 *b*. Enlarged. The only known specimen is in the Zoological Department, British Museum.

PLATE III.

- Figs. 1–3 *a*. *Terebratulina Wyvillii*, Davidson. 1, 1 *a*, 1 *b*. Exterior of the type, dredged by 'Challenger' Expedition, N.W. of St. Thomas, West Indies, in 390 fathoms. 2. Interior of dorsal valve from same species (Zool. Dept., British Museum). 3. Smaller specimen dredged by the 'Travailleur' Expedition, from the western coast of Africa (Muséum d'Histoire Naturelle, Paris). 3 *a*. Shell-sculpture enlarged.
- Figs. 4–5 *a*. *Terebratulina Crossii*, Davidson. 4, 4 *a*. Type from Japanese water (Coll. of M. H. Crosse). 5, 5 *a*. A smaller specimen, same habitat (Davidson Collection, Geol. Dept., British Museum).
- Fig. 6. *Terebratulina Crossii*, Davidson. A large light salmon-coloured specimen obtained from Sagami Bay, Japan, by Dr. H. Döderlein, and from his Collection.
- Figs. 7–11 *a*. *Terebratulina japonica*, G. B. Sowerby, sp. A series of specimens from Japanese waters (British Museum and Davidson Collection). 11, 11 *a*, represent the *Terebratulina angusta*, Adams and Reeves, which is only a young *Terebratulina japonica*.
- Figs. 12, 12 *a*. *Terebratulina caput-serpentis*, Linné, sp., from Sagami Bay, Japan, found associated with *T. japonica*. In the Collection of Dr. Döderlein.

PLATE IV.

- Figs. 1-5. Various stages of growth of *Terebratulina caput-serpentis*, Linné, sp. 1, 2. Large specimens in the Zool. Dept., British Museum. 3, 4, 4a, 4b. Specimens of ordinary size from the coast of Scotland. 5. A very young specimen (after E. Deslongchamps).
- Fig. 6. A young specimen in which the crura of the loop are not yet united.
- Fig. 7. Interior of the dorsal valve of *Terebratulina caput-serpentis* (after E. Deslongchamps), showing the labial appendages and the cirri. *ac*, cardinal process; *vc*, visceral cavity; *b*, mouth; *bc*, descending or primary portions of the arms; *br*, ascending or reflected portions of the arms; *mi*, interbranchial membrane; *e*, shield; *sp*, median portion of the arms spirally coiled.
- Fig. 8. Dorsal view of *Terebratulina caput-serpentis* deprived of its shell (after Hancock). *a*, pallial lobe; *b*, body; *c*, peduncle; *d, d*, great pallial or genital sinuses; *e*, ramifications of ditto; *f*, muscular ties passing between the walls of ditto; *g*, genitalia seen through the walls of ditto; *h*, marginal fold; *i*, setæ; *j*, extreme pallial margin; *k*, depressions corresponding to the bases of the crura; *l*, ridge formed by the union of the pallial margins; *m*, edge of dorsal mesentery; *n*, liver seen through the walls of the perivisceral chamber; *o*, extremities of anterior oclusors; *p*, ditto of posterior oclusors; *q*, ditto of divaricators; *r*, ditto of dorsal adjustors.
- Fig. 9. Ventral view of the same (after Hancock); the letters up to *j* agree with those of fig. 8. *k*, extremities of oclusor muscles; *l*, ditto of divaricators; *m*, ditto of ventral adjustors; *n*, ditto of peduncular muscle; *o*, peduncular nerves.
- Fig. 10. Interior of the ventral valve of *T. caput-serpentis* (after E. Deslongchamps), showing the mantle in its entirety. *vc*, visceral cavity; *ap*, raised portion of the mantle forming the wall of the visceral cavity; *o*, organs of reproduction; *svl*, lateral venous sinuses; *svl**, trunks accruing from the division of the sinuses; *svm*, median venous sinus; *r*, small branches of the venous sinuses; on all the circumference one sees distinctly the festooned muscular border of the mantle, whereon are implanted a multitude of hairs (enlarged 5 diameters).
- Fig. 11. Interior of the dorsal valve of the same species (after E. Deslongchamps), showing the mantle also in its entirety. The same letters indicate the same parts on this valve, and the parts correspond with those of the ventral valve, with the exception of *ab*, which shows the brachial apparatus in the shape of a ring (enlarged 5 diameters).

PLATE V.

- Figs. 1-30. *Terebratulina septentrionalis*, Couthouy, after Morse, "Embryology of *Terebratulina*," Mem. Soc. Boston Nat. Hist. vol. ii. pl. viii., 1873.—1. A cluster of eggs from the genital band. 2, 3, 4. Eggs from the perivisceral cavity immediately after their escape from the pallial sinuses. 5. First ciliated stage. 6, 7, 8. Successive stages of transverse division of embryo, showing long tuft of cilia at cephalic extremity. 9, 10, 11. From a single embryo, showing various outlines assumed while swimming. 12, 13, 14. Different embryos showing first appearance of caudal or peduncular segment. 15, 16. Representing the same embryo contracted and expanded. 17, 18, 19. Embryos just attaching themselves by their caudal segment. 20. Embryo showing first appearance of ventral (?) area by the bulging of the thoracic ring. 21, 22. Embryos in various stages showing widening of thoracic ring, and its gradual growth towards enclosing the cephalic ring. 23, 24, 25. Successive stages of the embryo, showing formation of dorsal and ventral areas by the folding and growth of the thoracic ring. In fig. 26 the head is still seen projecting from the dorsal and ventral folds of the thoracic ring. 27 to 30. Succeeding stages of embryos drawn in various positions. The deciduous setæ appear in these stages.
- Fig. 31. One of the earliest stages of *Terebratulina septentrionalis*, after Morse (*op. cit.* pl. i. fig. 3), "in which the body has rapidly lengthened, and the peduncle is equal in length to the

remaining portion of the animal." *p*, peduncle; *pa*, point of attachment of peduncle; *cæ*, pallial cæca; *se*, setæ. ("Early Stages of *Terebratulina*," Mem. Soc. Boston Nat. Hist. vol. ii. pl. i. fig. 3.)

Fig. 32. A spiculum of *Terebratulina caput-serpentis*: highly magnified (after Hancock).

Fig. 33. Portion of the internal surface of the shell of *T. caput-serpentis*, showing the imbricated arrangement, and the internal orifices of the vertical perforations arranged in rows: magnified 100 diameters (after Carpenter).

Figs. 34, 34 *a*. *Terebratulina caput-serpentis*: very young, dredged by the 'Travailleur' Expedition. Coll. of the Marquis de Folin.

Figs. 35–37. *Terebratulina caput-serpentis*, var. *emarginata*, Risso, = var. *mediterranea*, Jeffreys; from the Mediterranean Sea; 37, was dredged alive between Pantellaria and Sciacca, Sicily. All in Davidson Collection, Geol. Dept., British Museum.

Figs. 38–40. *Terebratulina caput-serpentis*, var. *unguiculata*, Carpenter: 38, 39, exterior of shell; 40, interior of dorsal valve. Off Vancouver Island. (Davidson Collection, Geol. Dept., British Museum.)

Figs. 41–42 *a*. *Terebratulina Cailleti*, Cross e. 41–41 *c*. From Barbados (Davidson Collection, Geol. Dept., British Museum); 42, 42 *a*. Type, after Crosse's figure.

Figs. 43–52. *Terebratulina septentrionalis*, Couthouy, from off Halifax, Nova Scotia, and Lockport, Maine, at different ages (Davidson Collection, Geol. Dept., British Museum). 43, 44. Full-grown individuals. 47. A cluster of specimens attached to one another. 50, 51. Young examples. 48. Interior of dorsal valve and loop; 49. Dorsal valve, showing labial appendages. 52. Portion of shell, magnified.

Figs. 53–53 *b*. *Terebratulina Cumingi*, Davidson: type. From China seas.

Fig. 54. *Terebratulina abyssicola*, Adams and Reeve, sp.: after Reeve's figure. Off Cape of Good Hope. (Davidson Collection, Geol. Dept., British Museum.)

PLATE VI.

Figs. 1–8 *b*. *Terebratulina cancellata*, Koch, sp. 1–7. Specimens dredged by the 'Challenger' Expedition off Moncœur Island, Bass Strait, in from 38 to 40 fathoms; 7. Interior of the dorsal valve, enlarged, showing the labial appendages. 8–8 *b*. A small dwarfed form, dredged by Mr. J. Brazier at Inner North Head, Port Jackson, N.S.W. (Davidson Collection, Geol. Dept., British Museum.) Specimens 1–7 in Zool. Dept., British Museum.

Figs. 9–14. *Terebratulina radiata*, Reeve, sp. Different forms from the Cape of Good Hope (Davidson Collection, Geol. Dept., British Museum). 12. Specimen showing black colour-markings.

Figs. 15–17. *Terebratulina*? *Murrayi*, Davidson. 15. Outline of shell, natural size. 15 *a*, *b*, enlarged. 16. Dorsal valve, with loop magnified. 17. Dorsal valve, showing mantle, enlarged. Dredged by the 'Challenger' Expedition near Kermadec Island, south of the Fiji Islands, depth 600 fathoms. (Zool. Dept., British Museum.)

Figs. 18–20. *Terebratulina*? *tuberata*, Jeffreys, sp. From Josephine Bank, 340 to 430 fathoms (Collection of Dr. Gwyn Jeffreys, now in the Smithsonian Institution, Washington, U.S.A.): figures all much enlarged. 19. Interior of the dorsal valve.

Figs. 21, 22. *Terebratulina*? *trigona*, Jeffreys, sp. A young and immature shell, dredged by Mr. Saville Kent off the coast of Portugal, at a depth of 500 fathoms (Coll. of Dr. Gwyn Jeffreys, Smithsonian Institution, Washington, U.S.A.). 22. The shell opened, to show the loop.

Figs. 23–25 *a*. *Terebratulina*? *incerta*, Davidson. 23, 24 *a*. Dredged by the 'Challenger' Expedition in lat. 1° 47' N., long. 24° 26' W. (Zool. Dept., British Museum, and Davidson Collection). 25, 25 *a*. Interior of dorsal valve, with loop, after Dall, magnified 2½ diameters; dredged by the 'Blake' Expedition off Bequea. (Smithsonian Institution, Washington, U.S.A.)

PLATE VII.

- Fig. 1. *Terebratulina (Agulhasia) Davidsoni*, King: natural size. 1 *a*, 1 *b*, 1 *c*. Shell enlarged.
- Fig. 2. Ditto, interior of ventral valve, enlarged.
- Figs. 3, 4. Ditto, interior of dorsal valve and sectional diagram of same, enlarged.
- Fig. 5. Tubuli characteristic of *Terebratulina caput-serpentis* (1-5 after King). (Davidson Collection, Geol. Dept., British Museum.)
- Figs. 6-14. *Waldheimia flavescens*, Lamarck, sp. (= *Terebratula australis*, Quoy). Different specimens from Port Jackson, N.S.W., showing modifications in shape and in ribbing. 10, (after G. B. Sowerby), shows the extent the animal opens its valves. 13, represents the var. *recurva*, Quoy, from Shark Point, Port Jackson. 12, 12 *a*. A young smooth shell, from Bottle and Glass Rocks, Port Jackson. All the specimens, with the exception of fig. 10, from the Davidson Collection, Geol. Dept., British Museum.
- Fig. 15. Dorsal view of *Waldheimia flavescens*, the shell having been removed (after Hancock). *a*, pallial lobe; *b*, body; *c*, peduncle; *d*, great inner pallial sinuses, giving off branches to the margin; *e*, outer ditto; *f*, genitalia seen through the pallial membrane; *g*, red matter of ditto; *h*, marginal fold; *i*, setæ; *j*, circumpallial vessel; *k*, extreme pallial margin; *l*, median fissure, corresponding to longitudinal plate in valve; *m*, depression occasioned by the hinge-plate and the bases of the crura; *n*, orifices for the passage of the crura; *o*, ridge formed by the union of the pallial margins; *p*, extremities of anterior oclusors; *q*, ditto of posterior oclusors; *r*, ditto of divaricators; *s*, ditto of dorsal adjustors; *t*, liver seen through the wall of the body.
- Fig. 16. Ventral view of the same; the letters up to and including *k* correspond to those of fig. 15. *l*, extremities of the oclusor muscles; *m*, ditto of divaricators; *n*, ditto of accessory divaricators; *o*, ditto of ventral adjustors; *p*, ditto of peduncular muscles; *g*, cæcal extremity of intestine, seen through the wall of the perivisceral chamber; *r*, peduncular nerves.
- Fig. 17. Thin lamella of shell of *Waldheimia flavescens* (after Carpenter), showing the parallel disposition of the flattened prisms, and the regular arrangement of the passages which intervene between them; and at *a* the outcrop of these prisms of the internal surface of the shell. Magnified 100 diameters.
- Fig. 18. Internal surface of *Waldheimia flavescens* (after Carpenter), showing the imbricated arrangement of the extremities of the prisms, which are seen longitudinally at *a*. Magnified 100 diameters.
- Fig. 19. External surface of shell of *Waldheimia flavescens* (after Carpenter), showing the large trumpet-shaped ends of the vertical passages, covered in by opercular disks, which have radiating lines proceeding from them. Magnified 100 diameters.

NOTE.—Figures of the dorsal and ventral valves, of the muscular system, and of the labial appendages and animal of *W. flavescens* are given in the text at pp. 42, 45, 47, and 48 (woodcuts, figs. 3, 4, 5, 6, 7, & 8).

PLATE VIII.

- Fig. 1. *Waldheimia venosa*, Solander, sp. Type. (See Dixon, 'Voyage round the World,' 1789.) From Falkland Islands.
- Figs. 2-2 *c*. *Waldheimia venosa*. The largest specimen known, dredged alive in Stanley Harbour, Falkland Islands, by Admiral B. J. Sullivan in 5 or 6 fathoms, 1844.
- Fig. 3. *Waldheimia venosa*. Type of the *Terebratula globosa* of Lamarck, which was badly figured in the Encyclopédie Méthodique, pl. 239. fig. 2, 1789. (Muséum d'Histoire Naturelle, Paris.)
- Fig. 4. Ditto. Type of the *Terebratula physema*, Valenciennes, after Reeve's figure, 'Monogr. of *Terebratula*, Conch. Icon.' pl. vi. fig. 23. From Coquimbo. (Muséum d'Histoire Naturelle, Paris.)
- Fig. 5. Ditto. Interior of the dorsal valve of a full-grown specimen, to show the loop.
- Fig. 6. Ditto. Type of the *Terebratula Fontaineana*, A. d'Orbigny, 'Voyage dans l'Amérique Méridionale,' vol. v. pl. 85. figs. 30 & 31, 1847. (Reduced $\frac{2}{3}$ natural size.) From Coquimbo.

PLATE IX.

- Figs. 1, 1 *a*. *Waldheimia venosa*, var. *dilatata*, Lamarck. Type of Lamarck's *Terebratula dilatata*, from a drawing made in Paris from the type in the Muséum d'Histoire Naturelle, Paris.
- Figs. 2-13. *Waldheimia lenticularis*, Deshayes, sp. A series of specimens of different forms and ages from the Straits of Foveau, New Zealand, in 15 fathoms water (Davidson Collection, Geol. Dept., British Museum). 2-3 *b*. Large specimens. 10. Interior of dorsal valve, showing loop in the adult condition. 11-13 *a*. Modifications assumed by the loop previous to its attaining the adult state.

PLATE X.

- Figs. 1-4. *Waldheimia Grayi*, Davidson. 1, 1 *a*, 2. Typical form. 3. Interior of dorsal valve, to show the loop. 4, 4 *a*, 4 *b*. Variety *transversa*. All dredged from Japanese waters except the specimen shown in fig. 2, which was dredged by Sir E. Belcher in the Straits of Corea. (Davidson Collection, Geol. Dept., British Museum.)
- Figs. 5, 6. *Waldheimia Wyvillii*, Davidson. 5, 5 *a*. Exterior of shell. 6. Interior of dorsal valve, enlarged, showing the loop. Dredged by 'Challenger' Expedition, lat. 33° 31' S., long. 74° 43' W., depth 2160 fathoms. (Zool. Dept., British Museum.)
- Figs. 7-17. *Waldheimia kerguelensis*, Davidson. 7-15. A series of forms of different ages. 16. Interior of dorsal valve, showing the loop. 17. Interior of dorsal valve, showing the labial appendages. All dredged by the 'Challenger' Expedition south of Kerguelen Islands, also off Marion Islands. Found in depths varying from 100 to 150 fathoms. All the specimens figured are in the Zool. Dept., British Museum.

PLATE XI.

- Figs. 1-10. *Waldheimia septigera*, Lovén, sp. 1-1 *c*. From Faeröe Sound, 570 fathoms (Davidson Collection, Geol. Dept., British Museum). 2-2 *b*. A very large broad specimen, after Sars. 3. A young example, from the British sea, between Shetland and Faeröe ('Porcupine' Expedition), depth 345 fathoms. 4. Interior of dorsal valve with its fully developed loop. 5. Profile view of the same (after Sars). 6. View of the loop seen from the front, much enlarged. 7-10. Different stages in the development of the loop, after H. Friele, figures magnified; all from Norwegian waters.
- Figs. 11-13. *Waldheimia Raphaelis*, Dall. 11-11 *c*. A very perfect example from off the coast near Jeddo, Japan (Davidson Collection, Geol. Dept., British Museum). 12, 12 *a*. A large typical example from Sagami Bay, coll. Dr. Döderlein. 13. Interior of dorsal valve, showing its fully developed loop. 12-13. In the Museum of the Smithsonian Institution, Washington, U.S.A.

PLATE XII.

- Figs. 1-5. *Waldheimia floridana*, Pourtales, at different ages, dredged by Agassiz and de Pourtales off Florida reefs, between 100 and 200 fathoms (Davidson Collection, Geol. Dept., British Museum). 5. Interior of dorsal valve, to show the loop.
- Figs. 6-10. *Waldheimia* (or *Macandrevia*) *tenera*, Jeffreys, sp. 6-6 *b*. Specimen of the natural size. 7. Interior of ventral valve, enlarged. 8. Interior of dorsal valve, to show the loop. 9. Posterior portion of the interior of the dorsal valve very much enlarged, to show the hinge-plate and septa. 10. A very young example in which a small pillar-shaped ventral septum is seen in the interior of the dorsal valve. These figures, drawn by Davidson, were published by Jeffreys in the Proc. Zool. Soc. for 1878. Dredged by Dr. Gwyn Jeffreys, lat. 56° 11' N., long. 37° 41' W., 'Valorous' Expedition, at a depth of 1450 fath. (Jeffreys Collection, Smithsonian Institution, Washington, U.S.A.)

Figs. 11–23. *Waldheimia* (or *Macandrevia*) *cranium*, Müller, sp. 11–15 *b*. A series of different ages. 11. A very large example (after Sars). 15–15 *b*. A very minute or first stage (after Fricke). 16. The first stage of the development of the loops with a coherent apophysis (Fricke). 17, 17 *a*. Specimen a little further advanced: *a*, lamellæ; *b*, vertical walls; *c*, tube (Fricke). 18, 18 *a*. Opening of the posterior end of the tube (Fricke). 19. Perforation of the lateral walls (Fricke). 20, 21. Connection with the septum is broken (Fricke). 22. The lamel-processes separated. 23. Adult and fully developed loop. All the specimens from Norwegian waters. (See Fricke, Develop. Skelet. in *Waldheimia*, Archiv Math. og Naturvidenskab, ii. p. 380, pls. i. & ii., 1877.)

PLATE XIII.

- Fig. 1. *Waldheimia* (or *Macandrevia*) *cranium*, Müller, sp. Dorsal view of the animal, much enlarged (after Hancock). *a*, pallial lobe; *b*, body; *c*, peduncle; *d*, great inner pallial sinus; *e*, outer ditto; *f*, genitalia appearing through the walls of the sinuses; *g*, marginal fold; *h*, setæ; *i*, circumpallial vessel; *j*, extreme pallial margin; *k*, two lobes corresponding to cavities at the sides of the foramen; *l*, similar lobes corresponding to the cavities in the umbonal region of dorsal valve; *m*, ridge resulting from the union of the pallial margins; *n*, extremity of anterior oclusors; *o*, ditto of posterior oclusors; *p*, ditto of dorsal adjustors; *q*, ditto of divaricators.
- Fig. 2. Ventral view of the same (after Hancock). The letters up to *k* as in fig. 1:—*l*, extremities of the oclusor muscles; *m*, ditto of divaricators; *m**, portions of ditto, corresponding to the accessory divaricators; *n*, ditto of ventral adjustors; *o*, ditto of peduncular muscle; *p*, peduncular nerves.
- Figs. 3–9 *a*. *Terebratella coreanica*, Adams and Reeve, sp. 3. Type specimen figured by Reeve. 4–4 *c*. A fine and large specimen. 5, 5 *a*. A still larger example (*Terebratula miniata*, Gould), dredged by L. V. Schrenk, from Hakodadi, Japan. 6, 6 *a*. An elongate globose variety, dredged by Capt. St. John in Tsuga Strait. 7. Interior of dorsal valve, to show loop. 8, 8 *a*. Elongate variety (*T. Bouchardi*, Davidson). 9, 9 *a*. Var. *Bouchardi*, Davidson. Specimens 4, 5, 6, 7, and 8, in the Davidson Collection, Geol. Dept., British Museum; 9, in the Zool. Dept., British Museum.

TEREBRATULINÆ

1-12. *LIOTHYRIS VITREA*. 13. *L. VITREA* var. *MINOR*. 14-16. *L. VITREA* var. *DAVIDSONI*.
17-18. *L. ARCTICA*. 19. *L. CERNICA*. 20-21. *L. BARTLETTI*.

TEREBRATULINÆ.

Van Iterson lith.

1-4. *LIOTHYRIS MOSELEYI*. 5-7. *L. UVA*. 8-14. *L. WYVILLII*.
15-16. *L. SUBQUADRATA*. 17-22. *L. SPHENOIDEA*. 23-23^b. *L. (or TEREBRATULINA) DALLI*.

West, Newman. imp.

TEREBRATULINÆ.

TEREBRATULINÆ.

TEREBRATULINÆ.

1-31^a & 43-52. TEREBRATULINA SEPTENTRIONALIS.32-34^a T. CAPUT-SERPENTIS. 35-37^a T. CAP-SERP. var. MEDITERRANEA. 38-40. T. CAP-SERP. var. UNGUICULATA.

Hammond lith.

41-42^a T. CAILLETI. 53-53^b T. CUMINGI. 54. T. ABYSSICOLA.

Hanhart imp.

TEREBRATULINÆ.

1-8^b. TEREBRATULINA CANCELLATA. 9-14. T. RADIATA. 15-17. T. MURRAYI.
18-20. T. TUBERATA. 21-22. T. TRIGONA. 23-25^a. T(?) INCERTA.

TEREBRATULINÆ.

TEREBRATULINÆ.

TEREBRATULINÆ.

Van Herson lith.

1. WALDHEIMIA VENOSA var. DILATATA. 2-13^a W. LENTICULARIS.

West, Newman & Co imp.

TEREBRATULINÆ.

Van Iterson lith.

1-3. WALDHEIMIA GRAYI. 4. W. GRAYI var. TRANSVERSA.
5-6. W. WYVILLII. 7-17. W. KERGUELENENSIS.

West, Newman imp.

TEREBRATULINÆ.

TEREBRATULINÆ.

TEREBRATULINÆ.

TEREBRATELLINÆ.

1-2. WALDHEIMIA (MACANDREVIA) CRANIUM.

3-9. TEREBRATELLA COREANICA. (5. = T. MINIATA Gould. 8-9. = T. BOUCHARDI, Dav.)