PROCEEDINGS OF THE SOCIETY.

MEETING OF 14TH JUNE, 1882, AT KING'S COLLEGE, STRAND, W.C.,
THE PRESIDENT (PROFESSOR P. MARTIN DUNCAN, F.R.S.) IN
THE CHAIR.

The Minutes of the Meeting of 10th May last were read and confirmed, and were signed by the President.

The List of Donations (exclusive of exchanges and reprints) received since the last meeting was submitted, and the thanks of the Society given to the donors.

	From
Cole, A. C.—Studies in Microscopical Science, Vol. i. Nos. 1 and 2, and two preparations in illustration	<i>The Editor.</i>
Geological Magazine, Vols. i.-xviii. (1864-81)	<i>Mr. Crisp.</i>
Heurck, Dr. H. Van.—Synopsis des Diatomées de Belgique, Fasc. V. Crypto-Raphidées, 1 ^{re} Partie, pls. 78-103	<i>The Author.</i>
Hitchcock, R.—Synopsis of the Fresh-water Rhizopods. (8vo, New York, 1881) 56 pp., with 4 pls. now added	<i>The Author.</i>
Jones, T. Rupert.—Catalogue of the Fossil Foraminifera in the Collection of the British Museum, pp. xxiv. and 100. (8vo, London, 1882)	<i>The Trustees.</i>
Micrographic Dictionary. 4th ed. Part 12	<i>Mr. Van Voorst.</i>

The President called special attention to the donation of a complete set of the 'Geological Magazine,' and Mr. Stewart referred to the 'Studies in Microscopical Science,' edited by Mr. A. C. Cole, the illustrations of which he considered were very good, and the letterpress appeared to be equally so. The novel feature was that a microscopical preparation accompanied each part.

Mr. Crisp exhibited a $\frac{1}{4}$ -inch objective by Tolles, with a very tapering front, and stated that it was claimed on Mr. Tolles' behalf that he was the first to make such fronts about ten years ago.

Mr. Ingpen said that he had one of Andrew Ross's $\frac{1}{2}$ -inch objectives with a triplet front, dated 1848, which was similarly tapered.

Mr. Beck said that he had made them in the same way for the last fifteen years.

The President, referring to J. L. de Lanessan's 'Traité de Zoologie—Protozoaires' (8vo, Paris, 1882, pp. vii. and 336, and 281 figs.), said that it treated of *Amœbæ* on a somewhat grand scale, particularly as regarded the larger kinds, and he would suggest to some of the Fellows that they should search for these species as being of great interest. He obtained one from the Hampstead ponds which he found full of minute refractile points.

Mr. Stewart said he had examined some large *Amœbæ*, a short time ago, which came from Mr. Ingpen's aquarium, and were crowded with refractile points. They were distinctly crystalline in character, and had a vesicular nucleus.

Mr. Hartog said that he used to find the nuclei in the large *Amœbæ* with a very distinct network, although one which he found had a hollow nucleus.

The President inquired if Mr. Hartog had ever been able to iodize *Amœbæ*?

Mr. Hartog said he had stained them with picro-carmin, but had not done so with iodine.

Dr. Ralph, the President of the Victoria (Australia) Microscopical Society, responding to a welcome from the President and the Meeting, said he should like to take the opportunity of bringing before the Fellows of the Society the examination of the leaves of various kinds of plants by means of the action of prussic acid and ammonia. The process was a very simple one, it being only necessary to place the section under the Microscope, and then introduce the compound. He had only had time as yet to examine a few kinds of leaves in this way, but the vine had hitherto given the most marked instances of the chemical action to which he referred. A longitudinal section as thin as possible being made of the leaf and placed under the Microscope, the fluid should be added, and would be found to penetrate the structure, chiefly acting on the ducts, which were within reach of the bark. In a few moments the most extraordinary colours made their appearance, such as claret, amber, port-wine colour, and others having all the appearance of a coloured injection of the tissue, only that in about a quarter of an hour it all disappeared. Other leaves which he had tried behaved much in the same way, though they did not all respond to the reagent in an equal degree. It was only in the brittle sappy stems that it was possible to get the best results. In consequence of these observations, he had been able to treat sections of the human subject with prussic acid with marked effects. Whenever he got in the plasma amorphous particles distinctly blue in colour, he also found the formation of amyloid bodies in the blood. These bodies were very well defined, and under favourable circumstances, in polarized light, the black cross could be seen. The production of amyloid forms by chemical means, not only under the action of hydrocyanic acid, but by chloral, formic acid, or solution of copper in ammonia, was a point of considerable interest. If a portion of either of these reagents were added to fresh blood, and examined carefully under the Microscope, they would, in all probability, find these starch-like bodies developed in the field. It was not easy to show the process in a room to a number of persons; but if any one present was interested in these subjects, he should be very glad to demonstrate what he had been describing.

Mr. Stewart inquired whether these bodies were supposed to be formed by the reagent, or were they supposed to be really present before, but only to be made visible by the reaction?

Dr. Ralph said that under favourable circumstances they might see a globule which they would be disposed to say was oil; when the reagent was added it would increase in size from about the $\frac{1}{3000}$ inch to about the $\frac{1}{1000}$ inch; later it would suddenly become

opaque, and then it would colour in such a way that any one looking at it would say at once that it was starch. So far as he had carried out the observations, it seemed as if these starch-grains really did develope, and he thought it might be an instance of the commencement of the process of amyloid deposit.

Dr. Morris, of Sydney, was introduced to the Meeting by the President, and detailed some experiments which he had made in mounting diatoms in phosphorus in such a way as not to be inflammable. As he had only just arrived from Sydney, he had not had time to write anything on the subject.* He had seen some of the specimens which had been mounted in England, and was under the impression that the solution was too strong. He proposed, therefore, to reduce it to such a strength that if a piece of white blotting-paper was put into it, it would not blaze up. By such a solution all difficulty as to using the medium was done away with, the only necessary precaution being to have a basin of water near at hand to dip the fingers into.

Mr. Stephenson said that if they used a weak solution they would get a lower refractive index, whereas the principal object in mounting in phosphorus was to get as great a difference as possible between the refractive index of the medium and that of the object, for on this difference alone the increase of visibility depended.

Mr. Crisp said that if phosphorus was used diluted to a refractive index of 1.7, the visibility of the diatom would be proportionately reduced, and as there was no virtue in phosphorus, except for its high refractive index, it would be better not to use it at all in such a condition, but to take some non-inflammable substance which had the lower refractive index.

Dr. Morris said that, with regard to the value of the process, he might mention that he had tried some *Naviculæ*, and that with an oil-immersion lens the object mounted in phosphorus was resolved in the most perfect manner—far superior to anything that could be seen with balsam.

Mr. Crisp said that it must be borne in mind that the danger incidental to the use of phosphorus was not confined to the process of mounting. A case recently occurred in which an object-glass, brought down too hard, broke the cover-glass, and the observer, having wiped off the exuding phosphorus with his handkerchief, put it into his pocket and set himself on fire.

Mr. Ingpen exhibited and described some high-power achromatic eye-pieces, by Steinheil, of Munich (Series A F). They were constructed for astronomical purposes, but he thought they would prove serviceable for use with the Microscope. Those exhibited were of $\frac{3}{4}$, $\frac{1}{2}$, and $\frac{1}{3}$ inch focus, each having a field of 40° (see p. 551). Mr. Ingpen said that, whatever differences of opinion there might be as to

* See note by Dr. Morris written after the meeting, *supra*, p. 579.

the value of deep eye-pieces, there was probably none as regarded the desirability of securing the best possible definition, whatever amplification was used. He considered that with Huyghenian eye-pieces of less than an inch focus there was a serious deterioration of the image. The eye-pieces now exhibited gave the sharpest definition of any of similar foci he had hitherto been able to examine.

Mr. Crisp explained the new method devised by Dr. Ehrlich, the assistant of Dr. Koch, for preparing the bacteria of tuberculosis, which constituted a considerable improvement upon the original process. It produces a more intense colour in the bacteria, so that they appear larger, and can be recognized with a lower power, even, it is said, of 250 diameters (see p. 572).

Prof. Abbe's process of increasing the consistency of pure cedar-oil, so as to render it less fluid and therefore more convenient for use as a homogeneous-immersion fluid, was explained by Mr. Crisp. The oil is spread out in thin layers, and exposed to the action of the air and light for a long time, until it becomes of the consistency of castor-oil (see p. 551).

Mr. W. F. Petterd's letter was read accompanying a collection of diatoms from Tasmania.

Mr. A. Certes' letter, reiterating his claim to the discovery of a method of staining the nucleus of *living* Infusoria, was read by Mr. Crisp, who said that the author of the interpolation by which such a claim was described as "erroneous," now agreed that he was mistaken in the view he had expressed, not having in fact sufficiently noticed that the claim related essentially to the nucleus of the *living* animals (see p. 576).

Mr. Crisp called attention to a process devised by E. Korschelt, of Freiburg, for preserving Infusoria and *Amæbæ* by the use principally of osmic acid and chromic acid, no method having hitherto been found for the latter organisms. Dr. A. Gruber had also succeeded in preserving Heliozoa by the method, in excellent condition (see p. 574).

Mr. R. Miller's description of his caoutchouc cement for making and sealing cells was read, in which he stated that he did not claim to have discovered any new material, but to have accidentally hit upon a new method of combination. The material so obtained was singularly easy to work, and reliable in its results (see p. 579).

Dr. Van Heurck's letter was read describing favourably his further experiences with the Swan electric lamps and Faure-Reynier accumulators (see p. 557).

Professor Abbe's note was read with reference to the description of his camera lucida, *ante*, p. 261, in which he stated that the benefit of the arrangement consisted simply in the ease with which it may be used. In other respects it did not possess any advantage over those forms which fulfil the condition that the image is seen without reflection or other loss of light.

Mr. Ahrens exhibited and explained the construction of his new erecting binocular Microscope. In general outline it presented much the same appearance as that of Mr. Stephenson, the tubes being inclined, whilst the stage was horizontal. The erection of the image was, however, obtained by introducing a Nachet erecting prism over the objective, another prism similar to the Wenham being used above to divide the pencils.

Dr. Millar said that, whilst he could speak with great praise of the advantages of the erecting binocular as an instrument to work with, leaving nothing to be desired for comfort and convenience, he thought the one now before them did not appear to give an entirely satisfactory stereoscopic effect.

Mr. Michael said there was no doubt that a binocular Microscope, with a flat stage and an erecting arrangement like that of Mr. Stephenson, was an immense help to those who worked at living objects. In drawing from living insects he had found it of the greatest practical service.

Professor Liversidge's communication on the "silver-fish" of Sydney was read, in which he defended the accuracy of the observations of Hooke recorded in his 'Micrographia' (1665) as against the strictures of Mr. W. Blades in his 'Enemies of Books' (3rd edition, 1881). Specimens of the insects were sent, and labels and sheets of paper destroyed by them exhibited (see p. 500).

Mr. Hartog said there was no doubt as to the animal being *Lepisma saccharina*, the same that was often found in this country. It was mentioned by Emerson Tennant as being thought by some to prey upon books, but the author was of opinion that it rather preyed upon the insects which were found in the books, a view which he (Mr. Hartog) considered an error.

Mr. Thom's letter on *Saccharomyces* was read, in which he mentioned that he had devoted all his spare time for ten years in tracking these organisms, and now made all his bread without any yeast, so called.

Mr. Bennett, to whom the letter was addressed, made some remarks upon the subject.

Mr. Badcock's note was read (in his absence from illness) as to his observation of eyes in the adult forms of *Melicerta ringens*, *M. tyro* (or *tubicularia*), and *Stephanoceros Eichhornii*, also as to the development of the latter form from the egg. He also pointed out that

the tube with advancing age got filled up with mucilage, until only just room enough was left for the creature to move up and down (see pp. 345-6 and p. 512).

Mr. Badcock's note was read as to a criticism by Professor O. Bütschli of his paper on *Acinetina* (this Journal, III. (1880) p. 561), which had appeared in the 'Zoologischer Jahresbericht' for 1880 (p. 173). Professor Bütschli, in remarking on Mr. Badcock's comparison of the finely ciliated newly-born *Podophrya quadripartita* to *Megatricha partita*, referred to the latter as a Rotifer (!), being apparently surprised that such a comparison should have been made. At first Mr. Badcock was puzzled to know how the reporter could have fallen into such a mistake, but it had since occurred to him that the explanation was to be found in the fact that there were Rotifers named *Megalotricha*, and although these were very large Rotifers, while *Megatricha* is one of the most delicate of all the Ciliata, it was most probable that this was the origin of the mistake.

Mr. Wilson thought it was strange that Professor Bütschli, who was a very eminent authority on the Protozoa, should not have been familiar with *Megatricha*, as appeared to be the case. Many of the reviews in the 'Zoologischer Jahresbericht' were, however, obliged to be hastily done, and some excuse must be made on that account, as well as for the fact that the reporters had to deal largely with papers written in foreign languages.

Professor Abbe's paper "On the Relation of Aperture to Power," Part II., was laid before the meeting by Mr. Crisp, who said that as it was a very long communication, he had prepared a *résumé*, which presented the leading points in a condensed form suitable for being read to the Meeting. This he was proceeding to do, when

Mr. Beck, interposing, said he thought that as the paper was an important one, and required a good deal of consideration, a *résumé* of it would be of very little use to them. He would suggest, therefore, that the paper should be printed, and that when it had been before them *in extenso*, an evening should be specially devoted to its consideration and discussion.

The Chairman (Mr. Glaisher, in the absence of the President), on the contrary, thought that it would be very useful to have an abstract of the paper read.

Mr. J. Mayall, jun., also thought it very desirable that the Society should be in possession of Professor Abbe's views, without having to wait until the next issue of the Journal.

The Meeting having indorsed this view, Mr. Crisp read a *résumé* of the paper (see p. 460).

The Chairman said that the paper was obviously one of very considerable importance, and he had great pleasure in proposing that the warm thanks of the Society should be presented to Professor Abbe for it.

Mr. Stephenson seconded the suggestion, and said that, having

had the advantage of reading the paper in MS., he was able to say that it contained a most exhaustive discussion of the subject.

Mr. Crisp said that it was the first attempt that had been made to establish a definite relation between aperture and power, and amongst other benefits it could not fail to prevent the perpetration in future of such absurdities as had been issued in the way of low-power objectives with very large aperture.

The Chairman said that the report of the committee appointed to consider the question of standard gauges for eye-pieces and substages had been adopted by the Council. The gauges recommended for eye-pieces were 1.35 inch and 0.92 inch, and for substages 1.5 inch.

The following is the report of the committee :—

“The committee who were appointed under a resolution of the Council directing them to ‘consider the advisability of establishing standard gauges for eye-pieces and substages,’ after communication with the leading opticians and with Fellows of the Society, now present their report as follows :—

“The committee have found that a very general desire exists on the part of workers with the Microscope that there should be *standard gauges* both for eye-pieces and substages, and the committee therefore resolved ‘That in the opinion of this committee it is expedient that standard gauges should be established for the eye-pieces and substages of Microscopes.’

“In considering the question of the *number* of standards to be adopted, the committee were of opinion that it would not be practicable to establish a single uniform gauge for eye-pieces. On the one hand, a considerable number of Microscopes issued are of small size, and could not with convenience be fitted with a body-tube of large diameter; while, on the other hand, it would be undesirable to reduce the diameter in use with first-class instruments. The same considerations do not apply to the case of substages, which are frequently of as great a diameter in the smaller instruments as in the larger. The committee, therefore, resolved to recommend ‘That the standards for eye-pieces should be two in number, with a single standard for substages.’

“With regard to the *size* of the standards, the committee did not feel themselves able to treat the question as one entirely open, but considered that it would be preferable to select some sizes already in general use, so as to involve the minimum of change. They therefore resolved to recommend ‘That the two standard gauges for eye-pieces should be, for the No. 1, 1.35 inch, and for the No. 2, .92 inch (external diameter), and that the gauge for the substages should be 1.5 inch (internal diameter).’ The No. 1 gauge is generally used for the larger instruments in England, whilst No. 2 is that adopted by many Continental makers.”

Mr. Crisp said that it should be mentioned that there had been some difference of opinion, both on the part of the committee and the Council, as to whether there should not be a third intermediate

gauge; also, that in selecting the sizes mentioned by the Chairman, regard had been had to those which were at present in most frequent use, so as not to disturb existing arrangements more than could be helped.

The following Papers were taken as read :—

“Plant Crystals,” by Dr. Aser Poli, of Rome.

“Note on the Rev. G. L. Mills’ Paper on Diatoms in Peruvian Guano,” by Mr. F. Kitton.

“On the Estimation of Aperture in the Microscope,” by the late Mr. Charles Hockin, jun.

The following Instruments, Objects, &c., were exhibited :—

Mr. Ahrens :—New Erecting Binocular Microscope.

Mr. Bolton :—Spawn of Perch.

Mr. Crisp :—(1) Balkwill’s Slide of Foraminifera. (2) Hardy’s Compressorium (see p. 553). (3) Malassez’s Compte-Globules (see p. 559). (4) Tolles’ $\frac{1}{4}$ -inch objective with very tapering front. (5) Tolles’ Camera Lucida (see Vol. III. p. 527).

Mr. Iugpen :—Steinheil’s High-power Achromatic Eye-pieces, $\frac{3}{4}$ -inch, $\frac{1}{2}$ -inch, and $\frac{1}{3}$ -inch.

Prof. Liversidge :—“Silver-Fish” (*Lepisma*) from Sydney, N. S. Wales.

Mr. R. Miller :—Caoutchouc Cement.

Dr. Ralph :—*Vallisneria* from Melbourne.

Dr. G. M. Sternberg :—Photographs of Bacteria (see p. 571).

New Fellows.—The following were elected *Ordinary* Fellows :—
Messrs. William Borrer, jun., William E. Pickels, John Rookledge, Thomas C. Squance, M.B. and M.S., and Prof. Albert H. Tuttle, M.Sc.
(At p. 443 for Henry Pocklington read Christopher Pocklington.)

WALTER W. REEVES,

Assist.-Secretary.

Fig. 1. $\frac{250}{1}$

Fig. 2. $\frac{250}{1}$

Fig. 3. $\frac{400}{1}$

Fig. 3. bis.

Fig. 4.

Fig. 5.

Fig. 6.

Fig. 7. $\frac{10}{1}$

Fig. 8. $\frac{40}{1}$

West Newman & Co. sc.