patient it was over forty to the c.m.m. A predominance of polymorpho-nuclear cells over the lymphocytes was present. Eighteen showed positive cultures. All of these ended fatally, except one. A characteristic organism was found in the spinal fluid in a 81.8 per cent. of the cases. The organism is a bacillus, gram-negative, facultative anaërobe, nonmotile, acid producing, and forms spores.

Rovsing, C. CERVICAL RIBS. [Hospitalstidende, Vol. 62, no. 22.]

The details in four cases of cervical ribs are here reported. Neuralgias, scoliosis of the cervical spine, atrophy, and vasomotor disturbance, and aneurysm in the subclavian artery were among the common symptoms observed. He describes the technical surgical procedures of removal.

Strickler, Jr., F. P. FRACTURE OF CERVICAL VERTEBRÆ. [Kentucky State Med. Assoc., Sept. 22, 1919. J. A. M. A.]

Fractures of the cervical vertebræ constitute from 25 to 36 per cent. of all spinal fractures, and occur most frequently in the male of middle age. Several vertebræ may be fractured and dislocated at one time, but the parts most frequently involved are the body. When this is the case the fracture is usually accompanied by dislocation and compression which produce a rather sharp kyphosis. The spinous and transverse processes are also frequently the site of fracture, but in fractures of this nature there may not be much deformity. The site of these fractures, fracture dislocations and compression fractures is in the larger number of cases found in the lower cervical vertebræ. The nerve symptoms found in fracture and dislocation of the cervical vertebræ are produced by confusion or crushing of the cord at the time of injury, or by persistent compression of the cord by fragments of bone, effusions, blood clots, and later on by callus or the products of inflammatory changes.

If few or no cord symptoms are manifested, either the conservative or operative treatment may be employed, depending on which form of treatment will give the best results in the individual case. The conservative treatment is carried out by immobilizing the spine in plaster-of-Paris jackets, the Calot jacket being the type most frequently used in fractures of the cervical region. If the operative treatment is decided on, one of two operations may be used—the Albee or the Hibbs. The prognosis is guarded. In those cases in which there are marked cord lesions it is very grave, for few, if any, of these patients show improvement. The case usually terminates in death.

Brown, T. G., and Stewart, R. M. FLEXION REFLEX. [Jl. Royal Army Med. Corps., 32, 1919, No. 6.]

The phenomena of flexion reflex have not been studied with the same thoroughness in man as they have in animals. The reason for

this is the difficulty of obtaining suitable environment for observation. These authors have endeavored to carry the teaching of Sherrington further and to apply it to the human subject. They have arrived at the conclusion that under normal conditions the only portion of the receptive field that can be stimulated with effect is the area containing sensory nerve endings in connection with the foot and possibly the leg. By pressing the tendo Achilles against the posterior surface of the tibia the great toe flexes at the interphalangeal joint. At times flexion is obtained by pressure applied to the vastus externus, the vastus internus and the gastrocnemius-soleus muscles. On the other hand, when the cord has been injured the receptive field for effective stimulation corresponds more closely to that of the lower animals. The authors give examples substantiating this claim. They point out that the response in the great toe is usually extensor, but in the case of a person affected with syringo-myelia superficial stimuli applied to the sole of the foot gave rise to a flexion of the great toe, while stimuli applied to the gastrocnemius or to the postero-internal border of the tibia resulted in extension. While the flexion reflex in a normal man is very limited, in a subject suffering from a spinal lesion all or nearly all the muscles of the limb are involved. In distinguishing between the flexion and extensor relaxation it is necessary to recognize that the contraction takes place in the physiological flexors. The immediate reflex phenomena usually take the form of flexion at the hip, knee and ankle, extension of the great toe and abduction of the toes. At times the anterior abdominal muscles contract, leading to a flexion of the lumbar spine. At the same time there is crossed extension in the opposite limb. Rarely there may be crossed flexion. The movements of the two limbs are noted to be accurately reciprocal. At times no movements occur in the stimulated limb, while there is flexion after a prolonged period of latency in the other. They have observed in the case of complete division of the cord a reflex reversal. This took the form of extension in the stimulated limb and flexion in the opposite limb. Successive reflex phenomena of a distinct type were observed. They speak of this reaction as the rebound reflex. Similarly, the rhythmic phenomena which have been described in connection with the lower mammals, have been observed in man after division of the cord. The authors claim that many of the reflexes described by them have not before been noted in man, but they suggest that this is due, not to their infrequent occurrence, but to the fact that they have not been sought. They suggest that some of them may be of diagnostic significance. For example, they point out that the rhythmic alternate movements of the lower limbs in response to stimuli applied to one or both lower limbs are in fact forms of the reflex involved in progression. When they occur, it is usually possible to determine the level of the lesion. But, apart from this possible significance, the observations are of importance, demon-

strating, as they do, the fact that the flexion reflex in man is comparable to that of the lower mammals.

Galant, S. SPINAL REFLEX. [Schweiz Arch. f. Neur. u. Psych. J., Vol. 2, No. 2. J. A. M. A.]

Galant noticed that stroking along the spine of a quite young infant, held with its abdomen on his left palm, elicited a characteristic reflex movement. It is rapid, and resembles the curving of the body of a lizard as it winds its way rapidly through the grass. The infant keeps this lateral curve for a time. He examined 150 young infants and thirty-six idiots between seven and thirty years old. The spine reflex occurs more constantly in infants than the Babinski, and is stronger and persists longer. It can be elicited by stroking with the handle of a hammer or by pricking. The reflex movement is so pronounced that the infant is liable to slip off the supporting hand unless one is careful. It grows weaker as the months pass, and it could not be elicited in one seven months infant. All but one of the 105 infants presented this reflex, but only six of seventeen epileptics, and it was typical only in one in this group.

Walshe, F. M. R. THE GENESIS AND SIGNIFICANCE OF SPASTICITY. [Brain, Vol. XLII, Part I, 1919.]

This paper discusses spasticity and the functional relations of the pyramidal system in their bearings upon the principles of physiology enunciated by Sherrington. He has previously advanced evidence that there are two clinical types of spasticity, as seen in the lower limb, the extended and the flexed. The extended type, wherein the limbs lie fully extended with the feet plantar flexed, is seen in hemiplegia, or in any spastic paralysis confined to the pyramidal system. The flexed type, with flexed position and flexor spasms of great frequency and violence, is seen in severe spinal lesions. The extended type may be identified with "decerebrate rigidity"; the flexed type with the "spinal man." The extended type is not only analogous to "decerebrate rigidity," but is an expression of activity of the proprioceptive system (Sherrington) as a whole. This system induces a tonic reflex, which, unmasked in the decerebrate animal, has been shown to possess the significance of a postural reflex—reflex standing. In man, the rigidity seen in hemiplegia has the same significance. There is an anomaly, however, inasmuch as the upper limb shows an attitude of flexion, the hypertonus being resident in the flexors, not in the extensors, as in the lower limb. This may be traced to the altered functions the upper limb exercises in man and to its complete lack of the locomotor function. Walshe denies the accuracy of Förster's theory of the origin of spasticity, namely, that "the pyramidal tract exercises both excitatory and inhibitory functions," because it disregards the fundamental principals laid down by