under the arms. In the sitting posture, the body become bunched up. There was considerable relaxation of the ligaments of the hip and ankle joints, so that the trunk and lower extremities could be placed in various fantastic positions. Extreme hyperextension of the ankle joint could be made, so that the dorsum of the foot lay along the front of the tibia. There was a looseness of all the joints excepting those of the knee, which were held in flexor contraction to a moderate degree. In the dorsal position the lower extremities were abducted and rotated outwards at the hip joints; flexed at the knees, and the soles of both feet were directed inwards and upwards. When the abducted lower limbs, flexed at the knee, were brought into close adduction, the patient was able to hold them in that position for a short time, showing that there was some power in the adductors, but the limbs would soon fall back into their former position. He was able to move the extremities at all the joints. Locomotion was carried on by a process of rolling over from one side to another. By this means he went from one room to another all over the house. Recently, he had been making attempts to get on his hands and knees.

Muscular system: There was a generalized loss of muscle tone of varying degrees. Voluntary power was generally preserved, more or less. The muscles were small and weak, and had not the power they should have. The lower extremities were the most deeply involved. There was no fibrillation. There was a strictly symmetrical flaccidity; a weakness, without paralysis. The Wassermann test was negative. The urine was normal; the blood picture was one of mild secondary anemia. The superficial reflexes were present, with the exception of the epigastric, the abdominal and the cremasteric. The plantar reflexes were flexor in type; no Gordon nor Oppenheim. The knee jerk and ankle jerks were absent. Sensation to touch was normal, and while there was an apparent defect of pain sensibility, no complete loss was demonstrable.

The patient showed no demonstrable evidences of involvement of the central nervous system. The characteristic feature was the loss of muscle tone, and a diminution of voluntary power. The condition began congenitally. The question of rickets could be raised, but the early age of onset and the absence of all bony deformity seemed to exclude it as a possible cause. The main feature seemed to be a loss of muscular tone, which might possibly be due to a defect in the sensory mechanism of the muscles or in their reciprocal innervation.

A CASE OF DYSTONIA MUSCULORUM PROGRESSIVA. (OPPENHEIM'S NEW DISEASE)

By C. C. Beling, M.D.

The patient was a girl, born December 10, 1897, in New York City, of Hebrew parents, both of them native of the town of Vilna in Russia. The family history was good. Both the parents were living and well. The patient had two brothers and four sisters who were in normal health. Two children, twins, died soon after birth.

The history of this case, which was then given by Dr. Beling in detail, showed that an insidious disease appeared at the age of seven years in this otherwise normal girl. Its chief feature consisted of a progressive, generalized muscular spasm, of a tractile and torsion-like

character. The disease began in the right foot and exhibited itself as a disturbance of gait, in which hypertonia and spasmodic components were evident. These spasms produced peculiar deformities in the trunk and extremities. The movements were almost always slow, rather stereotyped and not so graceful and rhythmic as those of bilateral athetosis. Occasionally, jerky muscular contractions were observed. The muscular spasms were not under the influence of the will. In the early stages, voluntary movements were not seriously disturbed, but at the present time they were strongly interfered with. Complicated motor acts, such as rising, walking, eating, etc., had been profoundly affected. There had been no diminution in the power of the muscles excepting that produced by the contractures. The movements ceased during sleep. The intellect had been entirely unaffected and the emotional state was normal. Neither suggestion nor hypnosis had had any influence; there were no hysterical stigmata.

Dr. Beling said the disease showed a marked similarity to bilateral athetosis, chronic chorea and degenerative tic. The characteristic participation of the facial muscles, the peculiar grimacing, noted by Lewandowsky, the combination with it of signs of spastic diplegia, which was a frequently associated phenomenon, and the enormous influence of emotional processes seen in bilateral athetosis were lacking in this case.

Dr. Smith Ely Jelliffe thought the Society was very fortunate in having had the opportunity of seeing the two cases shown by Dr. Beling; they were rare cases, and he saw no reason to question the diagnosis in either. The first case was undoubtedly one of amyotonia congenita. It was interesting to note the marked hypotonia and the frog-like attitude assumed by the patient, which was a distinct feature of the disease. Spiller had given the first careful pathological description of this disease, although Oppenheim was the first to describe it. Could the disorder possibly be an intrauterine poliomyelitis. Such a point of view would separate the disorder from the Werdnig-Hoffmann type of atrophy, with which the disorder might be confounded.

Dr. Jelliffe said the second case shown by Dr. Beling was also quite classical. It was quite analogous to three patients whom he had had the opportunity of seeing in Prof. Ziehen's Clinic in 1909. These were exact counterparts of this patient. At a recent meeting of the Society, when Dr. Fraenkel read his paper upon Oppenheim's new disease, Dr. Jelliffe said he had referred to a colored boy who was under his observation at the City Hospital, and in whom the symptoms were such that the differential diagnosis between a double athetosis, a pseudobulbar palsy and dystonia musculorum was very difficult. That boy had been shown before this society as an example of Huntington's chorea, which it certainly was not.

The speaker said that through the courtesy of Dr. Flatau, of Breslau, he had the opportunity of showing a number of photographs of a patient suffering from this general type of disorder which well illustrated the conformational phases of the disease. He also showed some photographs of the boy at the City Hospital.

Dr. Jelliffe said that in the discussion of Dr. Fraenkel's paper upon the subject of Oppenheim's dystonia musculorum, he recalled that Dr. Kennedy had said that he regarded these cases of Oppenheim's so-called new disease as being rather ordinary, and as belonging to the tic group. Dr. Jelliffe asked Dr. Kennedy whether he still would maintain his previ-

ously expressed view now that he had seen a patient whose entire course and symptoms agreed so closely with the type which Oppenheim had in view and which he, Dr. Jelliffe, knew from personal experience in both Oppenheim's and Ziehen's Clinics was of the group to which Oppenheim had given the name *dystonia musculorum progressiva*?

Dr. Foster Kennedy said that during the discussion referred to by Dr. Jelliffe he had defined a tic as a purposive movement retained after the purpose for which it had been undertaken had been lost. As to the case shown by Dr. Beling, the speaker said he did not think it belonged to the tic group, according to the definition or description of that term.

Dr. William M. Leszynsky said that when he showed the first case of this disorder before the Neurological Society about seven or eight years ago, the true condition was not recognized. Now, since the attention of the profession had been called to it by Oppenheim and Fraenkel, he supposed that more cases would come to light. It was apparently a form of muscular spasm regarding the pathology of which we were ignorant. Personally, he would not be willing to classify the case just presented as among the tics. It may have shown some evidence of a tic in its onset, but now we had absolute tonicity of the muscles and an absence of the purposeful movements of tic. One characteristic of these cases, which was spoken of by Dr. Beling, was that the spasm disappeared during sleep.

Dr. Richard B. Kruna said that in a disorder of this kind, in which we had tonicity of the muscles, he did not regard the disappearance of the movements during sleep as a very important factor, as we might have a state of hyper- or hypo-tonicity which was really secondary, due to muscle fatigue.

Referring to the case mentioned by Dr. Leszynsky, the speaker said that that patient was now doing farm-work, and was apparently showing signs of improvement.

The President, Dr. Clark, referring to Dr. Beling's case of amyotonia congenita, said that several authors have reported at length upon the occurrence of the flaccid palsies in cerebral diplegics. They cited them as freaks or paradoxical diplegias (Little's type), when a close analysis of many of such supposed cases of agenesis of the pyramidal tracts following in the wake of so-called Little's disease one might find were really cases of amyotonia congenita. At least one needs to scan closely these flaccid types of infantile cerebral diplegia with the aforementioned point in mind.

The same speaker said that, in his Randall's Island service he had seen one case at least which he now knew he should have clinically diagnosed as *dystonia musculorum progressiva* instead of bilateral athetosis. He believed this new disorder would ultimately be proven an organic nervous disease.

A CASE OF TRAUMATIC PSYCHOSIS ASSOCIATED WITH AN OLD DEPRESSED FRACTURE OF THE FRONTAL REGION: OPERATION: RECOVERY

By Charles E. Atwood, M.D. and Alfred S. Taylor, M.D.

Dr. Atwood stated that this patient, who had been shown at a meeting of the Society about a year ago, was again presented for the follow-