wounded bladder were pared and brought together with fine wire sutures, and then the external incision was closed with stronger sutures." The patient entirely recovered from the operation and died suddenly six months later. At the autopsy the bladder was found to be converted into a tube eight inches in length by one to one and a half inches in diameter. After briefly describing the tumor Dr. Neale says "The interest in the case chiefly lies in: (1.) The unusual position of the bladder. (2.) The perfect recovery of that organ after the extensive wound. (3.) The great benefit of the exploration upon the general health and growth of the tumor for a time. (4.) The great advantage of carbolic acid dressings. (5.) The fact that, on account of the universal adhesions, no urine was effused into the peritoneal cavity. In Dr. Eustache's case the bladder was wounded and was sewn up with three catgut sutures including the mucous membrane. On the second day urine was passed freely and without pain per urethram. Perfect recovery followed. Here are nine cases in which the bladder had been wounded during abdominal section, with five recoveries and four deaths. In four of these cases the bladder wound was sewn up and replaced in the abdomen. The sewing material in two of these cases was silk, in one silver wire, and in one catgut. In one case the stitches passed through the abdominal parietes and the walls of the bladder. It should be noted that in all the fatal cases¹ reported by Dr. Thomas all the patients died within thirty hours of the operation, except the case related by Dr. Neale, which seems in my opinion to have recovered, *i. e.*, it is the severity of the operation and the other complications that bring about a fatal result and not the wounding of the bladder. Indeed, the latter seems to heal readily as a rule if cleanly cut and neatly sewed up. The question of the best treatment for a bladder wounded during ovariectomy cannot be set down as invariable. Each case will probably have peculiarities necessitating appropriate methods. But the main facts to be learned are: first, that the ovariectomist must be on the lookout for the bladder in making his incision, and, second, he may feel encouraged that nature will heal the wound if it is carefully sewn up. Of course, as the bladder is a distensible organ, and is necessarily so, a method of union which will leave it in its normal distensible condition is better than one in which the viscus is tied to undistensible structures, but sometimes, as in Dr. Thomas's case, it is unsafe to trust to primary union of an unsupported bladder, and he very wisely avoided the risk of urinary infiltration by uniting the wound as he did.²

A CASE OF OSTEOTOMY.³

BY A. T. CABOT, M. D.

A SPECIMEN from a recent osteotomy of the femur must ever be a rarity; for recovery from this operation is the rule almost without exception, and intercurent

¹ Vide Transactions of the American Gynecological Society for 1881.

² The best article on the treatment of intra-peritoneal wounds of the bladder that I have seen is one by Dr. E. Vincent, Surgeon of the Charity Hospital of Lyons. It is entitled "Plaies pénétrantes intrapéritonéales de la Vessie," and is published in the consecutive numbers of the *Revue de Chirurgie* for June and July, 1881. Dr. Vincent quotes freely from the excellent article by Dr. Bartel's, *Die Trauman der Harnblase*, von Dr. Max Bartels, in Berlin. Arch. für klin. Chirurgie von Langenbeck. Bd. XII, pages 518, 715, 1878.

³ Read before the Surgical Section of the Suffolk District Medical Society, January 7, 1882.

maladies seldom supply the chance to study the results of one of these operations till after the displaced parts are so changed by absorption and by the deposition of new tissue that conclusions as to the exact nature of the original injury are impossible. I have, therefore, thought it worth while to publish a wood-cut of such a specimen, which, being cut from a photograph upon the block, reproduces the appearances very exactly.

The history of the case is as follows:—

Mary T., a colored child six years of age, was sent to me by her parents in consequence of extreme rachitic deformity of the legs. The father was healthy; the mother had suffered from hip disease in her youth, and had recovered with ankylosis and apparently considerable shortening. The older brother of my patient was a well-formed boy of nine or ten. A younger brother, three years of age, however, had had an extreme degree of curvature of the tibiae, which had been corrected by osteoclasia. This was done at so early an age because a marked compensatory deformity of the ankles was developing.

Mary's condition when I examined her was the following: Both femora were considerably bowed outwards and forwards. The left knee was in a marked condition of genu valgum, and the right knee showed a slight degree of the same deformity. Both tibiae were bent sharply forward at the junction of the middle and lower thirds.

The child entered the Children's Hospital in July, 1881, and the lower limbs were straightened by osteoclasia, as will be reported at some future time in a series of these operations. During her stay in the hospital she had whooping-cough mildly. When the legs were strong again the child, though in excellent health apparently, was sent into the country for a few weeks, to avoid a too long continuous stay in the hospital.

In October she reëntered with us, and on October 13th was operated upon for the correction of the deformity of the left knee. I employed Macewen's method in this operation.

The knife was entered on the inner aspect of the knee, about one inch above the articulation. It was thrust immediately down to the bone, and an incision one inch in length was made upwards in the axis of the femur through the whole thickness of the soft parts. Before withdrawing the scalpel a chisel 1.5 cm. in diameter was introduced along it, turned at right angles across the axis of the bone, and the shaft of the femur was divided through about three fourths of its diameter. As the bone at the point operated upon was somewhat broader than the chisel, this had to be withdrawn once or twice and reintroduced before the whole width of the bone was severed. Care was taken, however, never to drive the chisel across the whole transverse diameter of the bone, in order to avoid injury of the periosteum on the outside.

The femur being now three quarters divided, the leg was seized and straightened, which was accomplished with but little exercise of force.

There was almost no bleeding from the wound, which was done up carefully with carbolic gauze, the whole operation having been conducted under spray, and with all antiseptic precautions.

Lastly, the leg was bandaged from the toes to the upper part of the thigh with a plaster bandage, and carefully held in good position till this had set and was firm.

No constitutional disturbance followed this operation. The temperature never rose to 100° F., and after the second day continued perfectly normal. The dressing did not soak through, and was consequently left in place.

November 7th. Three and a half weeks after the operation the plaster bandage and Lister dressing were removed. The wound had wholly healed, and the bone was apparently firmly united, with no perceptible callus. The child was still kept in bed, however, in the fear that the leg was not yet strong enough to bear her weight.

November 13th. She became feverish, and soon developed marked typhoid symptoms, — headache, epistaxis, diarrhœa, tympanites, with a characteristic temperature curve. In spite of every care her strength steadily failed, and she died November 22d, five and a half weeks after the osteotomy. Figure 1 is taken

FIG. 1.

FIG. 2.

from a drawing made in May, Figure 2 from one after death.

A complete autopsy was not allowed, but I was permitted to examine the left leg, and obtained the specimen from which the wood-cut is taken.

FIG. 3.

In this figure (Figure 3) the inner side of the bone upon which the chisel was entered is to the left. At *c* we have the line of the epiphysis. Three fourths of an inch above this is the line of division. A glance at

this shows that while on the outer side of the bone the line of the shaft is pretty well preserved, on the inner side a considerable displacement has occurred. The compact wall of the shaft has been driven down into the cancellated tissue to the point *a*. The tissue in the middle of the shaft, on the other hand, was less resistant than the more densely cancellated tissue below, so that here the centre of the lower fragment is impacted into the upper. A very firm locking of the fragments is the consequence, and this no doubt greatly facilitated the rapid recovery. That there has been but slight reaction in the parts about is shown by the absence of callus. The only true callus formation is seen at *b*, where a little new bone has been thrown out over the free end of the lower fragment; besides this there is only a very thin layer of new bone under the periosteum on the outer side.

The course of this case and the post-mortem appearances furnish strong evidence in favor of this method of operating, which leaves the outer side of the bone intact. If the bone were chiseled from the outer side, when it was straightened, a wedge-shaped gap would be left between the fragments, which would have to be filled with new bone before the healing would be complete. This would, of course, require more time than a method by which the fragments were closely applied to each other and really locked together. This point has been very clearly argued by Macewen in his excellent monograph on the subject.

Another advantage in this method is the immediate stoppage of the hæmorrhage. Most of the bleeding comes from the cut surfaces of the bone. This is stopped when these surfaces are forced together, but allowed to take place when they are held apart. The limited amount of hæmorrhage in this case allowed the wound to entirely heal under the first dressing, thus avoiding the necessity of cutting a window in the plaster or disturbing the wounded parts by a change of dressings.

REPORT ON PROGRESS IN THE TREATMENT OF THORACIC DISEASES.

BY F. I. KNIGHT, M. D.

DIGITALIS IN DISEASES OF THE CHEST.

DR. H. C. WOOD made some very valuable and timely remarks on the use of this drug at a meeting of the Philadelphia County Medical Society.¹ In opening the discussion he referred to the current views in regard to the action of digitalis upon the nervous apparatus of the heart, and claimed for it a peculiar effect upon the heart-muscle.

This influence, which had been fully demonstrated by physiological experiment, and sustained by chemical observation, renders digitalis particularly serviceable in the condition of heart disease in which the increased work required of the heart is greater than the increase of the power, without regard to the particular valve which may be affected. It improves the nutrition of the heart by regulating its contractions and lengthening the diastolic interval, doing away with the rapid, imperfect contractions which interfere with the blood supply of the cardiac muscle. In such cases the nutrition of the heart suffers because it is necessary to have lateral distension of the aorta in order to fill the arteries in the muscular tissue. A little digitalis steadies

¹ Philadelphia Medical Times, vol. xi., No. 355.