in the British *Lancet* that all cases of tetanus, whether of the mild or severe type, should be treated from the beginning with magnesium sulphate; the mild type by subcutaneous injections, the more severe type by rectal injections. It is always well to bear in mind that in using large doses, that is, over 4 c. c., a paralysis of the respiratory center may develop, and means for artificial respiration should be at hand.

THORACIC EMPYEMA*

BY MARTIN D. DELANEY, A.M., M.D.,
Alexandria, Va.

It is an interesting fact that the tree, the most familiar and beautiful of all natural objects, is used as a symbol from the opening to the closing of the story of man's destiny upon this globe. We read of "the tree of knowledge of good and evil" in the green fields of Eden; and "of man's first disobedience and of the fruit of that forbidden tree, which brought sin into the world and death by sin" * * * And in that wonderful prophecy, "The Revelation of Saint John the Divine," we read also of "A New Heaven and a New Earth" in the latter days, and of "the tree of life whose leaves were for the healing of the nations."

Of these leaves may we not claim a few in the ministry of surgery, that almost divine "revelation for the healing of the nations?" The practitioner it ennobles and to the patient it often gives the strength and vigor of "the tree of life." In every application, whether for life or for death, it is a consecrated work and we, its followers, should realize and seek, "in season and out of season," to be worthy of our "high calling."

From the standpoint of the railroad surgeon no subject should receive more attention than pus in the thoracic cavity, or thoracic empyema. When we consider the number of patients with fractured ribs that apply to us for relief, it should open our eyes to the fact that some of these cases are punctured wounds and that infection may be carried into the pleural cavity, causing abscess that must be treated surgically or our patients will surely die.

We who have done surgical work in the region of the pleura, when operating upon the liver or kidney for instance, may have at some of these operations (while dealing with adhesions) accidentally opened the pleural sac in reaching these organs without collapse or serious complications following. If the lung does collapse it will in time expand. Hence, why should we not enter here as boldly as elsewhere?

During recent years I have noticed that the chief attention of surgeons in society meetings has been directed to the abdominal viscera, viz., the appendix, the gall-bladder, gastric ulcer, etc., but papers read on so important a subject as empyema are in the minority. Again reflecting on the untold suffering: the wrecked constitutions and the unsightly chests that present themselves either in our consultation rooms, at their homes, or in the hospitals, for physical examination (after having been subjected to the most serious of surgical operations resulting either from procrastination or ignorance) it seems to me that empyema, or pus, in the pleural cavity should be a subject of more frequent discussion in order that recent thought and methods of treatment could be constantly before us.

All ages are certainly susceptible to this lesion,—the infant, the child, the youth, the middle-aged and the aged. I have seen the infant with a diagnosis of gastroenteritis promptly recover after a thoracotomy. I have seen adults with a diagnosis of "indigestion" likewise recover after proper scientific discrimination and resection of ribs.

Taking too much for granted and not going over a chest carefully often results in the most serious disaster. Why should we wait over eight or ten days if only a few of the symptoms of pleurisy are present? If we are not positive where the physical signs are concerned, it is so easy and so simple a procedure to introduce a needle attached to a hypodermic syringe and withdraw some of the contents. If serum, blood or pus is present, we have clinched our diagnosis and can proceed to treat the case in an intelligent manner. If, on the contrary, our tap has been a "dry" one, we have done no harm, but can rest easily in knowing that we have done our full duty to our patient.

*Read by title, Southern States Association of Railway Surgeons, Auxiliary Southern Medical Association, Memphis, Tenn., Nov. 12-15, 1917.

If serum or blood is found we can aspirate; if pus is obtained, a thoracotomy will save the patient if performed early enough.

I know that many physicians insist upon leaving large quantities of plain serum in the pleural cavity, throwing the responsibility upon Nature to remove it. To me this does not seem reasonable. What assurance have we that pathogenic bacteria will not invade this most excellent culture medium and produce a luxuriant growth, giving rise to a chain of symptoms that are at all times alarming not only to us but to the family and attendants?

The avenues by which the micro-organisms can enter the body are many; in children the most frequent route is by way of the lungs, complicating pneumonia. A pneumonia that does not clear up within two weeks in a child needs a puncture between the ribs.

In railroad cases contused and punctured wounds of the chest, either from foreign bodies or from fractured ends of ribs, give rise to the trouble. Tuberculosis and the exanthematous diseases are the next exciting factors.

The infection may begin in the urethra or other genital organs and ascend to the kidney, producing abscess that may rupture into the pleural sac.

Gall-bladder infection, appendicitis, pyosalpinx or any other region affected with pus-producing organisms, may cause infection in the pleural sac either by continuity, by the lymph channels, or by the blood-stream.

The prompt recognition of the chest complication is imperative and the contents must be evacuated, or (as I have had occasion lately to operate upon persons referred to me with large cavities of a leathery consistency usually with the base in the axilla and apex low down) this sac draining pus and varying in thickness from $\frac{1}{8}$ to $\frac{1}{4}$ of an inch, will be presented, necessitating a resection of nearly all the ribs, leaving a deformed chest and perhaps a sinus draining for months, finally metastasis, the whole body eventually being honeycombed, and suffering and death the result. Hence, why delay? Action brings relief and restoration to health.

Before closing this brief appeal for the fuller consideration of empyema, I would urge my co-laborers of this Association that when this hydra-headed enemy invades the citadel of life, to be on their guard and to remember that, although the fabled hydra possessed a multitude of heads which as soon as one was struck off, two others immediately sprang up, still a Hercules was found to dispatch the monster and rid the earth of such a pestilential foe. As fast as Hercules struck off a head, his charioteer seared the place with a hot iron and soon the hydra was no more, showing in this fable the force of physical and mental strength wisely applied and the adaptation of one of Nature's most powerful elements as a supplementary agent under control.

And, now, why may not we in our labors conquer if we apply strength, courage and all the gracious gifts with which the gods provide us?

POST-OPERATIVE RESIDUAL URINE

BY R. M. HARBIN, M.D., F.A.C.S.,
Rome, Ga.

During the last three years of hospital experience, I have endeavored to maintain through the assistance of colleagues and nurses, a critical observation for the discovery of the presence of residual urine in the bladder in routine cases of abdominal surgery, especially in those who do not complain of the bladder, and as a result of these clinical studies the prevalence of cystitis has been well-nigh eliminated. It is needless to say that the early discovery of the presence of residual urine well-nigh establishes the cure with the adoption of the familiar routine treatment, and we are warranted in making the conclusions:

1. That unrecognized residual urine is a more frequent cause of cystitis than a lack of proper aseptic technique in the use of the catheter, and that the menace of residual urine cystitis lies in cases that do not complain of the bladder and can not be ascertained by palpation.

2. That even when the amount of urine voided is apparently normal the presence of residual urine may be painless without