ventricles are found dilated on autopsy. We found them dilated in our three autopsies, the lateral ventricles in one, and the third in two. In the two in which the third ventricle was dilated the lateral ventricles were also slightly dilated. Some of the other symptoms may also be lessened if we do more frequent lumbar puncture.

Dr. Douglas VanderHoof, Richmond, Va.—Just one or two points as to the differential diagnosis in cases of suspected epidemic encephalitis. There are two acute conditions which must be promptly differentiated. Attention has been drawn especially to the confusion which may arise in cases of encephalitis with an acute onset. I refer particularly to instances of botulism, cases of poisoning after eating canned vegetables—string beans, olives, etc. In cases of botulism the onset is acute. There are eye disturbances, paralysis of the muscles of deglutition, etc., and the differential diagnosis is extremely important.

The second condition is one that arose in a patient whom I saw in consultation and concerning whom there was the possibility of wood alcohol poisoning. That again has to be considered in the differential diagnosis.

The third condition is the one Dr. Gaines mentioned, and that is paresis. Early in 1919 I saw two patients who turned out to be instances of encephalitis. The first patient had an onset three weeks before and presented symptoms apparently like those of paresis associated with Argyll-Robertson pupils. The pupils were unequal; they did not respond to light. This patient was referred to Dr. Tucker as a probable case of paresis. The serological examinations for syphilis were negative, however, and the patient went on to a satisfactory and prompt recovery.

The second patient was seen in consultation at an institution where they take care of nervous and mental cases, and while his pupils were not fixed to light, clinically he was a typical parietic. His onset followed an acute ethmoid infection. I do not know the subsequent course of this second patient for the reason that he left this institution and fell into the hands of a neurologist who insisted, in spite of all arguments we could bring to bear, that it was a case of paresis and he intended to treat it as such, although a blood Wassermann and a spinal fluid Wassermann were negative.

These cases that resemble paresis are particularly interesting. In other cases with acute onset we thought of the possibility of botulism or wood alcohol poisoning.

Dr. Gaines (closing).—I am very glad these points in the differential diagnosis have been taken up in this discussion. They were omitted in the reading of the paper on account of lack of time. I wish to mention, however, one other point in the differential diagnosis which is of some value, particularly to surgeons, and that is that these cases may simulate some acute abdominal condition. Climenko mentioned a case at Mt. Sinai Hospital which was operated on for an acute abdominal condition, which turned out to be a case of encephalitis, and one of my cases was sent into the hospital for an operation for appendicitis. It eventually turned out to be a

case of this disease. Some of the cases I have seen I did not abstract in this paper. One case which was not reported in my paper presented every indication of beginning paresis, such as Dr. VanderHoof mentioned. This case showed remarkable somnolence. He was a carpenter by occupation. He would go to sleep at his work. People would find him asleep on the floor. They awoke him and shortly after, when they went back, they found him again sleeping on the floor, simulating cerebrospinal syphilis rather than actual paresis.

No one said anything in the discussion about the fact that not so many cases of this disease are being seen at the present time. I have not seen a case for several months, and my impression is that the disease is not appearing at the present time, although it may later on.

ERYSIPELAS IN CHILDHOOD*

BY J. H. MASON KNOX, JR., M.D., PH.D.,
Baltimore, Md.

Erysipelas is generally understood to be an inflammation of the skin caused by invasion by a member of the streptococcus group of organisms and characterized by erythema, fever and toxemia. The condition has been recognized from the time of Hippocrates, but many diverse diseases probably were included under this name by early writers. The contagiousness of the malady was pointed out by Lorry, in 1777; and Fehleisen, a century later, isolated and described a micrococcus, the streptococcus erysipelatus, found in the lymph vessels of the skin and subcutaneous tissue, which subsequent study has proved to be the infecting agent. That there is a specific variety of streptococcus alone responsible for this form of infection is now considered doubtful. It is a disease of all ages and produces its greatest ravages at the beginning and end of life, in early infancy and old age.

Erysipelas of the new-born is one of the serious manifestations of pyogenic infection, the organism gaining entrance through the umbilical cord or abrasion of the skin. Because of the feeble resistance of the tissues the inflammation spreads rapidly over large areas. The constitutional symptoms, prostration, high temperature and rapid wasting are all severe and death usually ensues after a brief pe-

*Read in Section on Pediatrics, Southern Medical Association, Fourteenth Annual Meeting, Louisville, Ky., Nov. 15-18, 1920.

riod. We are all familiar with the picture, although more cleanly obstetrical practice has greatly reduced the number of these cases.

Again erysipelas in the aged is a serious disease, having a mortality in those over seventy of more than 25 per cent. The fatal termination is often brought about by myocardial changes. Alcoholics especially exhibit little resistance. It is usually stated, on the other hand, that healthy adults rarely succumb and that the average death rate in general hospitals is about 5 per cent.

This figure does not represent the danger of erysipelas among infants and young children who have passed the period of the new-born. Throughout the early months and years this type of infection is a dangerous one and is attended with a high death rate. It was with the hope of adding a little to our knowledge of the disease in childhood that the present study was undertaken. The cases analyzed, fifty-three in number, were admitted to the wards of the Harriet Lane Home of the Johns Hopkins Hospital over a period of six years. They were of average severity and were selected because the observations upon them were fairly complete. It is believed that they represent the disease as it affects children of the given ages and condition in Baltimore.

Age.—The children ranged from one month to nine years in age. The number of cases at age periods arbitrarily selected were as follows:

1 to 2 months.....	10
2 to 6 months.....	10
6 to 12 months.....	9
12 months to 2 years.....	9
2 to 7 years.....	10
7 to 9 years.....	5

Total..... 53

It will be noted that more than half the cases (29) occurred in the first year, and two-thirds of these (20) in the first six months, and that the incidence of the disease decreases with age. This probably corresponds to an increased resistance to infection acquired in later childhood and adult life.

Sex.—Sex seemed to have no bearing upon the incidence of the malady. Of our cases, twenty-seven were females and twenty-six males.

Race.—Of the fifty-three cases studied, forty-three occurred in white (including many foreign) and ten in colored children. These figures represent roughly the proportion of white and colored children admitted and indicate that each race is equally susceptible to the disease.

Season.—Erysipelas is said to be a disease more prevalent in cold weather. This is certainly true of erysipelas among adults and aged. In our cases there was fairly equal distribution throughout the year, with a smaller number of cases in July, August and September. Apparently in early childhood the season has less influence in the incidence of the disease, which depends more upon the nature of the injury than upon climate or season. Distribution by months of our cases is given in the following table:

January-March.....	15
April-June.....	17
July-September.....	5
October-December.....	16

Total..... 53

It will be noted of the fifty-three cases that forty-eight were fairly evenly distributed between October and July, and that only five occurred during July, August and September.

Nature of Initial Lesion and Mode of Origin.—The nature of the initial lesion and mode of origin was most varied. In practically every case a definite form of injury with subsequent skin infection could be determined. An idea of these lesions can only be understood by a detailed report of the cases as follows:

Abrasion or bruise.....	16
Eczema.....	4
Pin scratch.....	4
Vaccination.....	3
Furunculosis.....	3
Irritation of nares (from discharge).....	3
Irritation of vulva (from discharge).....	3
Irritation of face (from discharge following otitis media).....	2
Umbilical infection (delayed).....	2
Adenitis, suppurative.....	2
Pimples (scratched).....	2
Burn.....	2
Infection of breast.....	1
Infection of foot.....	1
Infection of eye.....	1
Tuberculous sinus.....	1
Splinter.....	1
Irritation from turpentine.....	1
Circumcision.....	1

Total..... 53

It is evident that a great variety of lesions can become the portal of entry for the causal organism and that in all instances the resistance of the integument is lowered by injury before infection. The importance of cleanliness in the care of patients suffering with a discharge from mucous membranes and of antiseptic treatment even of simple injuries in children is a manifest conclusion from this tabulation.

Area of Body Involved.—The site of the original infection is briefly indicated as follows:

Face	10
Scalp	8
Legs	19
Arms	6
Genitalia	4
Trunk	5
Buttocks	1
Total	53

It will be seen that no part of the body is immune and that the site is largely determined by the nature of the injury.

Area Involved.—The area of the body involved in the course of the disease was exceedingly variable and bore but little relationship to the severity of the infection. Some of the more severe cases resulted from a comparatively small skin infection, while in others large areas of the body were affected with but moderate constitutional symptoms. Four general groups can be indicated according to the general area of the body involved:

1. Scalp and face, singly or together, with occasional involvement of trunk or arms	19
2. One or more extremities, with occasional involvement of hip, buttocks and trunk	25
3. Genitalia and buttocks, with occasional involvement of extremities and trunk	6
4. Trunk alone	3
Total	53

The areas of the body most often involved are: one leg, 11 cases; head and neck, 8 cases; two extremities (legs or arms with involvement of trunk), 8.

Incubation Period.—The time which elapsed between the initial injury and the beginning of symptoms was from two to twenty-eight days. In the latter case the period was delayed somewhat owing to injury from abrasion of the skin of the arm from a splint applied to set a broken bone. In fifty-one cases in which the incubation period could be determined with

some definiteness, the average time in which the constitutional symptoms began after injury was 5.8 days.

Range of Temperature.—Nearly all the cases showed considerable elevation of temperature and symptoms of intoxication. In one instance the temperature reached 108°. In but one case did the temperature remain below 100°. In all the cases the temperature curve exhibited an irregular or intermittent character. In thirty-three cases of the total number the temperature reached 104° or over. In nine instances it dropped in the course of the disease to 97°. In other words, the temperature range of the cases was typical of septic infection.

Leukocyte Count.—The leukocytes were counted in forty-nine cases. There was a pronounced leukocytosis in nearly every instance. The highest count was 55,000 in a case complicated by otitis media. There were but two counts below 10,000, both of these in critically ill children, one month of age, who were overwhelmed by the infection. Of the forty-nine cases in which counts were made, twenty-nine had more than 20,000 leukocytes to a cubic millimeter. The average leukocyte count of the forty-nine cases was 23,500. In two instances the counts varied markedly in the same patient. In one the initial count was 18,000 and the second count made a few days later was 40,500. This child recovered. In another case the initial count was 45,300 and the second count 21,600, and the patient died. The importance of the white blood cells in resisting the spread of the infection in the lymph channels has frequently been pointed out and the findings in this series would add further evidence in support of this view.

Duration of Illness.—The illness lasted after the initial symptoms twenty-three days in three cases; one case terminated fatally in three days in a weak syphilitic child in which the symptoms were also complicated by pneumonia and empyema. In two instances complete recovery took place in four days after the initial symptoms appeared. The length of illness could be determined in fifty-one cases and its average duration was 12.1 days. It would seem, therefore, that erysipelas in young children is a particularly acute illness in

which the result is determined in a comparatively brief period.

Cultures.—Cultures from the circulating blood were made in eighteen cases; streptococci isolated in six cases; in twelve the cultures were sterile. In one of these twelve cases staphylococcus aureus was found in an abscess in the involved area. In three additional cases streptococci were cultured from an abscess. Streptococci were found in the spinal fluid in addition to the circulating blood in one instance and in two instances in which the blood gave negative results streptococci were found once in the pleural cavity and once from the deeper layer of the skin. It is felt that unquestionably more complete bacteriological study of the blood in cases of erysipelas with marked constitutional symptoms would disclose a larger incidence of septicemia, as has been shown to be true in typhoid fever, tuberculosis and a number of other serious diseases.

Complications.—As would be expected in an infection of this kind, involvement of other tissues than the skin were frequent and complications were found in many of the cases. The most frequent one was inflammation of the middle ear (otitis media), present in nine cases; abscess with suppuration involving deeper structures than the skin was present in seven cases; general septicemia in six cases. Broncho-pneumonia occurred five times and lobar pneumonia once; empyema, secondary to pneumonia, three times; streptococcus meningitis three times. Cervical adenitis was noted twice and tuberculous ulcer of the ankle, once. Four of the children showed evidences of congenital syphilis.

Results.—Of the fifty-three cases, thirty-eight recovered and fifteen died, a mortality of 28.3 per cent. This indicates without further emphasis the serious character of erysipelas in young children. A brief analysis of the fatal cases along the lines already followed will indicate certain features which are more marked in the severe forms of the disease.

ANALYSIS OF FATAL CASES

Age.—The oldest case in which the disease was fatal was four years; the youngest, one month. Twelve of the fifteen cases occurred under six months, which

shows conclusively the increased danger of erysipelas in the very young. The average age of the fatal cases was 3.6 months.

Sex.—Seven of these patients were girls and eight were boys.

Race.—Nine were white and six black.

It will be noted that in the whole series there were but ten colored children. Of these six died, a mortality of 66 per cent. These numbers are too few for general deduction, but they suggest certainly a lessened resistance to streptococcus infection in the colored infant. This is probably due to a lowered state of nutrition and poorer average hygienic surroundings.

Season.—These fatal cases were distributed throughout the year, two of them only occurring between November and June.

Mode of Infection.—A large variety of lesions may form the portal of entry of the infection as indicated by the following table:

Abrasion of extremities.....	4
Nasal discharge	3
Infected umbilicus	2
Infection of the eye.....	1
Abrasion of the face.....	1
Furunculosis of head	1
Vaginal discharge	1
Vaccination	1
Circumcision	1

Total..... 15

The severity of the disease is not determined by the extent of the initial lesion, but rather by the virulence of the infecting organism which may find entrance through an otherwise insignificant injury. The two cases of infection of the umbilicus probably belong to the group of erysipelas neonatorum, but in each instance the constitutional symptoms commenced at about one month of age after the babies had been discharged well from the obstetrical service. The children are, therefore, included in this series.

Area of Body Involved.—The areas of the body involved in the fatal cases were as follows:

Legs, genitalia and trunk.....	4
Head, face and trunk.....	4
Head and face.....	2
Arm and trunk.....	2
Trunk alone	2
Genitalia and buttocks.....	1

Total..... 15

In most of these cases the tendency of the infection to migrate from area to area was noted.

Incubation Period.—The beginning of symptoms after the initial injury varied from two to twenty-eight days in the fatal cases, the average being 6.6 days, which is a little longer average than that for the whole group.

Leukocyte Count.—The leukocyte count was always high, averaging about 23,000 for all the fatal cases. A fall in leukocytes seemed to indicate lowered resistance.

Duration of Illness.—The average duration of illness was 12.2 days and was terminated often by the presence of complications, particularly septicemia and pneumonia.

Cultures.—Most of the positive cultures were obtained in this series of fatal cases, showing evidences of marked intoxication. Ten of these cases were cultured and in six the blood was positive, and in but two negative. Other positive cultures of streptococci were obtained from abscesses at the site of injury, the spinal fluid and pleural cavity.

Complications.—The number of complications in the fatal cases was of course large. Septicemia occurred six times; pneumonia, six times; meningitis, three times; empyema, three times, and peritonitis, twice. Three of the fatal cases were syphilitic.

Treatment.—A discussion of the treatment would not be helpful. No drugs or external applications were found to be specific. Supporting treatment consisted of diet suitable for age and condition of the patient, and hydrotherapy for pyrexia. Occasional sedatives for restlessness were employed. In several instances applications of non-flexible collodion beyond the advancing edge of the skin infiltration seemed temporarily to check the progress of the disease; at other times it had no effect. Various ointments or lotions were used to allay itching and to make the patient more comfortable. Frequently compresses were found to be more helpful. Serum treatment with anti-streptococcal serum and vaccine treatment were not employed. It is felt as yet that its value has not been demonstrated, and that it may do harm.

The most favorable group of cases are

those in which the erysipelas spreads from a lesion where the infection is under considerable tension, often under a scab which has formed over a small infected area. When this focus is opened, the scab removed and treated with an antiseptic dressing the surrounding area of redness and heat usually quickly recedes and the constitutional symptoms subside. In the treatment therefore of all cases it is important to examine carefully for these areas of infection and when found to treat them vigorously.

SUMMARY

A review of the cases here presented would indicate that erysipelas occurring in young children is an exceedingly serious disease and that its danger is in direct proportion to the youth of the patient. The largest mortality occurs in the first half year of life, even when the cases of so-called erysipelas of the new-born are eliminated.

The incubation period is variable; the average is about five or six days. The symptoms are those of local infection and general constitutional intoxication. Leukocytosis is a prominent feature. A low count would indicate decreased resistance.

Both sexes are equally susceptible. There is probably but little difference in susceptibility between the black and white races. The negro seems to be less resistant to the infection. The disease occurs in children throughout the year, but is more frequent from October to June. The incidence of the disease depends largely upon the frequency of the presence of infection and the resistance of the patient following the initial lesion. Any form of injury which breaks the continuity of the skin may form a portal of entry. The intensity of the course of the illness is somewhat independent of the extent of the area involved and generally an advancing erythema and infiltration mean an increase in the constitutional symptoms.

The duration of the disease may vary from three days to three weeks. The average is about twelve days. A fatal termination is usually the result of a general septicemia, pneumonia or involvement of the meninges or the peritoneum. The mortality rate in the whole series was 28 per cent, including fifteen children over

two years old, in whom the rate was only 6 per cent. Of the twenty cases between two and six months, twelve died, a death rate for that period of 60 per cent.

The disease is best prevented by avoiding injury of the skin and in being scrupulously careful in the treatment of these injuries when they occur, and by the prevention of irritation of the skin through infectious discharges.

DISCUSSION

Dr. O. H. Wilson, Nashville, Tenn.—I have had four cases of erysipelas during the last year. The striking difference from Dr. Knox's description is that they lasted longer, usually over two weeks. Only one of the four died. A little two-months syphilitic with erysipelas involving the entire body recovered, where a perfectly healthy nine-months-old child with only partial involvement died. Some seem to resist the infection better than others.

As to treatment as summed up by Dr. Knox, I agree that nothing seems to do any good. Ichthylol is usually used as a placebo. The most important thing is to keep up nourishment, at any rate the quantity of fluids.

One striking point in my cases has been the freedom from complications, especially subcutaneous suppuration. Edema is quite frequent, but I rarely see suppuration.

Dr. P. F. Barbour, Louisville, Ky.—I went through the contagious wards of the Cook County Hospital during the meeting of the American College of Physicians in Chicago. At that time the professor said to us: "I have exhausted every trick and every treatment and nothing does it any good. They die or get well without any help from us." I have not felt quite so hopeless in my own experience. In the last year I saw four cases in new-born babies and but one died. It died one day after being seen. I have had eight cases of erysipelas recently with no deaths except this one. One agent that I have used seems to me to give good results and that is the erysipelas phylacogen. I have used antistreptococcic serum without any effect on the disease, but I have gotten satisfactory results with the phylacogen. At least my patients have recovered. I have seen the temperature drop and the spread of eruption cease. In view of what Dr. Knox has said, I shall not claim that phylacogen has done it. Some of my friends who have used it have also gotten good results in cases of erysipelas.

Dr. W. McKim Marriott, St. Louis, Mo.—When it comes to the treatment of erysipelas, I can agree with what has been said about the futility of using drugs. My experience with vaccines and antistreptococcic serum is that they do not help. I think there are very few things that do help. Recently Woodyatt mentioned the fact that while ordinary flexible collodion painted around the erysipelatous area was of little value, the non-flexible collodion was distinctly advantageous. Following his suggestion we have used non-flexible collodion and I have really felt pleased with the results. The idea is to compress the deeper skin lymphatics and thus limit the extension of infection. Local applications have very

little effect. The giving of fluids in large amounts is important. It has been our experience that serious sequelae, such as nephritis, seem to be mitigated by having the patient drink two or three quarts of water a day.

Dr. L. R. DeBugs, New Orleans, La.—I should like to ask Dr. Knox two questions. He made blood counts in 18 cases. What I would like to know is at what stage of the disease, after its onset, were the blood cultures made? Were they all made at the same period and when were the six positive findings?

Dr. G. G. Thornton, Lebanon, Ky.—I am surprised at the doctor's opinion regarding the futility of the use of vaccines in erysipelas. I have been either curing my cases or I have been misled. Within the last two months I have had two cases, one six months old, the other three years, and I had, as I almost universally have, good results in the use of antistreptococcic serum with other things. I have been attributing most of my good results to the antistreptococcic serum. I give to young children, 6 months old, 20 mls. (20 c. c.) of antistreptococcic serum, and within 24 hours the erysipelas usually ceases to spread. With this I give them internally fluid extract of echinacea. Sometimes I give in connection with that sulphid of calcium if the case does not get better in 48 hours. In the child three years old that I saw two months ago I gave only 10 mls. (10 c. c.) of the antistreptococcic serum at the first visit, but the erysipelas spread rapidly, so I gave 20 mls. the second day; 20 the third day, and 20 the fourth day, when the case recovered rapidly. In addition to this I made application of a formula that I took from a medical journal a number of years ago that runs about this way:

Phenol	} aa	1 oz.
Acidi picrici		
Acidi boracici		
Thymol		
Glycerine		2 drs.
		4½ oz.
Alcoholis qs. ad.		16 oz.

M.—Apply as directed.

I apply this full strength not necessarily over the erysipelatous area, but just a little ahead of it. In all my cases where it was inclined to spread and the antistreptococcic serum did not stop it within the first 24 hours, the spread seemed to be a little lighter. In the case of an old lady with erysipelas in the lower extremity before the days of antistreptococcic serum, I apparently stopped the spread by the use of a strong solution of carbolic acid. I saturated a cloth with the solution and bound it around the leg above the erysipelatous area and this resulted in a blister. When the disease came to the area circled by the carbolic acid it stopped, notwithstanding that it was the most malignant case of erysipelas I have had occasion to see. Over the erysipelatous area the serum in the blisters that formed was absolutely black under the blebs. The tissue sloughed out in some places nearly to the bone and the old lady, 80 years of age, got well.

Dr. J. J. Waller, Oliver Springs, Tenn.—My experience has been better than those which I have heard. I have been using 50 per cent, sometimes 25 per cent, ichthylol in lanolin, and it has invariably given me good results. In my case I could not get the ichthylol mixture immediately and used a preparation I copied from a medical

journal, and it worked better than the ointment. The formula contained ammonium muriate and tincture of stramonium in water. It is put on a wet cloth and applied to the surface involved. It seemed to do good almost immediately. The formula was as follows:

Muriate ammonia	1 oz.
Tr. robelia fol.	4 drs.
Aq. dist. qs.	3 oz.

M. Sig.—Apply 3 or 4 times a day.

As to the mixture the doctor spoke of, I remember years ago in Chicago the celebrated Nicholas Senn gave a prescription in his clinic that almost invariably gave good results. It was a 50 per cent solution of alcohol in water. Simply plenty of buttermilk applied locally in saturated gauze is best of all.

Dr. Eugene Rosamond, Memphis, Tenn.—Some five or six years ago some one suggested painting the erysipelatous area with pure carbolic acid, then immediately after it had blanched the skin, dissolving it off with its perfect solvent, alcohol. I thought at the time that I had discovered an ideal treatment for erysipelas and was very much interested in this treatment. I had an associate with me, one of the best internists in the city of Memphis, and I saw his cases and he saw mine, and we applied this treatment in a series of nine cases, getting beautiful results. Since that time for some mysterious reason it has failed to work, and I have tried phylacogen and I have tried antistreptococcic serum and I have had the same experience that everybody else has had. It pursues its own sweet way, sticking its scarlet finger out from under the painted area and spreading across the skin as has been described by the other speakers. I want to mention this because it became quite the vogue at one time to use this treatment. Certainly if local applications can stop this infection the use of pure carbolic acid would do so, using it until we get a superficial burn and immediately dissolving it out of the tissues with alcohol.

Dr. Knox (closing).—Dr. DeBuys asked about the cultures. In the very ill on admission they were made at once. In these cases the constitutional symptoms were severe. Many of the positive cultures were obtained in the fatal cases. I do believe, however, that more cases of septicemia occur in erysipelas than have been heretofore demonstrated by routine cultures.

We must be very careful in making deductions as to therapy in a disease of this kind. We all know the diverse and specific therapies that have been suggested for lobar pneumonia. Literally pneumonia has been cured a hundred times and in a hundred different ways, and yet there are many of us who feel that no specific treatment for pneumonia has been found. Erysipelas tends to attack the patient with a lowered resistance. Probably any treatment which would increase the leukocytosis and attack the infecting agent would be helpful. The pathology of the disease shows us that the inflammation is deep down in the lymph channels under the skin and is out of reach of antiseptics that we can put on the skin without destruction of the tissues.

My object in preparing the summary was simply to call attention to the serious nature of erysipelas, especially in very young children. In these patients it is not a disease of a 5 or 10 per cent mortality as many have believed.

THE INTRAPERITONEAL USE OF DEXTROSE AND NORMAL SALT SOLUTION IN THE TREATMENT OF MARASMUS AND THE SEVERE TYPE OF MALNUTRITION*

BY H. LESLIE MOORE, M.D.,
Dallas, Tex.

We are all familiar with the splendid and often spectacular results obtained in acute dehydrated conditions from the administration of fluids intrasinus and intraperitoneally. Some three years ago, at the Dallas Baby Hospital, we began the use of this method in the treatment of certain types of chronic conditions in which there was more or less dehydration—the advanced cases of marasmus and the severe type of malnutrition.

Those of you doing hospital and dispensary work have always on hand feeding cases that refuse to gain in weight regardless of proper nursing and scientific feeding. Some of these cases show a complete digestion and take full feedings; many of them have normal appetites and the stools show a moderate degree of indigestion. Such cases have dropped below a certain level from which they can not climb. The purpose of treatment is to give them a lift and start them on an upward course. This is done by giving daily 250 to 400 c. c. of a 5 per cent dextrose in normal salt solution intraperitoneally until there has been a steady gain in weight over a period of several days. Within a few hours following the first injection the vitality improves and by the third or fourth treatment the stools have become normal in number and appearance. Usually after the second day the loss of weight is checked and by the fourth day they begin to gain. After gaining steadily for a few days treatment is discontinued, and rarely needs repeating. Occasionally the weight comes to a standstill after gaining steadily for a number of days. Under such conditions, if it remains stationary or a loss begins, treatment must again be instituted as before. I do not recall having to give more than the second lift.

*Read in Section on Pediatrics, Southern Medical Association, Fourteenth Annual Meeting, Louisville, Ky., Nov. 15-18, 1920.